The role of solid-state reactivity in the prebiotic selection of nucleobases and formation of peptides

Stolar, Tomislav

Doctoral thesis / Disertacija

2022

Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj: University of Zagreb, Faculty of Science / Sveučilište u Zagrebu, Prirodoslovno-matematički fakultet

Permanent link / Trajna poveznica: https://urn.nsk.hr/urn:nbn:hr:217:985093

Rights / Prava: In copyright/Zaštićeno autorskim pravom.

Download date / Datum preuzimanja: 2025-03-22



Repository / Repozitorij:

Repository of the Faculty of Science - University of Zagreb







University of Zagreb FACULTY OF SCIENCE

Tomislav Stolar

THE ROLE OF SOLID-STATE REACTIVITY IN THE PREBIOTIC SELECTION OF NUCLEOBASES AND THE FORMATION OF PEPTIDES

DOCTORAL DISSERTATION

Supervisor: Dr. Krunoslav Užarević

Zagreb, 2022.



Sveučilište u Zagrebu PRIRODOSLOVNO-MATEMATIČKI FAKULTET

Tomislav Stolar

ULOGA REAKTIVNOSTI U ČVRSTOM STANJU U PREBIOTIČKOM ODABIRU NUKLEOBAZA I STVARANJU PEPTIDA

DOKTORSKI RAD

Mentor: Dr. sc. Krunoslav Užarević

Zagreb, 2022.

Acknowledgements

I am thankful to my wife Jasna Alić Stolar, my mum Snježana, and my brothers Dado, Josip, and Angelo for their support. Also, to Rea for being the best dog ever and doing what only dogs can do. I love you with all my heart.

Thanks to Krunoslav Užarević for being the best possible mentor for me. For giving me space to grow and being alongside me through ups and downs.

Thanks to Ernest Meštrović for supporting me throughout my professional career: for advice, motivation, and access to experts and equipment to work with.

Thanks to José G. Hernández for being a great mentor and friend. José is one of the most talented scientists that I know, and he will do great things for the Colombian and Latin American scientific community.

Thanks to Saša Grubešić for helping and teaching me about analytical chemistry. Saša will soon obtain his PhD degree which he undoubtedly deserves given his expertise.

Thanks to Luca Grisanti for believing in my crazy ideas and working with me. I am looking forward to publishing the results that we obtained.

Thanks to Martin Etter for mentoring me about crystal structure solution from powder X-ray diffraction data and providing me with the software to work with. You helped me tremendously.

Thanks to my scientific brother Stipe Lukin for joining me in many scientific adventures. I enjoyed the time, discussions, and learned a lot from you.

Thanks to my collaborators: Dr. Nikola Cindro (University of Zagreb, Chemistry Department, Croatia), Dr. Ivica Đilović (University of Zagreb, Chemistry Department, Croatia), Dr. Nenad Tomašić (University of Zagreb, Geology Department, Croatia), Dr. Mario Cindrić (Ruđer Bošković Institute, Croatia), Mrs. Lucija Dončević (Ruđer Bošković Institute, Croatia), Dr. Krešimir Molčanov (Ruđer Bošković Institute, Croatia), Dr. Ivor Lončarić (Ruđer Bošković Institute, Croatia), Dr. Krešimir Molčanov (Ruđer Bošković Institute, Croatia), Dr. Ivor Lončarić (Ruđer Bošković Institute, Croatia), Dr. Gregor Mali (National Institute of Chemistry, Slovenia), Dr. Andraž Krajnc (National Institute of Chemistry, Slovenia), Dr. Ben K. D. Pearce (Johns Hopkins University, USA). I hope we will soon publish our results if we already didn't do that.

Thanks to: Dr. N. V. Hud (Georgia Tech, USA), Dr. Jason Dworkin (NASA), Dr. Michael Callahan (Boise State University, USA), Dr. Ramanarayanan Krishnamurthy (The Scripps Research Institute, USA), Prebiotic Chemistry & Early Earth Environments community supported by NASA Astrobiology programme, and participants of Molecular Origins of Life conference organized by Emergence of Life Initiative for the discussion.

Thanks to my PhD committee members: Prof. Mirta Rubčić, Prof. Sanda Rončević, and Prof. Oliver Trapp. I enjoyed the discussion.

Thanks to my best man Mario Špadina for being supportive whenever needed. Thanks to all other friends who helped me in my life.

Contents

ABS	TRACT	IX
SAŽETAKXIII		
PROŠIRENI SAŽETAKXIII		
§ 1.	INTRODUCTION	
§ 2.	LITERATURE REVIEW	
2.1.	Prebiotic selection of nucleobases	5
2.2.	Prebiotic peptide synthesis	8
2.3.	Mechanochemical reactions in prebiotic chemistry	11
§ 3.	EXPERIMENTAL	
3.1.	Molecular recognition of nucleobases	15
3.1.1.	General conditions	15
3.1.2.	. Variable-temperature synchrotron powder X-ray diffraction monitoring	16
3.1.3.	. Crystal structure solution from powder X-ray diffraction data	16
3.1.4.	Other solid-state analytical techniques	
3.2.	Peptide synthesis	
3.2.1.	General conditions	18
3.2.2.	Ion-pairing high-performance liquid chromatography	18
3.2.3.	Mass spectrometry	19
§ 4.	RESULTS AND DISCUSSION	
4.1.	Prebiotic emergence of molecular recognition on a nucleobase level	23
4.1.1.	. 2,6-Diaminopurine and uracil	23
4.1.2.	. 2,6-Diaminopurine and thymine	27
4.1.3.	Adenine and hypoxanthine	
4.1.4.	. Selectivity of base-pairing and other combinations	
4.2.	Mechanochemical prebiotic peptide synthesis	46
4.2.1.	Amino acid oligomerization	46
4.2.2.	. Amino acid and α -hydroxy acid oligomerization	60
§ 5.	CONCLUSION	
§ 6.	REFERENCES	
§ 7.	APPENDIX	XXIII
§ 8.	CURRICULUM VITAE	LXXXV



University of Zagreb Faculty of Science Department of Chemistry

ABSTRACT

THE ROLE OF SOLID-STATE REACTIVITY IN THE PREBIOTIC SELECTION OF NUCLEOBASES AND THE FORMATION OF PEPTIDES

Tomislav Stolar Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia

Within this thesis, solid-state reaction medium and mechanochemical methodologies were explored to unveil weather nucleobases can self-assemble and amino acids can oligomerize under these conditions. X-ray diffraction and differential scanning calorimetry analysis showed that 2,6-diaminopurine (D) selectively self-assembles with uracil (U) and thymine (T), while adenine (A) self-assembles with hypoxanthine (H). Crystal structures of D-U cocrystal hydrate, D-T cocrystal hydrate, A-H cocrystal hydrate, and A-H cocrystal anhydrate were determined from powder X-ray diffraction and single-crystal X-ray diffraction data. Ion-pairing high performance liquid chromatography and mass spectrometry analysis showed that glycine and alanine undergo oligomerization by mechanical activation. Based on the results, it can be concluded that minerals as additives, time, mechanical loading, and temperature are all important parameters for the mechanochemical prebiotic peptide synthesis. In summary, solid-state reactivity of nucleobases and amino acids might have been an important factor in their prebiotic chemical evolution.

(160 pages, 119 figures, 14 tables, 161 references, original in english)

Thesis deposited in Central Chemical Library, Horvatovac 102A, Zagreb, Croatia and National and University Library, Hrvatske bratske zajednice 4, Zagreb, Croatia.

Keywords: amino acids/ mechanochemistry/ molecular recognition/ nucleobases/ peptides/ prebiotic chemistry/ solid-state reactivity

Supervisor: Dr. Krunoslav Užarević, Senior Research Associate

Thesis accepted: 29.06.2022.

Reviewers:

Prof. dr. sc. Mirta Rubčić Prof. dr. sc. Sanda Rončević Prof. dr. Oliver Trapp **Doctoral Thesis**



Sveučilište u Zagrebu Prirodoslovno-matematički fakultet Kemijski odsjek

SAŽETAK

ULOGA REAKTIVNOSTI U ČVRSTOM STANJU U PREBIOTIČKOM ODABIRU NUKLEOBAZA I STVARANJU PEPTIDA

Tomislav Stolar Institut Ruđer Bošković, Bijenička c. 54, 10000 Zagreb, Hrvatska

U ovoj doktorskoj disertaciji koristili smo čvrsto stanje kao reakcijski medij i mehanokemijske metodologije da bismo istražili dolazi li do prepoznavanja nukleobaza i oligomerizacije aminokiselina u tim uvjetima. Analiza difrakcijom rentgenskog zračenja i diferencijalnom pretražnom kalorimetrijom pokazala je selektivno molekulsko prepoznavanje 2,6-diaminopurina (D) s uracilom (U) i timinom (T), kao i adenina (A) s hipoksantinom (H). Riješene su kristalne strukture D-U kokristalnog hidrata, D-T kokristalnog hidrata, A-H kokristalnog hidrata te A-H kokristalnog anhidrata iz podataka difrakcije rentgenskog zračenja. Analiza tekućinskom kromatografijom visoke djelotvornosti i spektrometrijom mase pokazala je da glicin i alanin oligomeriziraju u peptide uslijed mehanokemijske aktivacije. Rezultati su pokazali da su minerali kao aditivi, vrijeme, prenošenje mehaničke energije, i temperatura ključni parametri za mehanokemijsku prebiotičku sintezu peptida. Zaključeno je da je reaktivnost nukleobaza i aminokiselina u čvrstom stanju mogla biti važan faktor u njihovoj prebiotičkoj kemijskoj evoluciji.

(160 stranica, 119 slika, 14 tablica, 161 literaturnih navoda, jezik izvornika: engleski)

Rad je pohranjen u Središnjoj kemijskoj knjižnici, Horvatovac 102a, Zagreb i Nacionalnoj i sveučilišnoj knjižnici, Hrvatske bratske zajednice 4, Zagreb.

Ključne riječi: aminokiseline/ čvrsto stanje/ mehanokemija/ molekulsko prepoznavanje/ nukleobaze/ peptidi/ prebiotička kemija/

Mentor: dr. sc. Krunoslav Užarević., viši zn. sur.

Rad prihvaćen: 29.06.2022.

Ocjenitelji:

prof. dr. sc. Mirta Rubčić
prof. dr. sc. Sanda Rončević
prof. dr. Oliver Trapp

Doktorska disertacija



Sveučilište u Zagrebu Prirodoslovno-matematički fakultet **Kemijski odsjek**

PROŠIRENI SAŽETAK

1. UVOD

Prebiotička kemija je kemija koja prethodi nastanku života [L. E. Orgel, *Crit. Rev. Biochem. Mol. Biol.* **39** (2004) 99–123.]. Kroz kemijsku evoluciju raste kompleksnost na molekulskoj razini te ona prethodi biološkoj evoluciji. S obzirom da je površina Zemlje danas znatno izmijenjena uslijed različitih geoloških procesa tijekom vremena, nema povijesnog dokaza koji bi upućivao na način na koji je život nastao. Kao najizglednije lokacije na kojoj je život mogao nastati smatraju se kopno ili plitki kopneni bazeni [B. Damer, D. Deamer, *Astrobiology* **20** (2020) 429–452.]. Jedna od nepoznanica je razlog iz kojeg su kanonske nukleobaze odabrane i ugrađene u nukleinske kiseline s obzirom da se međusobno ne povezuju Watson-Crick načinom prepoznavanja u vođenim otopinama [D. M. Fialho, T. P. Roche, N. V. Hud, *Chem. Rev.* **120** (2020) 4806–4830.]. Naša hipoteza je da su druge kombinacije nukleobaza preferirane u odnosu na kanonske kombinacije. S druge strane, povezivanje slobodnih aminokiselina u oligomerne lance zahtjeva eliminaciju molekula vode. Iz tog razloga je oligomerizacija aminokiselina do peptida termodinamički nepovoljan proces u vođenom mediju [G. Danger, R. Plasson, R. Pascal, *Chem. Soc. Rev.* **41** (2012) 5416–5429.]. Naša hipoteza je da mehanokemijska aktivacija slobodnih aminokiselina na površinama minerala može omogućiti nastanak peptida.

Doktorska disertacija

2. LITERATURNI PREGLED

2.1. Prebiotički odabir nukleobaza

Nukleobaze (dušične baze) djeluju kao prepoznavajuće jedinice unutar nukleinskih kiselina i njihova komplementarnost je najpoznatiji primjer molekulskog prepoznavanja u prirodi. Pretpostavlja se da je na ranoj Zemlji bio dostupan veliki izbor dušikovih heterocikličkih spojeva koju su mogli poslužiti kao nukleobaze [D. M. Fialho, T. P. Roche, N. V. Hud, *Chem. Rev.* **120** (2020) 4806–4830.]. Argumenti za takvu hipotezu su brojni sintetski eksperimenti koji rezultiraju velikim brojem različitih purinskih i pirimidinskih baza [C. Menor-Salván, M. R. Marín-Yaseli, *Chem. Soc. Rev.* **41** (2012) 5404–5415.]. Također, pronalazak različitih nukleobaza u meteoritima [M. P. Callahan, K. E. Smith, H. J. Cleaves II, J. Ruzicka, J. C. Stern, D. P. Glavin, C. H. House, J. P. Dworkin, *Proc. Natl. Acad. Sci. U.S.A.* **108** (2011) 13995–13998.], kao i njihova sinteza u interstelarnim uvjetima pri niskim temperaturama [Y. Oba, Y. Takano, H. Naraoka, N. Watanabe, A. Kouchi, *Nat. Commun.* **10** (2019) 4413.], snažno upućuju u širok bazen nukleobaza koje su vjerojatno bile prisutne prije u Sunčevom sustavu prije nastanka života. Prema tome, velika je nepoznanica na koji način je došlo do prebiotičkog odabira nukleobaza koje su ugrađene u nukleinske kiseline s obzirom da se kanonske nukleobaze međusobno ne prepoznaju u vođenom mediju.

2.2 Prebiotička sinteza peptida

Smatra se da su peptidi zbog svojih katalitičkih svojstava i simbioze s nukleinskim kiselinama igrali važnu ulogu na ranoj Zemlji [M. Frenkel-Pinter, M. Samanta, G. Ashkenasy, L. J. Leman, *Chem. Rev.* **120** (2020) 4707–4765.]. S obzirom da je u vodenom mediju nastajanje peptidne veze termodinamički nepovoljan proces, prebiotička sinteza peptida u vodi je izazovna. Postoje brojni međusobno oprečni modeli koji objašnjavaju prebiotičku sintezu peptida, međutim oko njih ne postoji konsenzus u znanstvenoj zajednici [T. D. Campbell, R. Febrian, H. E. Kleinschmidt, K. A. Smith, P. J. Bracher, *ACS Omega* **4** (2019) 12745–12752.]. Brojni dokazi upućuju da je aminokiselina glicin bila prisutna na ranoj Zemlji, u vanzemaljskim tijelima [J. R. Cronin, C. B. Moore, *Science* **172** (1971) 1327–1329.], i interstelarnom mediju [S. Ioppolo, G. Fedoseev, K. J. Chuang, H. M. Cuppen, A. R. Clements, M. Jin, R. T. Garrod, D. Qasim, V. Kofman, E. F. van Dishoeck, H. Linnartz, *Nat Astron* **5** (2021)197–205.] prije nastanka života.

2.3. Mehanokemijske reakcije u prebiotičkoj kemiji

Mehanokemija je održiva sintetska metoda koja potiče kemijsku reaktivnost putem mehaničke aktivacije [K. J. Ardila-Fierro, J. G. Hernández, *ChemSusChem* **14** (2021) 2145–2162.]. Neki od izvora mehaničke energije na ranoj Zemlji mogli su uključivati udare meteorite i kometa, tektonske poremećaje, eroziju tla, potrese, i vremenske utjecaje poput vjetra ili kiše. Nedavno je objavljeno nekoliko radova koji pokazuju važnost mehanokemijske reaktivnosti za prebiotičku kemiju. Oni uključuju sintezu prekursora aminokiselina [C. Bolm, R. Mocci, C. Schumacher, M. Turberg, F. Puccetti, J. G. Hernández, *Angew. Chem. Int. Ed.* **57** (2018) 2423–2426.], modifikaciju aminokiselina [J. E. Elsila, N. M. Johnson, D. P. Glavin, J. C. Aponte, J. P. Dworkin, *Meteorit. Planet. Sci.* **56** (2021) 586–600.], i sintezu šećera [S. Lamour, S. Pallman, M. Hass, O. Trapp, *Life* **9** (2019) 52.; M. Hass, S. Lamour, S. B. Christ, O. Trapp, *Commun. Chem.* **3** (2020) 140.].

3. EKPERIMENTALNI DIO

3.1. Molekulsko prepoznavanje nukleobaza

3.1.1. Opći dio

Nukleobaze koje su korištene u ovome radu kupljene su od proizvođača Carbosynth. Mehanokemijske reakcije provedene su koristeći InSolido Technologies kuglični vibracijski mlin i posudice od stakloplastike, zapremnine 14 mL. Dvije čelične kuglice mase 1,4 g i promjera 7 mm korištene su za mljevenje. Reakcije su provedene pri frekvenciji od 30 Hz u trajanju od 2 sata. Za provođenje tekućinski potpomognutih reakcija mljevenja, 50 μL H₂O je ukupno dodano u 200 mg mase reakcijske smjese. Mehanokemijska reakcija u kriogenim uvjetima provedena je u istim uvjetima. Posudice s reakcijskom smjesom su uronjene u tekući dušik prije početka mljevenja te je tijekom reakcije osiguran kontinuirani dotok tekućeg dušika. Reakcije u vodi provedene su miješajući ekvimolarne količine reaktanata (ukupna masa reakcijske smjese iznosila je 100 mg) u 3 mL vode pri sobnoj temperaturi. Miješanje uz pomoć magnetske miješalice provedeno je u okruglim tikvicama volumena 10 mL, a trajalo je nekoliko dana. Jedinični kristali spojeva **2** i **10** dobiveni su hlapljenjem metanolnih otopina pripravljenih spojeva **1** i **4** (10 mg u 5 mL metanola). Hidrotermalna sinteza **9** provedena je zagrijavajući

ekvimolarne količine adenina (A) i hipoksantina (H) na 120 °C preko noći u kvarcnoj epruveti koja podnosi visoki tlak. Za analizu nuklearnom magnetskom rezonancijom u čvrstom stanju, uzorak je pripremljen suhim mljevenjem ekvimolarne količine A i H u čeličnoj posudici koristeći dvije čelične kuglice mase 1,4 g te naknadnim grijanjem tako priređene smjese do 300 °C u eksperimentu diferencijalne pretražne kalorimetrije (posudica od platine, 5 °C/min brzina zagrijavanja, atmosfera dušika). Difraktogrami praškastih uzoraka, koji su korišteni za rješavanje strukture odgovarajućih uzoraka, prikupljeni su u rotirajućoj staklenoj kapilari na laboratorijskom difraktometru (Stoe Stadi-P, $\lambda = 1,54059$ Å). Podaci difrakcije na jediničnom kristalu prikupljeni su na Oxford Diffraction Xcalibur Nova R instrumentu ($\lambda = 1,54719$ Å) opremljenom s Oxford CryoJet instrument koji koristi hlađenje tekućim dušikom. Programski paket CrysAlis PRO12 [CrysAlisPro, Rigaku Oxford Diffraction, 2018, version: 1.171.39.46, Rigaku Corporation, Oxford, UK] je korišten za redukciju podataka i numeričku absorpcijsku korekciju. Strukture iz podataka na monokristalu su određene i utočnjene pomoću SHELXL [G. M. Sheldrick, *Acta Cryst.* **A71** (2015) 3–8.]. U **2**, kristalizacijska molekula vode (O3) je u neredu kroz dvije pozicije (označene s A i B) te okupancijama od 63 % i 37 %.

3.1.2. Difrakcija sinkrotronskog rentgenskog zračenja na polikristalnom uzorku pri promijenjivim temperaturama

Mjerenja su provedena na P02.1 beamline na PETRA III, DESY pri čemu su korištene staklene kapilare promjera 0,5 mm. Uzorci su grijani pomoću Oxford Cryostream sustava u temperaturnom području 300-500 K. Udaljenost detektora i uzorka je iznosila 2010 mm. Korišteno je rentgensko zračenje od 0.20741 Å i difrakcijska slika je prikupljena na vrtećoj kapilari. Svaki uzorak je termalno uravnotežen na 300 K te je grijan brzinom od 5 K/min. Difrakcijske slike su prikupljene u razmaku od 10 K. Nakon što je uzorak postigao željenu temperaturu, termalno je uravnotežen, te je potom prikupljena difrakcijska slika.

3.1.3. Rješavanje kristalnih struktura iz difrakcijskih podataka na polikristalnom uzorku

Nakon indeksiranja, ćelija koja je bila najbolja kandidat je korištena za Pawley fit [G. S. Pawley, *J. Appl. Crystallogr.* **14** (1981) 357.]. Struktura je rješena metodom kaljenja krutog tijela u direktnom prostoru. Konačno utočnjavanje po Rietveldu [H. M. Rietveld, *J. Appl. Crystallogr.* **2** (1969) 65–71.] je napravljeno koristeći geometrijska ograničenja za duljine veza

i iznose kuteva, te planarnih ograničenja po potrebi. Svi računi su napravljeni u programu Topas [TOPAS, version 4.2; Bruker-AXS: Karlsruhe, Germany, 2009.].

3.1.4. Ostale analitičke tehnike u čvrstom stanju

ATR-FTIR spektri su snimljeni pomoću spektrometra PerkinElmer SpectrumTwo opremljenog s dijamantnom ćelijom u rasponu 4000–400 cm⁻¹. Difraktogrami praha su prikupljeni na PanAlytical Aeris difraktometru (CuKα zračenje, Ni filter) u Bragg-Bretano geometriji, pri čemu je uzorak bio nanesen na pločicu od silicija. Eksperimenti diferencijalne pretražne kalorimetrije provedeni su na Simultaneous Thermal Analyzer (STA) 6000 (PerkinElmer, Inc.) u platinskim posudicama te brzinom zagrijavanja od 5 °C/min u atmosferi dušika protoka 20 mL/min.

3.2. Sinteza peptida

3.2.1. Opći dio

Glicin, 2,5-diketopiperazin, TiO₂, SiO₂, i Montmorilonit K10 kupljeni su od Sigma-Aldrich. Gly₂–Gly₄ i L-Ala-L-Ala kupljeni su od Bachem. L-Ala kupljena je od Carbosynth. Mica (MICA-V-4) kupljena je od SPI. Svi mehanokemijski eksperimenti provedeni su koristeći InSolido Technologies vibracijski kuglični mlin. Za eksperimente s TiO₂, reakcije su provedene pri frekvenciji od 30 Hz u trajanju od 18 h. Ukupna masa reakcijske smjese iznosila je oko 400 mg (0,8 mmol Gly i 4 mmol TiO₂). Korištene su 14 mL posudice od čelika specifično dizajnirane za provođenje mehanokemijskih reakcija pri povišenoj temperaturi uz tri čelične kuglice mase 1 g i promjera 6 mm za mljevenja. Za eksperimente sa SiO₂, reakcije su provedene pri različitim frekvencijama i vremenima mljevenja. Ukupna masa reakcijske smjese iznosila je 500 mg (1,33 mmol Gly i 6,66 mmol SiO₂). Korištene su čelične posudice volumena 28 mL specifično dizajnirane za provođenje mehanokemijskih reakcija pri povišenoj temperaturi te dvije čelične kuglice mase 1,4 g i promjera 7 mm za mljevenje.

3.2.2. Tekućinska kromatografija visoke djelotvornosti

Oligopeptidi su analizirani tekućinskom kromatografijom visoke djelotvornosti na Agilent 1290 Infinity instrumentu koji je bio opremljen Phenomenex Luna C18 kolonom (250 mm dužine, 4,6 mm promjer, i 3 µm veličina čestica). Mobilna faza je bila vodena otopina 50 mM KH₂PO₄ i 7,2 mM natrijev heksansulfonat. pH je podešen na 2,5 dodatkom 85 % H₃PO₄ (HPLC čistoća). Mobilna faza je korištena izokratno s protokom od 1 mL/min. Uzorak je injektiran u volumenu od 1 μL, a vrijeme metode je bilo 20 min. Kolona je uravnotežena na 30 °C. Analiza se temeljila na UV-VIS detektoru pri absorbanciji od 195 nm. 2,5 mg Gly, 2,5-diketopiperazina, Gly₂, Gly₃, i Gly₄ prebačeno je u 1,5 mL HPLC vialu i napunjeno s mobilnom fazom. Viale su protresene na IKA Vortex u trajanju od 1 min kako bi se pospješilo otapanje. Za analizu uzoraka nakon provedenih reakcija, oko 40 mg svakog uzorka prebačeno je u Falcon epruvetu i napunjeno s 1,5 mL mobilne faze. Svaka epruveta je protresena na IKA Vortex u trajanju od 1 min kako bi se pospješilo otapanje oligopeptida. Nakon toga, epruvete su centrifugirane pri 15000 rpm u trajanju od 5 min. Naposljetku, 0,8 mL supernatanta prebačeno je u staklenu vialu te je analizirano pomoću HPLC.

3.2.3. Spektrometrija mase

Oligopeptidi su analizirani sa spregnutim sustavom tekućinske kromatografije koristeći Agilent 1290 Infinity instrument spojen na Q-TOF MS (Model 6545) instrument opremljen s Waters Acquity BEH HILIC amidnom kolonom (duljine 150 mm, promjera 2,1 mm i veličine čestica 1.7 µm). Mobilna faza (MF) je bila otapalo A: ACN i B: 5 mM HCOONH₄ pri pH 3,0 podešeno s HCOOH. Korišteno je izokratno eluiranje kroz 2 min (10 % MF B) nakone čega je uslijedilo gradijentno eluiranje (2-20 min, 10 % MF B do 50 % MF B). Protok je bio 0,4 mL/min. Volumen injektiranog uzorka iznosio je 3 µL, ukupno vrijeme metode bilo je 25 min te je kolona ekvilibrirana na 30 °C. Korišten je dvostruki AJS ESI izvor iona u pozitivnom (+) ESI modalitetu. Parametri temperature plina (300 °C), plina za sušenje (6 L/min), nebulizatora (60 psi), kapilarne struje (3000 V), fragmentora (100 V) i skimmera (65 V) optimizirani su koristeći standard glicina. Q-TOF MS je korišten u spektralnom masenom području 40-1700 m/z te je korišteno akvizicijsko vrijeme od 1,4 spektra/s (akvizicijsko vrijeme = 714 ms/spektru). Napravljeno je unutarnje utočnjavanje mase i kalibracija instrumenta Q-TOF prije analiziranja uzoraka. Instrument je kalibriran u visokom rezolucijskom modalitetu (4 GHz, 0-1700 m/z). Uzorci koji su korišteni za analizu tekućinskom kromatografijom visoke djelotvornosti su razrijeđeni u acetonitrilu (100 µL uzorka i 900 µL acetonitrila).

4. REZULTATI I RASPRAVA

4.1. Molekulsko prepoznavanje nukleobaza

4.1.1. 2,6-Diaminopurin i uracil

Provedeni su eksperimenti prepoznavanja nukleobaza u različitim uvjetima prebiotičke zastupljenosti vode. Za opisivanje uvjeta velikog suviška vode korištene su reakcije miješanja krute reakcijske smjese u zasićenoj vodenoj otopini. Za opisivanje uvjeta katalitičke prisutnosti vode korištene su tekućinski potpomognute reakcije mljevenja. Za opisivanje uvjeta odsustva vode korištene su reakcije grijanja u čvrstome stanju. Ovisno o reakcijskim uvjetima, 2,6-diaminopurin i uracil tvore kokristalni hidrat ili bezvodnu formu kokristala. Tehnikom difrakcije rentgenskog zračenja na jediničnom kristalu određena je kristalna struktura hidratnog kokristala 2,6-diaminopurina i uracila.

4.1.2. 2,6-Diaminopurin i timin

Ovisno o reakcijskim uvjetima, 2,6-diaminopurin i timin tvore kokristalni hidrat ili bezvodnu formu kokristala. Tehnikom difrakcije rentgenskog zračenja na jediničnom kristalu određena je kristalna struktura hidratnog kokristala 2,6-diaminopurina i timina. Tehnikom difrakcije rentgenskog zračenja na polikristalnom uzorku određena je kristalna struktura bezvodne forme kokristala 2,6-diaminopurina i timina.

4.1.3. Adenin i hipoksantin

Ovisno u reakcijskim uvjetima, adenin i hipoksantin tvore kokristalni hidrat ili bezvodnu formu kokristala. Tehnikom difrakcije rentgenskog zračenja na polikristalnom uzorku određene su kristalne strukture kokristalnog hidrata i bezvodne forme kokristala adenina i hipoksantina. Kao komplementara metoda difrakciji na polikristalnom uzorku korištena je nuklearna magnetska rezonancija u čvrstom stanju uz pomoć koje je utvrđeno u kojoj je tautomernoj formi prisutan hipoksantin.

4.1.4. Selektivnost molekulskog prepoznavanja

Rezultati su pokazali da u svim navedenim reakcijskim uvjetima dolazi do selektivnog prepoznavanja tri kombinacije nukleobaza: 2,6-diaminopurina i uracila, 2,6-diaminopurina i timina, te adenina i hipoksantina. Selektivnost u prepoznavanju je utvrđena u prisustvu kompetirajuće nukleobaze.

4.2. Prebiotička sinteza peptida

4.2.1 Oligomerizacija aminokiselina

Za prebiotičku sintezu peptida korištene su mehanokemijske reakcije u čvrstom stanju. Kao reaktanti su korišteni aminokiselina glicin i alanin. Reakcije su provedene uz prisustvo mineralnih aditiva. Reakcije su provedene u uvjetima različitog vremena trajanja reakcije, različite frekvencije mljevenja te različite temperature mljevenja. Rezultati su pokazali da povećanje temperature mljevenja uvelike pospješuje nastanje dužih oligomera glicina uz povećanje udjela diketopiperazina kao primarnog produkta reakcijske smjese. Tako je primjerice najveći udio ukupnih linearnih oligomera iznosio 10,2 % u uvjetima suhog mljevenja glicina uz TiO₂, frekvenciju od 30 Hz, reakcijsko vrijeme od 18 sati te temperaturu mljevenja od 100 °C. Pokazano je da povećanje duljine mljevenja povoljno utječe na uspješnost oligomerizacije glicina. Isto tako, ključnim se pokazala frekvencija mljevenja s obzirom da se veći unos mehaničke energije u reakcijski sustav očituje većim udjelom nastanka lineranih oligomera glicina. Međutim, čak i pri niskim frekvencijama mljevenja utvrđena je prisutnost nastalih peptida.

4.2.2. Oligomerizacija aminokiselina i α-hidroksilnih kiselina

Na temelju eksperimentalnih podataka utvrđeno je da je najjednostavniji oblik α -hidroksilne kiseline, glikolna kiselina, oligomerizira do poliesterskog oligomera duljine lanca četiri glikolne podjedinice. Također, zajedničkim mljevenjem glikolne kiseline i glicina nastaju njihovi heterooligomeri koji se sastoje od peptidnih i esterskih veza u oligomernom lancu.

5. ZAKLJUČAK

Rezultati istraživanja provedenih u okviru ove disertacije ukazuju da se neki od obrazaca molekulskog prepoznavanja koji su danas prisutni u biokemiji također očituju i na razini nukleobaza u prebiotičkim uvjetima. Molekulsko prepoznavanje 2,6-diaminopurina i uracila, 2,6-diaminopurina i timina, te adenina i hipoksantina je selektivno i odvija se pri različitim prebiotičkim reakcijskim uvjetima. S obzirom da mnogi prebiotički sintetski putevi rezultiraju smjesom nukleobaza, za očekivati je da su različite nukleobaze bile prisutne u smjesi na ranoj Zemlji prije nastanka života. Tijekom ranih faza kemijske evolucije, nukleobaze s

komplementarnim supramolekulskim interakcijama mogle su biti odabrane zbog svojih termodinamičkih i kinetičkih svojstava. Te karakteristike mogle su utjecati na njihovu kovalentnu reaktivnost i ugradnju u nukleinske kiseline.

Što se tiče prebiotičke sinteze peptida, pokazano je da mehanokemija vodi ka njihovoj sintezi. Mehanokemijska aktivacija aminokiselina potaknula je njihovu oligomerizaciju na mineralnim površinama. Pokazano je da je diketopiperazin produktivan intermedijer za sintezu peptida te da mehanokemijska sinteza peptidne veze tolerira prisustvo vlage. Reakcije s Gly₂, Gly₃, i diketopiperazinom su pokazale su kako je mehanokemijska sinteza peptidne veze dinamičan proces sa usklađenim nastajanjem i pucanjem peptidne veze. Zajedničko mljevenje glicina i alanine rezultiralo je nastankom njihovih hetero oligopeptida te je pokazalo da se ostale aminokiseline mogu ugraditi u rastući peptidni lanac. Rezultati ovog istraživanja pokazuju da su reakcijsko vrijeme i unos mehaničke energije važni parametri za prebiotičku mehanokemijsku sintezu peptida koja je u skladu s različitim geološkim i okolišnim uvjetima.



University of Zagreb Faculty of Science Department of Chemistry

§ 1. INTRODUCTION

Prebiotic chemistry is the chemistry that precedes the origin of life (Figure 1).¹ The hypothesis behind its underlying research is that chemical evolution, i.e., organization of simple matter into a more complex one, could have served as a transition to biological evolution, e.g., reproduction, variation, and selection of living systems over time.² Due to the alteration of Earth's surface by geological processes, origin of life "cannot be discovered, it has to be re-invented".³ Among the proposed sites where life on Earth could have originated, hydrothermal vents⁴ and land-based pools⁵ are the most popular.



Figure 1. Artistic depiction of origin of life research that includes prebiotic chemistry. Downloaded from: https://centerforchemicalevolution.com/mural-origins-life-0.

Forming important chemical connections and linking monomers of organic building blocks into corresponding biological polymers requires the formation of covalent bonds through condensation reactions where water molecules are eliminated. Hence, their formation is thermodynamically unfavorable in water media.⁶ This has led to the growing realization that, although water is essential for life, it is not necessarily desirable for the synthesis of life's

Doctoral Thesis

building blocks.⁷ To connect the conditions required for their synthesis with the plausible early Earth environment, wet/dry cycles have emerged as the most used experimental approach in the field of prebiotic chemistry (Figure 2).⁸ In the dry phase, chemical synthesis and the formation of covalent bonds is enabled whereas, in the wet phase, combinatorial selection through compartmentalization is operational. The idea behind using wet/dry cycles is that they closely resemble hydration (due to precipitation) and dehydration cycles (evaporation due to heating) in land-based pools.



Figure 2. Wet/dry cycling and its central role in the land-based hypothesis for the origin of life on Earth. Reprinted from the reference⁹ under the terms of Creative Commons public use license.

The availability of dry land on an early Earth is often discussed. Although it is estimated to have been only a fraction of today's landmass, it is considered that there was some land available on the emerging continents and near volcanoes.¹⁰ Among the reasons why dry land is thought to have played an important role in prebiotic chemistry is its capability to concentrate organic building blocks and induce their chemical transformations.¹¹

On the other hand, prebiotic chemistry is also a part of astrobiology, which includes interdisciplinary studies on the origin of life in the universe. Research efforts in prebiotic chemistry have been greatly influenced by the fall of Murchison meteorite which was proven to contain extraterrestrial amino acids,¹² nucleobases,¹³ and sugars,¹⁴ among other organic matter.¹⁵ These studies have offered us a glimpse into the organic inventory that was carried to Earth during the time of the late heavy bombardment.¹⁶ Meteorites, comets, and asteroids are solid planetary bodies whose composition is influenced not only by accretion, but also post-accretion events such as thermal metamorphism,¹⁷ aqueous alteration,¹⁸ and hydrous metamorphism.¹⁹ Taken together with millions of years of space weathering,²⁰ these processes present opportunities for the evolution of organic matter in extraterrestrial solid-state reaction environments. Therefore, recent Hayabusa2²¹ and OSIRIS-REx²² asteroid sample return missions have the potential to offer us unprecedented insight into the origin and evolution of organic matter in our solar system.

In this doctoral dissertation, solid-state reaction medium was used with the aim of solving some of the open question in prebiotic chemistry. In this regard, the inability of canonical nucleobases to self-assemble via Watson-Crick hydrogen-bonding in water is a long-standing bottleneck. It makes it unclear as to how did nature know which nucleobases to incorporate into nucleic acids if they were previously unable to base-pair and select between themselves. We hypothesized that canonical nucleobase pairs are not favored, and instead, other nucleobase pairs might form selectively. We worked with a pool of commercially available and plausibly prebiotic nucleobases. To characterize the reaction products, we primarily relied on powder X-ray diffraction (PXRD) and differential scanning calorimetry (DSC). When possible, crystal structures of novel crystalline phases were solved from single-crystal X-ray diffraction (SCXRD) and PXRD data using available software.

Furthermore, linking free amino acids into the corresponding oligomers requires the formation of peptide bonds along with the elimination of water molecules. Hence, oligomerization is thermodynamically unfavorable in water media. We hypothesized that

mechanochemical activation of free amino acids might enable their oligomerization on mineral surfaces. For studies on oligomerization of free amino acids on solid mineral surfaces, we mostly used glycine (Gly). The reason to choose Gly is its presumed availability on early Earth, its presence in meteorites, and the availability of analytical methods to characterize reaction products. We used diverse class of minerals that were available to us. For analysis, we primarily used the ion-pairing high-performance liquid chromatography (IP-HPLC) with ultraviolet–visible spectroscopy (UV-VIS) detection. Furthermore, the mass spectrometry (MS) was used to unambiguously determine the presence of different oligomers. Mechanochemical ball milling reactions were used to investigate the oligomerization of amino acids on mineral surfaces and some of these were performed as thermally controlled neat grinding reactions.

Some of the results presented in this thesis have been published while others are still in the publishing process.²³

§ 2. LITERATURE REVIEW

2.1. Prebiotic selection of nucleobases

All extant life utilizes DNA for storage and inheritance of genetic information. Besides contributing to the stability of the DNA double helix, specific hydrogen-bonding interactions between nucleobases act as a basis for molecular recognition. From the perspective of the origin of life, there are many puzzling questions regarding the abiotic origin of nucleic acids.^{24–28} Among them is the assumption that, on the prebiotic Earth, there was likely a pool of structurally similar nitrogen-bearing heterocyclic compounds that could have acted as nucleobases (Figure 3).^{29,30} Such a hypothesis is corroborated by many synthetic experiments that produce other purines and pyrimidines besides the canonical nucleobases.^{31–35} Furthermore, the discovery of the presence of extra-terrestrial nucleobases in carbonaceous meteorites,^{13,36} as well as nucleobase synthesis in interstellar ice analogues,^{37,38} strongly points to a pool of nucleobases that may have been present throughout the solar system. Interestingly, canonical nucleobases are unable to base-pair and select between themselves.^{39,40} Canonical nucleobases compete for hydrogen-bonding with water molecules, and vertical stacking is energetically favoured compared to horizontal hydrogen-bonding.^{41,42} This has led to the idea that other base-pairing combinations might have preceded canonical base-pairing that we observe today.39,43



Figure 3. Chemical space of molecules that could have acted as nucleobases on an early Earth. Reprinted with permission from the reference.³⁰ Copyright 2020 American Chemical Society.

As an example, 2,6-diaminopurine (D) could have been a competitor to adenine (A) because of its ability to form three Watson-Crick hydrogen bonds with both uracil (U) and thymine (T), while adenine (A) can only form two (Figure 4a and 4b). In fact, there are examples from contemporary biology showing that certain viruses use D instead of A as its DNA nucleobase.^{44,45} Recently, more examples and detailed mechanistic understanding have been found of how D replaces A in viral genomes.⁴⁶⁻⁴⁸ Additionally, a ribozyme that catalyses template-directed ligation of two RNA strands with 3',5'-phosphodiester linkages and is

7

composed of only D and U nucleotides, can be obtained in vitro.⁴⁹ Moreover, F. H. C. Crick and L. Orgel have argued that an earlier version of the genetic code may have consisted of only A and hypoxanthine (H) as its bases.⁵⁰ Their arguments were based on the relative ease of the prebiotic synthesis of A whilst H could have been produced from A by deamination. Importantly, such purine-only nucleic acid would not violate the principle of continuity in its plausible evolution to the extant nucleic acids.⁵¹ Their hypothesis has been brought up to the light by reports showing that synthetic DNA duplexes with only purine-purine base pairs,^{52,53} including A and H paired via Watson-Crick hydrogen-bonding,⁵⁴ have comparable stability with the extant DNA. Nevertheless, H is biologically relevant as its nucleoside inosine is commonly found in transfer RNAs (tRNAs) where it is involved in codon-anticodon wobble base-pairing (Figure 4c).^{55,56} Collectively, there is a large body of evidence for the presence of diverse nucleobases in prebiotic times. However, what kind of selection pressures operated between them remain a mystery.



Figure 4. Canonical nucleobases are labeled with blue color, whereas three nucleobases that have been detected in meteorites and are present in contemporary biology, are labeled with red color: a) canonical Watson-Crick base-pairing, b) nucleobases capable of forming three Watson-Crick hydrogen bonds, and c) nucleobases involved in codon anticodon wobble base-pairing.

2.2. Prebiotic peptide synthesis

Among many open questions in the field of prebiotic chemistry, the formation of the peptide bond is one of the most important ones. In an aqueous environment, difficulties arise when condensing free amino acids due to thermodynamic and kinetic unfavourability of the peptide bond formation.⁵⁷ On one hand, multiple evidence shows that amino acids such as glycine were present on Earth,⁵⁸ in extraterrestrial bodies,^{59,60} and in interstellar medium before life emerged.⁶¹ On the other hand, peptides are an important class of biological polymers with catalytic properties and are hypothesized to be included in early molecular symbiosis with nucleic acids (Figure 5).58

Notably, reports are suggesting the presence of extraterrestrial peptides in meteorites.⁶² Prebiotic peptide bond synthesis strategies include thermal condensation,^{63–65} different wet/dry cycling approaches,^{66–68} salt-induced peptide bond formation,⁶⁹ condensation in liquid SO₂,⁷⁰ catalysis by carbonyl sulfide,⁷¹ α -aminonitrile ligation in water,⁷² mineral catalysis,⁷³ hypervelocity impact studies,^{74,75} and high-pressure shearing (Figure 6).⁷⁶

In the search for energy needed to drive prebiotic formation of peptides,⁷⁷ the use of mechanical energy appears as a promising solid-state alternative that avoids the limitations related to solvation, solubility, and diffusion.⁶ In relation, it is known that the primordial Earth underwent a heavy meteoritic bombardment,16 which likely delivered diverse pool of extraterrestrial amino acids in ppm concentrations.^{78,79} Moreover, extraterrestrial impactors or terrestrial lithospheric activity could have provided mechanical and thermal energy to drive formation of peptides. As an example, studies simulating impact events in aqueous solutions of amino acids,⁷⁴ and those mimicking natural comet impacts under cryogenic conditions have revealed that glycine and alanine oligomerize due to the shock impact.⁸⁰ However, mostly linear dipeptides and tripeptides are formed in the process. In addition, simulations of glycine subjected to shear stress have shown that formation of linear diglycine could occur under 10 GPa compressive loads at RT, thus indicating the possibility for impact-based and tectonic shearing prebiotic scenarios.76 On the other hand, extreme mechanical stress can also cause peptide cleavage as demonstrated in hypervelocity impacts (mechanical loading of 2.4-5.8 km/s).⁸¹ This might explain the relatively short peptides that are formed under harsh mechanical conditions.



Figure 5. Peptides are hypothesized to have played an important role in the origin of life. Reprinted with permission from the reference.⁵⁸ Copyright 2020 American Chemical Society.



Figure 6. Some of the synthetic strategies towards prebiotic peptides. Reprinted with permission from the reference⁸² under the terms of Creative Commons public use license.

2.3. Mechanochemical reactions in prebiotic chemistry

The use of mechanical action to run chemical reactions has been used since antiquity⁸³ and modern synthesis technique is called mechanochemistry.^{84–87} It is highly efficient in inducing chemical reactivity in the absence of bulk solvents and kinetically accelerates solid-state reactions by overcoming slow solid-state diffusion.⁸⁸ Vibratory ball milling, planetary milling, and twin-screw extrusion are among the most used types of milling instruments (Figure 7). The field of mechanochemistry has experienced a significant growth in recent years due to its compliance with the principles of green chemistry and sustainability.⁸⁹ It has since been used in virtually all areas of chemical science due to different reactivity and selectivity when compared to the solution-based synthetic approaches.^{90–113}



Figure 7. Three main types of mechanochemical instruments: a) vibratory ball milling, b) planetary ball milling, c) twin-screw extrusion. Adapted with permission from the reference.⁸⁹ Copyright 2021 John Wiley & Sons, Inc.

More than two decades of intense research resulted in many innovations within the field. In liquid-assisted grinding (LAG) methodology, small amounts of liquids are added to solid reaction mixtures.^{114,115} It can have a strong reaction accelerating effect or can direct reaction courses into specific pathways. Importantly, several examples of temperature monitoring showed that bulk temperature of the reaction mixture rises during the milling process.^{116–118} This is most likely due to collisions and friction that occur in milling reactions. Notably, temperature oscillations can also be due to endergonic or exergonic nature of specific reactions, albeit to a lesser extent.¹¹⁹ Using sub-ambient temperature ball milling, it was established that temperature differences of 5 °C affect both yields and enantioselectivity on several reactions such as asymmetric aldol condensation catalyzed by (S)-proline, enantioselective multicomponent reactions between an aldehyde, an amine, and an alkyne, as well as Diels Alder reactions.¹²⁰

Furthermore, experimental setup was devised that used modified jars, thermocouple, and a ring heater element managed through a proportional-integral-derivative (PID) controller device (Figure 8).¹²¹ It was shown that temperature during milling can be precisely controlled up to 250 °C and with a deviation of ± 0.2 to ± 5 °C, depending on the milling reactor material. The utility and generality of such an experimental setup was showed for the synthesis of C-C, C-N bonds, and porous metal-organic frameworks. Notably, except for faster conversion rates to products, milling temperature had a profound effect on the selectivity as demonstrated on obtaining four different derivatives containing amido and urea groups in the mechanochemical reaction between 4-nitrobenzoyl azide and benzene-1,4-diamine just by varying milling temperature parameter. Furthermore, other experimental setups for high-temperature ball milling have been reported, for example using a heating gun that blows hot-air to the milling jar. First examples of such setup were described for the mechanochemical palladium-catalyzed Suzuki-Miyaura cross-coupling of insoluble aryl halides¹²² as well as for the synthesis of magnesium-based carbon nucleophiles, i.e., Grignard reagents, in air.¹²³ Overall, literature reports show that high temperature ball milling allows access to products that are characterized by high activation energies.



Figure 8. Thermally controlled ball milling setup that uses modified jars, thermocouple, and a ring heater element managed through a PID controller device. Reprinted with permission from the reference.¹²¹ Copyright 2019 American Chemical Society.

Additionally, ball milling photocatalytic setups have been developed which merge light and mechanical activation.^{124,125} Importantly, a growing number of mechanochemical reactions

include gaseous reactants and show great promise for the discovery of new reactivity of solidgas interfaces.^{126,127} Furthermore, *in situ* techniques for monitoring mechanochemical reaction courses have been established and are based mostly on powder X-ray diffraction^{128–130} and Raman spectroscopy.^{131–133} These offer mechanistic insights and means of kinetic analysis. Recently, mechanoredox reactions have been discovered and offer new avenues for redox chemistry.^{134,135} Finally, in accelerated ageing methodology, solids are initially mixed and are left to react on a scale longer than typical laboratory experiments.¹³⁶ The latter might be compatible with long-time prebiotic reactions.

Some of the primordial sources of energy that are coupled with prebiotic solid-state syntheses of various classes of molecules are mechanical and thermal energy. Mechanical energy on the prebiotic Earth could have been provided by impacts, tectonics, erosion, earthquakes, and weathering (Figure 9), whereas geothermal activity could have provided thermal energy. Recently, there have been several literature examples of rich solid-state reactivity relevant to prebiotic chemistry. The first reported use of mechanical activation in a prebiotic chemistry context has enabled the activation of highly stable iron cyanide complexes.¹³⁷ Under ball milling conditions with SiO₂, the authors observed mechanochemical activation of potassium ferricyanide(III) *via in situ* production of HCN that was subsequently incorporated into the corresponding α -aminonitriles. Furthermore, hydrolysis of α -aminonitriles into amino amides successfully showed that an impact scenario on an early Earth might offer a route towards amino acids. These conclusions were further supported by a different study where grinding processing of iron meteorites resulted in the formation of amino acids.¹³⁸ This confirmed that mechanical action might have offered a mechanism for the delivery of amino acids by iron meteorites.
§ 2. Literature review



Figure 9. Prebiotic sources of mechanical energy might have included impacts, erosion, earthquakes, tectonics, and weathering. Adapted with permission from the reference.²³ Copyright 2021 John Wiley & Sons, Inc.

In another study, mechanical energy was used to drive the synthesis of monosaccharides.¹³⁹ A wide range of linear and branched monosaccharides including ribose was obtained after ball milling formaldehyde, glycolaldehyde, and glyceraldehyde together with mineral additives as catalysts. These results showed that the formose reaction route to sugars is operational and is kinetically accelerated under solid-state reaction conditions devoid of aqueous media. The role of minerals in mechanochemical formose reaction was further investigated in a separate study.¹⁴⁰ In contrast to the aqueous formose reaction pathway that yields decomposition products, the solid-state pathway enriches monosaccharides over time. The selectivity towards individual sugars was demonstrated to be fine-tuned using a particular mineral catalyst. Moreover, the mechanochemical formose pathway was shown to be independent of the atmospheric conditions and proceeds under cryogenic conditions, thus offering both terrestrial and extraterrestrial prebiotic routes towards sugars. In addition, mechanochemistry was used in a unified route towards purine and pyrimidine nucleosides.¹⁴¹ The formation of critical N-glycosidic bond was achieved by either solid-state grinding and subsequent heating of N-isoxazolylurea, ribose, and boric acid at 95 °C, or by drying their aqueous solutions at the same temperature. Conversion of intermediate N-isoxazolyl-N'ribosyl-ureas to pyrimidine nucleosides was achieved by the catalytic action of iron-sulfur minerals.

Overall, the use of mechanochemical reactions in prebiotic chemistry is still in its beginnings, likely influenced by until recently dominant hypothesis that life originated in hydrothermal vents (deep in the ocean).

§ 3. EXPERIMENTAL

3.1. Molecular recognition of nucleobases

3.1.1. General conditions

Nucleobases used in this study were purchased from Carbosynth. Ball milling experiments were conducted using InSolido Technologies vibratory ball mill and in 14 mL poly(methyl methacrylate) (PMMA) containers. Two 1.4 g stainless steel (SS) balls (7 mm in diameter) were used as milling media, reactions were conducted at 30 Hz frequency and for the duration of 2h. When performing liquid-assisted grinding (LAG) reactions, 50 μ L of H₂O was added to overall 200 mg of the reaction mixture. Cryogenic ball milling experiments were conducted using the same abovementioned conditions. Jars were immersed in liquid nitrogen before the start of the experiment and liquid nitrogen was poured continuously for the duration of the reaction. Slurry reactions in water were performed by mixing equimolar amounts of reactants (overall mass of the reaction mixture was 100 mg) in 3 mL of H₂O at room temperature. Mixing was achieved in 10 mL round bottom flask using magnetic stirrer for several days. To grow single crystals of 2 and 10, 10 mg of 1 and 4 was dissolved in 5 mL of methanol and left for solvent to evaporate at room temperature. Hydrothermal synthesis of 9 was performed by heating equimolar amounts of A and H at 120 °C overnight in a pressure-resistant quartz tube. For solid-state nuclear magnetic resonance (ssNMR) analysis, we used sample that was prepared by neat grinding (NG) A and H equimolar mixture in stainless steel jars with two 1.4 g stainless steel balls and subsequent heating to 300 °C. High-resolution PXRD patterns which were used for crystal structure solution were collected in a glass capillary on a Stoe Stadi-P laboratory diffractometer equipped with copper radiation ($\lambda = 1.54059$ Å). Single crystal measurements were performed on an Oxford Diffraction X calibur Nova R ($\lambda = 1.54719$ Å) equipped with an Oxford Instruments CryoJet liquid nitrogen cooling device. Program package CrysAlis PRO12 [CrysAlisPro, Rigaku Oxford Diffraction, 2018, version: 1.171.39.46, Rigaku Corporation, Oxford, UK] was used for data reduction and numerical absorption correction. The structures were solved and refined with SHELXL [G. M. Sheldrick, Acta Cryst. A71 (2015) 3-8.]. Models were refined using the full-matrix least squares refinement; all non-hydrogen atoms were

refined anisotropically. Hydrogen atoms were located in a difference Fourier map and refined as a mixture of free restrained and riding entities. In **2**, crystallization water molecule (O3) is disordered over two positions (designated as A and B) with respective occupancies of 63 % and 37 %.

3.1.2. Variable-temperature synchrotron powder X-ray diffraction monitoring

In situ synchrotron PXRD monitoring of solid-state heating/cooling reactions was performed at P02.1 beamline at PETRA III, DESY. Glass capillaries of 0.5 mm width were filled with ball milled solid samples and were heated/cooled using Oxford Cryostream system. The latter was operating in 100-500 K temperature range. Distance between the sample and the detector was 2010 mm. X-rays of 0.20741 Å wavelength were used and X-ray diffraction pattern was collected on a spinning capillary. Each sample was thermally equilibrated at 300 K and was heated/cooled using a 5 K/min ramp. For each experiment, X-ray diffraction patterns were collected in 10 K steps. After reaching the desired temperature set-point, sample was thermally equilibrated, and X-ray diffraction pattern was collected. Exposure time for each collected diffraction pattern was 60 s for dark image followed by 60 s for real image. To obtain 2D plots, data were processed in MATLAB using an in-house developed script.

3.1.3. Crystal structure solution from powder X-ray diffraction data

For crystal structure solution from PXRD data, all calculations were performed using the program Topas.¹⁴²

The workflow for structure solution

1) The scan was loaded in xy file format.

2) Data indexing was performed by manually selecting 15-20 reflections. This can also be done in an automatic fashion. However, since successful unit cell hits are highly dependent on the selected reflections, we generally relied on our own assessment.

3) Unit cells with the best values were selected for Pawley refinement. The objective of Pawley refinement was to identify the best unit cell candidate. If suggested unit cells had the same parameters but different symmetry, we generally proceeded with the space group having higher

symmetry. The main criterium for identifying the unit cell was the assessment of the number of molecules in the asymmetric unit. When this was done, the data was exported as INP file.

4) To prepare the complete INP file, the coordinates of the crystallizing molecule(s) were extracted from the database CIFs containing the same molecule(s) (alternatively, this can be done by creating a Z-matrix). This was performed by exporting the target molecule(s) as pdb file in Mercury software. The objective of assembling the INP file is to place the target molecule(s) in the unit cell after which simulated annealing in direct space was used to find the position, orientation, and meaningful interactions of the crystallizing molecule(s).

5) The general criteria to recognize that simulated annealing process has finished was when R_{wp} remained constant for a significant period. This was either hours or days. However, the final decision was made after visually inspecting the CIF file.

An example of INP file ready for simulated annealing is given in Figure 7.1.

Rietveld refinement of crystal structures

1) The background and unit cell parameters were refined first.

2) To refine the atom positions, we used geometry restraints (using the penalty function) on bond distances and angles as well as planarity restraints. The values of bond distances and angles were extracted from CIF file in Mercury.

3) For validation of the structural parameters, we used Mogul Geometry Check option in Mercury. If there were discrepancies with respect to the database mean values, the database mean values of each specific bond distance or angle were further used as starting points for the next iteration of the refinement.

4) Final CIFs were deposited to Cambridge Crystallographic Data Centre (CCDC).

An example of INP file that was used in final Rietveld refinement is given in Figures 7.2. and 7.3.

3.1.4. Other solid-state analytical techniques

ATR-FTIR spectra were recorded on PerkinElmer SpectrumTwo spectrometer equipped with a diamond cell in a 4000-400 cm⁻¹ range. PXRD patterns were collected on PanAlytical Aeris diffractometer (CuK α , Ni filter) in Bragg-Bretano geometry, with the sample mounted on zero background silicon holder. DSC experiments were performed using Simultaneous Thermal Analyzer (STA) 6000 (PerkinElmer, Inc.) in platinum crucibles at 5 K/min heating rate under 20 mL/min nitrogen gas flow.

3.2. Peptide synthesis

3.2.1. General conditions

Glycine, 2,5-diketopiperazine (DKP), TiO₂, SiO₂, and Montmorillonite K10 were purchased from Sigma-Aldrich. Gly₂–Gly₄ and L-Ala-L-Ala were purchased from Bachem. L-Ala and glycolic acid were purchased from Carbosynth. Mica (MICA-V-4) grade was purchased from SPI. All milling experiments were conducted in InSolido Technologies vibratory ball mill. For the reactions with TiO₂, reactions were conducted at 30 Hz frequency and all reactions were performed for the duration of 18 h. Overall mass of reaction mixture was always around 400 mg (0.8 mmol Gly and 4 mmol TiO₂). There, we used 14 mL hardened stainless steel (SS) containers specifically designed for high temperature milling reactions and three 1 g SS balls (6 mm in diameter) as milling media. 25 μ L of H₂O was used as the liquid additive in LAG reactions. For the reactions mixture was always 500 mg (1.33 mmol Gly and 6.66 mmol SiO₂). There, we used 28 mL hardened stainless steel (SS) containers specifically designed for high temperature steel (SS) containers specifically designed for high temperature was always 500 mg (1.33 mmol Gly and 6.66 mmol SiO₂). There, we used 28 mL hardened stainless steel (SS) containers specifically designed for high temperature milling reactions and three 1 metation time. Overall mass of reaction mixture was always 500 mg (1.33 mmol Gly and 6.66 mmol SiO₂). There, we used 28 mL hardened stainless steel (SS) containers specifically designed for high temperature milling reactions and two 1.4 g SS balls (7 mm in diameter) as milling media.

3.2.2. Ion-pairing high-performance liquid chromatography

Oligopeptides were analyzed by Ion-Pairing High-Performance Liquid Chromatography (IP-HPLC) on Agilent 1290 Infinity instrument equipped with a Phenomenex Luna C18 column (250 mm long, 4.6 mm diameter and 3 μ m particle size). The mobile phase was an aqueous solution of 50 mM KH₂PO₄ and 7.2 mM sodium hexanesulfonate adjusted to pH 2.5 by the addition of 85 % H₃PO₄ (HPLC grade). The mobile phase was used isocratically with 1 mL/min flow rate. Sample injection volume of 1 μ L was used, the total method time was 20 min, and

the column was equilibrated at 30 °C. The instrument was equipped with multi-wavelength UV-VIS detector set to record absorbance at 195, 200, 210 and 214 nm (with reference set at 360 nm). The analysis relied exclusively on the data collected at 195 nm. Before the first separation experiment, the column was equilibrated with the mobile phase for 24 h.

Individual standards preparation

2.5 mg of Gly, DKP, Gly₂, Gly₃, and Gly₄ was transferred to 1.5 mL HPLC vial and filled with mobile phase. Vials were stirred with IKA Vortex for 1 min to aid with dissolution.

Sample preparation

Around 40 mg of each milled sample was transferred into Falcon tube and filled with 1.5 mL of mobile phase. Each tube was stirred with IKA Vortex for 1 min to aid with dissolution of oligopeptides. After that, all tubes were centrifugated at 15000 rpm for 5 min. Then, 0.8 mL of the supernatant phase was transferred into a glass vial and analyzed by HPLC.

Calculation

Quantification of oligopeptides was based on measurement (for Gly-Gly₄) and extrapolation (for Gly₅-Gly₁₁) of molar responses. The standards were accurately weighed and analyzed. Obtained peak areas were divided by molar concentration to get molar response factor. The molar response ratio was plotted against oligomer length. Linear relationship was confirmed with obtained correlation coefficient of 0.9959. The relationship was extrapolated to glycine oligomers up to Gly₁₁. The obtained calibration results closely correlated with data reported previously.⁸² From the obtained results (molar concentration of Gly oligomers in samples) the total yield percentage was calculated as the ratio of reaction yield to theoretical yield.

3.2.3. Mass spectrometry

LC/MS (Q-TOF)

Oligopeptides were analyzed by liquid chromatography (LC) on an Agilent 1290 Infinity interfaced to Quadrupole time-of-flight (Q-TOF) mass spectrometry (MS) (Model 6545) instrument equipped with a Waters Acquity BEH HILIC amide column (150 mm long, 2.1 mm diameter and 1.7 μ m particle size). The mobile phase (MP) consisted of solvent A: ACN and B: 5 mM HCOONH₄ at pH 3.0 adjusted with HCOOH. Elution was started with isocratic condition for 2 min (10 % MP B) followed by gradient elution (2-20 min, 10 % MP B to 50 %

MP B). The flow was set to 0.4 mL/min. Sample injection volume of 3 μ L was used, the total method time was 25 min, and the column was equilibrated at 30 °C. A dual AJS ESI ion source was operated in positive (+) ESI mode. The gas temperature (300 °C), drying gas (6 l/min), nebulizer (60 psi), capillary voltage (3000 V), fragmentor (100 V) and skimmer (65 V) parameters were optimized using glycine. The Q-TOF MS was operated in spectral mass range set at 40-1700 m/z and an acquisition rate of 1.4 spectra/s (acquisition time = 714 ms/spectrum). Internal mass tuning and calibration for the Q-TOF was preformed prior sample analysis. The instrument was calibrated in high-resolution mode (4 GHz, low mass range, 0-1700 m/z). A continuous two-point mass (121.050873 m/z and 922.009798 m/z) reference mass correction is further applied during data acquisition using reference solution.

Sample preparation

Samples prepared for IP-HPLC was subsequently diluted with acetonitrile (100 μ L of sample and 900 μ L of ACN) to match the starting mobile phase composition.

Calculation

Glycine and glycine oligomers were detected as the m/z equivalent to the $[M+H]^+$ ion of each analyte. From the obtained TIC chromatogram, extracted ion chromatograms were generated in m/z window of 3 units ($[M+H]^+$ of each oligomer to $[M+H]^+ + 3$). Obtained masses ($[M+H]^+$) were compared to theoretical masses and the difference is expressed in ppm. Glycine oligomers were confirmed with error lower than 3 ppm for the $[M+H]^+$ ions. Additionally, the isotopic distribution was verified for highest detected oligomer (Gly₁₀) in the sample where Gly₃ was milled with TiO₂ at 130 °C, with obtained error less than 2 ppm for three most prominent isotopes.

Direct MS (Q-TOF)

For selected samples direct analysis technique was chosen due to presumed lower solubility of alanine oligomers and simplicity. Samples of L-Ala milled with TiO₂ at 130 °C as well as Gly+L-Ala milled with TiO₂ at 100 °C were injected using Agilent 1290 Infinity instrument without the column. The samples were transferred into Q-TOF MS (Model 6545) instrument with mobile phase (10mM HCOONH₄ at pH 2.3 adjusted with HCOOH/acetonitrile, 50/50v/v) used only as a transfer media, with flow rate of 0.4mL/min. A dual AJS ESI ion source was operated in positive (+) ESI mode. The gas temperature (300 °C), drying gas (61/min), nebulizer

(60 psi), capillary voltage (3000 V), fragmentor (100 V) and skimmer (65 V) parameters were used. The Q-TOF MS was tuned and calibrated in extended dynamic range (2GHz) in 3200 m/z mass range mode. Spectral acquisition range was set from 50-950 m/z with an acquisition rate of 1.5 spectra/s (acquisition time =666 ms/spectrum). A continuous two-point mass (121.050873 m/z and 922.009798 m/z) reference mass correction was further applied during the data acquisition using reference solution.

Glycine and alanine oligomers were detected from the obtained accurate mass spectra (TIC) of the samples as the m/z equivalent to the $[M+H]^+$ ion of each analyte. From the obtained TIC chromatogram $[M+H]^+$ obtained exact masses ($[M+H]^+$) were compared to theoretical masses and the differences were expressed in ppm. All found oligomers were confirmed with error lower than 4 ppm for the $[M+H]^+$ ions.

LC/MS (Triple quadrupole)

Sample preparation

Around 40 mg of each milled sample was added to Falcon tube and filled with 1.5 mL of mobile phase. Each tube was stirred with IKA Vortex for 1 min. After that, all tubes were transferred to ultrasonic bath heated up to 35 °C and sonicated for 15 min. Finally, the content of each tube was filtrated thought 0.2μ M pore size regenerated cellulose filter, transferred into autosampler vial and analyzed by Triple quadrupole LC/MS.

Oligomers were analyzed by hydrophilic interaction liquid chromatography (HILIC-HPLC) on Agilent 1290 Infinity interfaced to Triple quadrupole MS (Model 6470B) instrument equipped with Waters Acquity BEH HILIC amide column (150 mm long, 2.1 mm diameter and 1.7 μ m particle size). The MF consisted of solvent A: 90 % ACN+10 % 200 mM HCOONH₄ at pH 3.0 and B: 20mM HCOONH₄ at pH 3.0. Elution was gradient, starting with 90 % MP A at 0 min to 80 % MP A at 10 min. The flow was set to 0.8 mL/min. Sample injection volume of 0.25 μ L was used and the total method time was 16 min. The column was equilibrated at 30 °C. An AJS ESI ion source was operated in positive (+) ESI mode. The gas temperature (300 °C), gas flow (7 L/min), nebulizer (45 psi), capillary voltage (3500 V), fragmentor voltage (25 V), sheet gas temperature (400 V) and sheet gas flow (11 L/min). The Triple quadrupole MS was operated in SIM mode (monitoring [M+H]⁺ ions of glycine and glycine oligomers to Gly₆) with Dwell time set to 50 ms.

MALDI TOF

For matrix-assisted laser desorption/ionization time-of-flight (MALDI TOF) analysis, we used 4800 Plus MALDI TOF/TOF Analzyer (Applied Biosystems, USA). Samples were dissolved in CHCA matrix. Ultra-pure water (Pliva d.d., Zagreb), acetonitrile HPLC grade (Merck), trifluoroacetic acid (Merck), and CHCA matrix (Merck) were used. CHCA matrix (5 mg/mL) was prepared in acetonitrile solution and 0.1 % trifluoroacetic acid (50/50, v/v) mixed with ultra-pure water. 2 mg of sample was centrifuged for 10 min in aqueous solution 1 mL, after which 2 μ L of sample was mixed with 8 μ L of CHCA matrix. 1 μ L of the latter was transferred to MALDI holder. Samples were calibrated with near external calibration method with standards of azithromycin and domicile peaks resulting from the matrix ionization. Analysis of the samples was performed in both positive and negative modes.

§ 4. RESULTS AND DISCUSSION

4.1. Prebiotic emergence of molecular recognition on a nucleobase level

It was recently shown that model nucleobases, having a methyl group attached at the glycosidic nitrogen atom, selectively self-assemble from multi-component nucleobase solid mixtures.¹⁴³ Although equivalent experiments with canonical nucleobases failed to yield base-pairing, the idea that solid-state can provide an alternative reaction medium for chemical selection based on self-assembly served as a motivation for this study.

In this section, the objective was to explore molecular recognition properties in a pool of plausibly prebiotic nucleobases. These included canonical nucleobases: adenine (A), thymine (T), uracil (U), guanine (G), cytosine (C) and those nitrogen heterocycles that have been detected in meteorites or are present in contemporary biology; 2,6-diaminopurine (D), hypoxanthine (H), and xanthine (X). We studied their interactions in reaction conditions that cover all extremes of prebiotic water activity and elucidated their selective self-assembly processes. We used slurry reactions in water to mimic prebiotic bulk water activity, LAG with water to mimic prebiotic conditions of catalytic water presence, and solid-state heating reactions to mimic dry prebiotic environments. Variable-temperature synchrotron powder X-ray diffraction (VT-SPXRD) monitoring enabled insights into the nucleobase cocrystal formation mechanisms. For example, cocrystallization from an amorphous matrix or cocrystal hydrate transformations upon wet/dry cycling.

4.1.1. 2,6-Diaminopurine and uracil

As a starting experiment, an equimolar amount of powdered D and U was milled together for 2 h. PXRD analysis revealed that the reaction product **1** was poorly crystalline (Figure 10). After ageing **1** in water vapor, as well as by LAG with the addition of catalytic amount of water, **1** is converted to a highly crystalline phase **2** that is absent of diffraction peaks corresponding to reactants (Figure 10). Furthermore, **2** can also be obtained by slurring equimolar amounts of untreated reactants in water at room temperature for six days (Figure 10).



Figure 10. PXRD patterns for the D and U system.

We grew single crystals of **2** and elucidated its crystal structure by SCXRD (Figure 7.4 and Table 7.1).^a **2** is a cocrystal channel hydrate and crystallizes in monoclinic C2/c space group. The asymmetric unit of **2** contains D and U assembled by three intermolecular Watson-Crick hydrogen bonds together with disordered water molecules (Figure 11a). Water content of **2** was determined by TGA and resulted in 5 % (Figure 7.5). Inside the crystal, D and U molecules form hydrogen-bonded layers (Figure 11b). Disordered water molecules fill structural voids and form channels along crystallographic *c*-axis (Figure 11c). Van der Waals interactions are acting between hydrogen-bonded layers that are distanced at 3.4 Å (Figure 12). Furthermore, disordered water molecules serve as vertical bridges between horizontal hydrogen-bonded layers by being involved in van der Waals interactions with D and U molecules within layers.

^a The crystal structure of **2** was solved by collaborating co-author Dr. Krešimir Molčanov (Ruđer Bošković Institute).



Figure 11. (a) The asymmetric unit of 2 contains D and U molecules assembled by Watson-Crick hydrogen-bonding as well as disordered water molecules (b) Hydrogen-bonded layers inside the crystal along crystallographic *b*-axis. (c) View from the crystallographic *c*-axis showing disordered water molecules that lie in the channel.



Figure 12. Distance in Å between hydrogen bonded layers of 2.

The heating effect of wet/dry cycling on D and U base-pair was monitored in situ by VT-SPXRD (Figure 13). At around 400 K, the formation of new Bragg reflections started to appear and reached maximum intensities at around 430 K. After 430 K, this new phase **3** which is presumably a D:U cocrystal anhydrate, is the only one present in the reaction mixture until the end of monitoring at 490 K. We attempted to solve the crystal structure of **3** from PXRD



data but were not successful. The best Pawley fit was achieved by using $P\overline{1}$ space group but simulated annealing failed to provide any reasonable structural model (Figure 14).

Figure 13. a) In situ SPXRD monitoring of heating **2** using synchrotron radiation ($\lambda = 0.20741$ Å). To obtain 2D plot, individual PXRD patterns were stacked upon each other. Note: some Bragg reflections are shifting towards lower 2θ values during heating due to thermal expansion of the unit cell.



Figure 14. The best Pawley fit for **3** was achieved by using $P\overline{1}$ space group (a = 10.22 Å, b = 7.20 Å, c = 16.58 Å, $\alpha = 112.00^{\circ}$, $\beta = 82.30^{\circ}$, $\gamma = 96.33^{\circ}$, V = 1144.02 Å³). Simulated annealing failed to provide any reasonable structural model.

To test whether the formation of D:U cocrystal hydrate proceeds under cryogenic conditions, cryogenic ball milling experiment was conducted. Rietveld refinement confirmed that after LAG an equimolar amount of D and U with H_2O for 30 min in the 80-100 K temperature range, D and U self-assembled as **2** (Figure 15).



Figure 15. Rietveld refinement of LAG an equimolar amount of D and U with H_2O for 30 min in the 80-100 K temperature range. The resulting product is mainly composed out of 2.

4.1.2. 2,6-Diaminopurine and thymine

After neat grinding an equimolar amount of powdered D and T, the resulting solid mixture was analyzed by PXRD and revealed that the reaction product **4** was poorly crystalline (Figure 16). After ageing **4** in water vapor, highly crystalline phase **5** absent of diffraction peaks corresponding to reactants was obtained and is presumably D:T cocrystal hydrate (Figure 16). Crystalline phase **5** can also be obtained by LAG equimolar amounts of D and T with the catalytic addition of water as well as by slurry mixing equimolar amounts of D and T in water for six days at room temperature.



Figure 16. PXRD patterns for the D and T system.

Attempts to grow single crystals of **5** from aqueous solution failed and resulted in single crystals of D hydrate that is known from the literature.¹⁴⁴ On the other hand, after dissolving **4** in methanol, we obtained needle crystals and solved the crystal structure using SCXRD data (Figure 7.6 and Table 7.2).^b This showed that D and T crystallized as a 1:2 stoichiometric cocrystal hydrate **10** in a monoclinic *Pc* space group. The asymmetric unit of **10** contains four symmetrically non-equivalent T and water molecules as well as two symmetrically non-equivalent D molecules (Figure 17). Crystal packing is characterized by hydrogen-bonded layers where each D molecule is connected to two distinct T molecules by three intermolecular Watson-Crick hydrogen bonds as well as to a third T molecule by R_2^2 (9) hydrogen bonding

^b The crystal structure of **10** was solved by collaborating co-author Prof. Ivica Đilović (Chemistry Department, Faculty of Science, University of Zagreb).

motif (Figure 18a). Furthermore, horizontal hydrogen-bonded layers are vertically interconnected by hydrogen-bonded water molecules (Figure 18b).



Figure 17. Asymmetric unit of **10** contains four symmetrically non-equivalent T and water molecules as well as two symmetrically non-equivalent D molecules.



Figure 18. Crystal packing in **10**. a) Hydrogen-bonded layers viewed down the crystallographic *c*-axis. b) Horizontal hydrogen-bonded layers are vertically interconnected by hydrogen-bonded water molecules. View along crystallographic *b*-axis.

Heating of **4** was monitored in situ by VT-SPXRD (Figure 19a). At around 420 K several low diffraction peaks appear which are a result of gradual re-crystallization of the starting mixture. Next, at approximately 460 K crystallization of **6**, which is presumably D:T cocrystal anhydrate, is observed as evidenced by the appearance of new high intensity diffraction peaks. The appearance of **6**, which remains as the only crystalline phase inside the reaction mixture until the end of monitoring at 490 K, coincides with the crystallization peak at 460 K in DSC trace of neat ground equimolar mixture of D and T (Figure 19b).



Figure 19. a) In situ SPXRD monitoring of heating **4** using synchrotron radiation ($\lambda = 0.20741$ Å). To obtain 2D plot, individual PXRD patterns were stacked upon each other. b) DSC trace of a neat ground equimolar mixture of D and T.

Crystal structure of **6** was solved from the laboratory PXRD data (Figure 20 and Table 7.3). **6** is indeed a D:T cocrystal anhydrate and crystallizes in triclinic $P\overline{1}$ space group. The asymmetric unit of **6** contains two symmetrically non-equivalent D and T molecules, making eight molecules in the unit cell (Figure 21). Crystal packing is characterized by hydrogenbonded layers. Each T molecule is connected to one D molecule by three intermolecular Watson-Crick hydrogen bonds as well as to a second D molecule by $R_2^2(9)$ hydrogen bonding motif (Figure 22). Hydrogen-bonded layers mutually interact through van der Waals interactions.



Figure 20. Rietveld fit of **6** crystal structure that was solved from the laboratory PXRD data ($\lambda = 1.5406 \text{ Å}$).



Figure 21. The unit cell of 6 is composed of four D and four T molecules.



Figure 22. Hydrogen-bonded layers in 6 seen along crystallographic *b*-axis.

4.1.3. Adenine and hypoxanthine

After neat grinding an equimolar amount of powdered A and H, the resulting solid mixture was analyzed by PXRD and revealed that the reaction product **7** was poorly crystalline (Figure 23). Slurry mixing equimolar amounts of A and H in H₂O at 65 °C (Figure 7.7), LAG equimolar amounts of A and H with H₂O at 65 °C (Figure 7.8), and heating **7** in DSC experiment to 300 °C (Figure 7.9) resulted in the formation of a new crystalline phase **8** that was absent of diffraction peaks corresponding to reactants (Figure 23). Furthermore, hydrothermal synthesis of equimolar amounts of A and H resulted in the formation of distinct new crystalline phase **9** (Figure 23).



Figure 23. PXRD patterns for A and H system.

Since crystal structure solution of **8** from the PXRD data indicated that three different structural models were possible, we resolved this conundrum by the combination of crystallography, DFT computation, and solid-state nuclear magnetic resonance (ssNMR) analysis (Figure 24).^c DFT computation showed that model A had by far the highest ground state energy (Figure 24a). The ground state energies of B and C were quite close one to another, although B still had notably lower energy. For model A, the calculated ¹³C and ¹⁵N chemical shifts did not match well with the experimentally obtained shifts (Figure 24b). This model was also excluded by the 2D DQ-SQ BABA (Back-to-Back) ¹H-¹H homonuclear correlation experiment (Figure 24d). However, the latter showed the presence of the diagonal cross peak at about 13.5 ppm in the ¹H dimension, a consequence of protons that resonate at this chemical

^c DFT computational and ssNMR analysis was performed by collaborating co-authors Dr. Gregor Mali (National Institute of Chemistry, Ljubljana) and Dr. Andraž Krajnc (National Institute of Chemistry, Ljubljana).

shift being close in space (i.e., two equivalent protons in two neighboring molecules are quite close one to another). For model A, this was not the case since equivalent N-H protons that would be close in space were absent. This resulted in exclusion of model A. Regarding models B and C, both models predicted ¹H, ¹³C and ¹⁵N chemical shifts that agreed with the experimental ones reasonably well. However, in the ¹³C-detected proton spin diffusion 2D experiment (Figure 24c), we would have expected a strong correlation peak between C2ads and H1ads for model C (highlighted by the blue circle). The peak would have had a similar intensity as the H1hyp-C2hyp cross peak. This was clearly not the case. On the other hand, the observed cross peaks agreed well with the proposed model B. Therefore, a combination of ssNMR experiments and DFT computation suggested that model B is the best model among the three models.



Figure 24. DFT computational and ssNMR analysis. a) Structural model A: adenine (tautomer H9) + hypoxanthine (tautomer H9), structural model B: adenine (tautomer H9) + hypoxanthine (tautomer H7), and structural model C: adenine (tautomer H1) + hypoxanthine (tautomer H7) with DFT ground state energies. b) ¹³C, ¹H, and ¹⁵N calculated and experimental chemical shifts. c) ¹³C-detected proton spin diffusion 2D experiment. d) 2D DQ-SQ Back-to-Back ¹H-¹H homonuclear correlation experiment.

As a result, structural model B which included adenine in 9H tautomeric form and hypoxanthine in 7H tautomeric form was chosen for Rietveld refinement of **8** (Figure 25 and Table 7.4). **8** crystallizes in $P2_1/n$ space group with one adenine and one hypoxanthine molecules in the asymmetric unit and overall, eight molecules in the unit cell (Figure 26). Adenine and hypoxanthine molecules interact through Hoogsteen R_2^2 (9) hydrogen-bonding mode and crystal packing is characterized by hydrogen-bonded sheets (Figures 27 and 28). They are composed of homomeric adenine strands that hydrogen-bond with complementary homomeric hypoxanthine strands (Figure 29). Hypoxanthine molecules are distanced at 3.4 Å and are involved in parallel displaced π - π stacking interactions (Figure 30).



Figure 25. Rietveld fit of 8 crystal structure that was solved from the laboratory PXRD data ($\lambda = 1.5406 \text{ Å}$) with inputs from DFT computation and ssNMR analysis. Note that experimental PXRD pattern contained small amounts of unreacted adenine which was modeled in Rietveld refinement.



Figure 26. The unit cell of 8 viewed from crystallographic *a*-axis.



Figure 27. Top view of hydrogen-bonded sheets in 8 seen along crystallographic *c*-axis. Adenine and hypoxanthine molecules interact through Hoogsteen $R_2^2(9)$ hydrogen-bonding mode.



Figure 28. Side view of hydrogen-bonded sheets in 8 seen along crystallographic *a*-axis.



Figure 29. Sheets in **8** are composed of homomeric adenine strands that hydrogen-bond with complementary homomeric hypoxanthine strands. View along crystallographic *c*-axis.



Figure 30. Hypoxanthine molecules are distanced at ca 3.4 Å and are involved in parallel displaced π - π stacking interactions: a) view from the crystallographic *a*-axis, b) view from the crystallographic *c*-axis.

To solve the crystal structure of **9** we used the identical structural model as in **8** (adenine in 9H tautomeric form and hypoxanthine in 7H tautomeric form) and solved it from the laboratory PXRD data (Figure 31 and Table 7.5). **9** is a channel hydrate and crystallizes in $P2_1/n$ space group with four adenine, four hypoxanthine, and eight water molecules in the unit cell (Figure 32). Hydrogen-bonded sheets are closely packed with water molecules filling structural voids and lying in the channel along crystallographic *c*-axis (Figure 33). Water molecules interact with the exposed C–H groups of adenine and hypoxanthine by van der Waals interactions. Each adenine molecule is involved in two R_2^2 (9) hydrogen bonding motifs and one C(2) chain motif with hypoxanthine molecules, and vice versa (Figure 34). Adenine molecules are distanced at 3.2 Å and are involved in parallel displaced π - π stacking interactions (Figure 35).



Figure 31. Rietveld refinement of **9** crystal structure that was solved from the laboratory PXRD data ($\lambda = 0.7107$ Å) using the same structural model as in **8** (adenine in 9H tautomeric form and hypoxanthine in 7H tautomeric form).



Figure 32. The unit cell of 9 is composed of four adenine, four hypoxanthine, and eight water molecules.



Figure 33. Hydrogen-bonded sheets in **9** are closely packed with water molecules filling structural voids and lying in the channel along crystallographic *c*-axis.



Figure 34. In hydrogen-bonded sheets of 9, each adenine molecule is involved in two $R_2^2(9)$ hydrogen bonding motifs and one C(2) chain motif with hypoxanthine molecules, and vice versa.



Figure 35. Adenine molecules are stacked at ca 3.2 Å and are involved in parallel displaced π - π stacking interactions.

4.1.4. Selectivity of base-pairing and other combinations

The summary of all base-pairing combinations that were investigated in this dissertation is given in Table 1 (Figures 7.10–7.13). D and U, D and T, as well as A and H base-pair in all reaction conditions of different prebiotic water activity. Furthermore, all of them base pair selectively when in the presence of competing nucleobases (Table 2, Figures 7.14-7.18).

Some of the reaction conditions needed for nucleobase self-assembly, such as catalytic presence of water/absence of water and cryogenic/elevated temperatures, might be of astrobiological significance. The individual nucleobases used in our study have been detected in meteorites and their extraterrestrial origin has been ascertained. Meteorites are asteroidal fragments that have fallen to the Earth's surface⁷⁸ and likely served as an exogenous source of prebiotic organic molecules during the time of the heavy bombardment.^{9,16} Moreover, since meteorite composition depends on aqueous alteration, thermal metamorphism, or hydrous metamorphism, these post-accretion events might have presented opportunities for the self-assembly of nucleobases. The latter represents an example of chemical evolution of important biomolecules in laboratory conditions that mimic their extraterrestrial environments.

Reactants	Bulk water	Catalytic presence of water	Dry conditions
D, U	D:U cocrystal hydrate	D:U cocrystal hydrate (2)	D:U cocrystal
	(2)		anhydrate (3)
D, T	D:T cocrystal hydrate	D:T cocrystal hydrate (5 and	D:T cocrystal
	(5 and 10)	10)	anhydrate (6)
D, X	poorly crystalline,	poorly crystalline,	poorly crystalline,
	inconclusive	inconclusive	inconclusive
A, H	A:H cocrystal	A:H cocrystal anhydrate (8)	A:H cocrystal
	anhydrate (8) and		anhydrate (8)
	hydrate (9)		
C, H	physical mixture	physical mixture	physical mixture
U, H	physical mixture	physical mixture	physical mixture
G, U	physical mixture	physical mixture	physical mixture

Table 1. Results of base-pairing experiments using different prebiotic water availability.

Reactants	Bulk water	Catalytic presence of water	Dry conditions
D, U, A	D:U cocrystal hydrate	D:U cocrystal hydrate (2) +	D:U cocrystal
	(2) + unreacted A	unreacted A	anhydrate (3) +
			unreacted A
D, T, A	D:T cocrystal hydrate	D:T cocrystal hydrate (5 and	D:T cocrystal
	(5) + unreacted A	10) + unreacted A	anhydrate (6) +
			unreacted A
D, U, T	D:U cocrystal hydrate	D:U cocrystal hydrate (2) +	D:U cocrystal
	(2) + D:T cocrystal	D:T cocrystal hydrate (5 and	anhydrate (3) +
	hydrate (5)	10)	D:T cocrystal
			anhydrate (6)
A, H, U	A:H cocrystal	A:H cocrystal anhydrate (8) +	A:H cocrystal
	anhydrate (8) +	unreacted U	anhydrate (8) +
	unreacted U		unreacted U
A, H, T	A:H cocrystal	A:H cocrystal anhydrate (8) +	A:H cocrystal
	anhydrate (8) +	unreacted T	anhydrate (8) +
	unreacted T		unreacted T

Table 2. Results of selectivity experiments in the presence of competing nucleobases.

4.2. Mechanochemical prebiotic peptide synthesis

4.2.1. Amino acid oligomerization

Pristine Gly was milled at room temperature conditions and the product was analyzed by IP-HPLC (Figure 36a).^d It showed that trace amount of 2,5-diketopiperazine (DKP) and Gly₂ have formed. After that, we milled Gly with five-fold molar excess of TiO₂. We used TiO₂ as a mineral additive because it is known to have good amino acid adsorption properties and is prebiotically relevant.⁶⁵ The excess of mineral additive was used to simulate prebiotic environments where it is regarded that organic material was present in limited amounts on mineral surfaces. Same is true for extraterrestrial matter, i.e., meteorites, albeit to a greater extreme. There, amino acids are present in ppm quantities in complex mineral matrix. In our case, milling Gly with TiO_2 at ambient conditions clearly showed the formation of Gly_2 and Gly₃, along with DKP and unreacted Gly (Figure 36b). Next, we conducted ball milling reactions with TiO₂ at temperatures ranging from ambient to 130 °C (Figure 37a-e), guided by recent literature examples for temperature limitation.^{67,68} These have shown that increase in milling temperature results in the formation of higher Gly oligomers, along with the formation of DKP as the dominant component of the reaction mixture. The highest Gly oligomer was obtained by milling Gly with TiO₂ at 130 °C, Gly₁₀ as detected by IP-HPLC (Figure 37e). The analysis by LC/MS (Q-TOF) confirmed that even Gly₁₁ was obtained in the same reaction (Figure 38).^e

^d IP-HPLC analysis was performed jointly with collaborating co-author Mr. Saša Grubešić (Xellia Pharmaceuticals).

^e LC/MS (Q-TOF) analysis was performed by collaborating co-author Mr. Saša Grubešić (Xellia Pharmaceuticals).



Figure 36. IP-HPLC chromatograms for: (a) pristine Gly milled at RT, (b) Gly milled with TiO₂ at RT, (c) Gly standard, (d) Gly₂ standard, (e) Gly₃ standard, and (f) DKP standard. The inset shows the zoomed region and reveals the presence of Gly₂ and Gly₃ in (a) and (b).



Figure 37. IP-HPLC chromatograms for Gly milled with TiO_2 at: (a) RT, (b) 40 °C, (c) 70 °C, (d) 100 °C, and (e) 130 °C. The inset shows the zoomed region of retention times in the 6.1–11 min range.



Figure 38. UPLC-MS analysis of Gly milled with TiO₂ at 130 °C. The inset shows the zoomed region in the acquisition time range from 14–17 min.

We also used IP-HPLC method to quantitatively determine the total yield of linear oligomers, according to the recent report (Table 3).⁸² Total yield of linear oligomers was highest for Gly milled with TiO₂ at 100 °C, which had almost identical yield as Gly milled with TiO₂ at 70 °C. On one hand, these two temperatures showed that elevated temperature is needed to increase the conversion of Gly to linear oligomers. On the other hand, 130 °C exhibited lower linear Gly oligomer yield due to high temperature favoring Gly cyclization to DKP.
Ball milling temperature	Longest detected oligomer ^[a]	Yield of linear oligomers
ambient	Gly_6	6.7 %
40 °C	Gly_6	6.7 %
70 °C	$\mathrm{Gly}_8^{[b]}$	10.1 %
100 °C	Gly ₈	10.2 %
130 °C	Gly ₁₁ ^[b]	8.5 %

Table 3. Total yields of linear Gly oligomers for milling Gly with TiO_2 at different temperatures. [a] According to IP-HPLC. [b] According to UPLC-MS.

Furthermore, we conducted experiments of milling Gly₂, Gly₃, and DKP with TiO₂ (Figure 39). These have shown that odd and even numbered oligomers have formed irrespective of the starting reactant, and that DKP is a productive intermediate in the mechanochemical prebiotic peptide bond formation.



Figure 39. IP-HPLC chromatograms for: (a) Gly milled with TiO_2 at RT, (b) Gly₂ milled with TiO_2 at 130 °C, (c) Gly₃ milled with TiO_2 at 130 °C, and (d) DKP milled with TiO_2 at RT. The inset shows the zoomed region of retention times in the 6.1–11 min range.

Besides TiO₂, we tested other mineral additives, namely SiO₂, mica, and montmorillonite (Figures 7.19–7.21). All of them resulted in Gly oligomerization. LAG reactions with milling Gly in the presence of water at RT and 130 $^{\circ}$ C showed that mechanochemical prebiotic peptide bond formation proceeds in the presence of moisture (Figures 7.22 and 7.23).

To test if oligomers of other amino acids oligomerize under mechanochemical conditions, we milled L-alanine (Ala) with TiO₂ at 130 °C. The analysis by IP-HPLC showed

that Ala₂ formed (Figure 7.24), while direct MS analysis showed that oligomers up to Ala₅ formed (Table 4). After that, we conducted milling Gly and Ala with TiO₂ at 100 °C to see if hetero oligomers might form under mechanochemical conditions. Indeed, IP-HPLC indicated that might be the case (Figure 7.25), which was confirmed by direct MS analysis (Table 4).^f

			Ala milled at 130°C		Gly+Ala milled at 100°C	
	Elementary	Theoretical	Experimental		Experimental	
Oligomers	formulae	[M+H] ⁺ (m/z)	$[M+H]^{+}(m/z)$	Δppm	$[M+H]^{+}(m/z)$	Δppm
Gly	C ₂ H ₅ NO ₂	76,0393			76,0395	2,63
Gly ₂	$C_4H_8N_2O_3$	133,0608			133,0611	2,25
G1y3	$C_6H_{11}N_3O_4$	190,0822			190,0825	1,58
Gly4	$C_8H_{14}N_4O_5$	247,1037				
G1y5	$C_{10}H_{17}N_5O_6$	304,1252				
Gly ₆	$C_{12}H_{20}N_6O_7$	361,1466				
Gly7	C14H23N7O8	418,1681				
G1y8	C ₁₆ H ₂₆ N ₈ O ₉	475,1896				
G1y9	C ₁₈ H ₂₉ N ₉ O ₁₀	532,2110				
G1y10	$C_{20}H_{32}N_{10}O_{11}$	589,2325				
Gly ₁₁	$C_{22}H_{35}N_{11}O_{12}$	646,2539				
Gly ₁₂	$C_{24}H_{38}N_{12}O_{13}$	703,2754				
DKP(Gly)	$C_4H_6N_2O_2$	115,0502				
Ala	$C_3H_7NO_2$	90,0550	90,0554	4,44	90,0554	4,44
Al a ₂	$C_{6}H_{12}N_{2}O_{3}$	161,0921	161,0924	1,86	161,0924	1,86
Al a ₃	C9H17N3O4	232,1292	232,1298	2,58		
Ala ₄	$C_{12}H_{22}N_4O_5$	303,1663	303,1675	3,96		
Al a ₅	$C_{15}H_{27}N_5O_6$	374,2034	374,205	4,28		
DKP(Ala)	$C_{6}H_{10}N_{2}O_{2}$	143,0815	143,0817	1,40		
Gly-Ala	$C_5H_{10}N_2O_3$	147,0764			147,0767	2,04
Ala-Ala-Gly	$C_8H_{15}N_3O_4$	218,1135			218,1136	0,46
Gly-Gly-Ala	C7H13N3O4	204,0979			204,098	0,49

Table 4. Summary of each oligomer detected by the direct injection to MS (Q-TOF).

^f Direct MS Q-TOF analysis was performed by collaborating co-author Mr. Saša Grubešić (Xellia Pharmaceuticals).

We further set out to explore some of the critical reaction parameters that represent different geological and environmental scenarios to make them more universally plausible. These are the effects of mechanical loading and time. The transfer of mechanical load needed to initiate oligomerization is a crucial factor for mechanochemical peptide bond formation. Molecular collisions in ball mills are typically in the range of m/s,^{145,146} and although they have much milder mechanical loading when compared to impact events which are in the km/s range, their relevance for prebiotic environments is hard to predict given the uncertainty and frequency of their appearance over time. On that note, subaerial geomorphological processes such as erosion and weathering are known to occur frequently and are capable of exchanging matter on the planetary surface.¹⁴⁷ Therefore, it is of interest to decouple the effect of solid-state mixing from that of mechanical activation in the solid-state.¹⁴⁸

We simulated different mechanical loading that is transferred to the reaction mixture by using 5 Hz, 15 Hz, and 30 Hz milling frequencies. We propose that low milling frequency of 5 Hz might simulate the erosion of land and subaerial weathering. At 5 Hz, typical sound coming from the collision of milling balls is absent, the reaction mixture is gently shaken, and homogenization is limited. Intermediate milling frequency of 15 Hz might simulate the conditions that arise during earthquakes.¹⁴⁹ Highest milling frequency of 30 Hz might best simulate post-impact conditions of extraterrestrial bodies falling to Earth surface or rocks hitting the ground during heavy winds and volcanic eruptions. Our results show that the highest amounts of peptides are formed when milling is conducted at 30 Hz. The dependance of the total yield of linear Gly oligomers that are formed under these conditions on the reaction time are presented in Figure 40. The comparison of the effect of all three milling frequencies on mechanochemical peptide bond formation are summarized in Table 5 (Figures 7.26–7.37). IP-HPLC and LC/MS (Triple quadrupole)^g analysis showed that 5 and 15 Hz frequencies enable peptide bond formation. The longest Gly oligomers were obtained by milling Gly with SiO₂ at room temperature for 48 h using 30 Hz frequency, Gly₈ (Table 5). Furthermore, milling Gly with TiO₂ using three stones for 18 h at 20 Hz and ambient temperature resulted in the formation of oligomers up to Gly₄ (Figure 41). Importantly, milling in N₂ and CO₂ atmospheres showed

^g LC/MS (Triple quadrupole) analysis was performed by collaborating co-author Mr. Saša Grubešić (Xellia Pharmaceuticals).

that mechanochemical peptide bond formation proceeds in the presence of these two gases which were likely the main components of prebiotic Earth atmosphere,¹⁵⁰ although it gives a more complex mixture of products (Figures 7.39–7.42). For example, there is a high intensity peak at 6.1 min of retention time which we did not identify. Nevertheless, LC/MS (Triple quadrupole) analysis identified the presence of Gly₆ and Gly₅ when reaction was performed in N₂ (Figure 7.40) and CO₂ (Figure 7.42) atmospheres, respectively.



Figure 40. The effect of reaction time for milling Gly with SiO₂ at 40 °C on total yield of linear Gly oligomers.

Table 5. The longest glycine oligomers detected by: [a] IP-HPLC and [b] LC/MS (Triple quadrupole) for Gly milled with SiO₂ at room temperature.

	5 Hz	15 Hz	30 Hz
18 h	[a] Gly ₃	[a] Gly ₄	[a] Gly ₈
40 11	[b] Gly ₂	[b] Gly ₃	[b] Gly ₈
06 h	[a] Gly ₂	[a] Gly ₃	[a] Gly7
70 II	[b] Gly ₃	[b] Gly ₃	[b] Gly ₄

54



Figure 41. IP-HPLC chromatogram for milling Gly with TiO₂ using three hand-picked stones (Figure 7.38) for 18 h at 20 Hz and ambient temperature.

After studying the effects of mechanical loading, time, temperature, and atmosphere, we turned to oligomerization on complex mineral surfaces. We have selected different mineral classes including silicates (quartz, talc), sulfides (pyrite), sulphates (gypsum), phosphates (apatite), carbonates (calcite, magnesite), micas (muscovite), and zeolites (analcime, clinoptilolite) (Figure 42). The elemental composition of minerals was verified by X-ray fluorescence^h (Table 6) and PXRD analysis combined with Rietveld analysis (Figures 7.43–7.52). Milling reactions were performed with the excess of minerals under the same molar ratio as those with TiO₂ and SiO₂, namely 5:1. We used 30 Hz frequency, two stainless steel balls as milling media, room temperature, and experiments were performed for 16 h. Reaction products were analyzed by IP-HPLC and LC/MS triple quadrupole (Figures 7.53–7.71). The results of these experiments are summarized in Table 7. Among all tested minerals, quartz yielded the longest oligomer detected by LC/MS triple quadrupole, Gly₇. Notably, unidentified peak at 6.3 min of retention time is present in all milling reactions with minerals alongside glycine

^h XRF analysis of minerals elemental composition was performed by Dr. Ivan Nemet (Chemistry Department, Faculty of Science, University of Zagreb).

oligomers (same as in Figures 7.40 and 7.42 showing chromatograms for milling glycine with SiO_2 in N_2 and CO_2 atmospheres).



Figure 42. Mineral samples that were used in this study.

Name	Formula	Method	w / %	Origin
analcime	NaAlSi ₂ O ₆ (H ₂ O)	XRF	SiO ₂ 55,26 Al ₂ O ₃ 22,40 FeO 0,07 CaO 3,12 MnO 0,06	Tirol, Austria
apatite	Ca ₅ (PO ₄) ₃	XRF	CaO 54,51 P ₂ O ₅ 39,40 FeO 0,196 MnO 0,06	Bancroft, Canada
calcite	CaCO ₃	XRF	CaO 55,52 FeO 0,02 MnO 0,03 SiO ₂ 0,05	Trepča, Kosovo
clinoptilolite	Ca1.9Na1.76K1.05Mg0.17Al6.72Si29.2O72	XRF	SiO ₂ 68,75 Al ₂ O ₃ 12,11 FeO 1,362 CaO 3,92 K 1,27 MnO 0,02	Serbia
gypsum	CaSO ₄ x 2(H ₂ O)	XRF	CaO 31,90 SO ₃ 46,1 SiO ₂ 0,14	Mt. Paderno, Italy
magnesite	MgCO ₃	XRF	MgO 47,60 K ₂ O 0,06 MnO 0,07 FeO 0,02 SiO ₂ 0,03	Goleš, Kosovo
muscovite	KAl ₂ (Si ₃ Al)O ₁₀ (OH,F) ₂	XRF	SiO ₂ 46,12 Al ₂ O ₃ 37,80 K ₂ O 10,50 MnO 0,02	Čanište, Macedonia
pyrite	FeS ₂	XRF	Fe 45,61 S 52,22 SiO ₂ 0,05 MnO 0,06	Trepča, Kosovo
talc	$Mg_3Si_4O_{10}(OH)_2$	XRF	SiO ₂ 60,95 MgO 31,50 Fe ₂ O ₃ 1,163 CaO 0,222	Rabenwald, Austria
quartz	SiO ₂	XRF	SiO ₂ 95,5	Trepča, Kosovo

Table 6. Information about minerals origin and elemental composition as determined by X-ray fluorescence (XRF). Minerals were provided by Prof. Nenad Tomašić (Department of Geology, University of Zagreb).

Mineral	Longest detected oligomer by LC/MS (triple quadrupole)
analcime	Gly ₂
apatite	Gly ₃
calcite	Gly ₄
clinoptilolite	Gly ₂
gypsum	Gly ₂
magnesite	Gly4
muscovite	Gly ₂
pyrite	Gly ₃
talc	Gly ₃
quartz	Gly7

Table 7. Summary of reactions where Gly was milled with different minerals.



Figure 43. LC/MS (triple quadrupole) chromatograms for milling Gly with quartz at room temperature for 16 h and at 30 Hz.

59

4.2.2. Amino acid and α -hydroxy acid oligomerization

Together with amino acids, among the soluble organic material that is found in meteorites are also α -hydroxy acids. Therefore, investigation of their mutual interaction and the production of prebiotic oligomers is desired.^{151,152} First, we milled glycolic acid (GA) with SiO₂ at room temperature, for 16 h, and at 30 Hz. The analysis by IP-HPLC showed several peaks which indicated that we might have synthesized ester bonds (Figure 7.72). Indeed, MALDI TOF analysisⁱ confirmed that oligomers up to four glycolic acid residues have formed under these conditions (Figure 44 and Table 8).



Figure 44. MALDI TOF spectra of the reaction product after milling glycolic acid with SiO₂ at room temperature, at 30 Hz, and for 16 h. Showing: a) glycolic acid, b) glycolic acid dimer, c) glycolic acid trimer, d) glycolic acid tetramer.

ⁱ MALDI TOF analysis was performed by collaborating co-author Dr. Mario Cindrić (Ruđer Bošković Institute).

Table 8. Summary of oligomers that were detected by MALDI TOF after milling glycolic acid with SiO_2 at room temperature, for 16 h, and at 30 Hz. There is no protonation due to the formation of oxazolone ion.

Product	Elementary formulae	Theoretical (m/z)	Experimental (m/z)	Δ ppm
1GA	$C_2H_4O_3$	76.0160	76.0160	0.00
2GA	$C_4H_6O_5$	134.0215	134.0211	-2.98
3GA	$C_6H_8O_7$	192.0270	192.0271	0.52
4GA	$C_8H_{10}O_9$	250.0325	250.0330	2.00

Encouraged by the formation of oligomers with ester bonds, glycine and glycolic acid were milled with SiO₂ at room temperature, for 16 h, and at 30 Hz. IP-HPLC chromatogram showed numerous peaks which indicated the rich reactivity under these reaction conditions (Figure 7.73). This was confirmed by MALDI TOF analysis which showed the formation of at least five different oligomers containing peptide and ester bonds (Figure 45). Furthermore, Table 9 contains detailed list of theoretical and experimentally observed molecular masses and Figure 46 shows molecular structures of products that were detected.



Figure 45. MALDI TOF spectra of the reaction product after milling glycine and glycolic acid with SiO₂ at room temperature, at 30 Hz, and for 16 h. Showing: a) glycine and glycolic acid dimer, b) glycine and glycolic acid trimer, c) glycine and glycolic acid tetramer, d) glycine and glycolic acid hexamer.

Table 9. Summary of oligomers that were detected by MALDI TOF after milling glycine and glycolic acid with SiO₂ at room temperature, for 16 h, and at 30 Hz.

Draduat	Elementary	Theoretical [M+H] ⁺	Experimental	A nnm
Product	formulae	(m/z)	$[M+H]^+(m/z)$	д ррт
1Gly+1GA	C4H7NO4	134.0453	134.0448	-3.73
1Gly+2GA	C ₆ H ₉ NO ₆	192.0503	192.0504	0.52
1Gly+3GA	$C_8H_{11}NO_8$	250.0558	250.0551	-2.80
2Gly+3GA	$C_{10}H_{14}N_2O_9$	307.0727	307.0717	-3.26
3Gly+3GA	$C_{12}H_{17}N_3O_{10}$	364.0987	364.0971	-4.39



Figure 46. Reaction scheme for milling glycine and glycolic acid with SiO₂ at room temperature, for 16 h, and at 30 Hz. Represented oligomers were detected by MALDI TOF (Figure 45 and Table 9).

§ 5. CONCLUSION

Our results show that some of the molecular recognition patterns present in contemporary biochemistry are available on a nucleobase level under prebiotic conditions. Molecular recognition of 2,6-diaminopurine (D) and uracil (U), D and thymine (T), and adenine (A) and hypoxanthine (H) is selective and occurs under all reaction conditions that were tested: bulk water, catalytic presence of water, and in the absence of water (in the solid-state). The crystal structures of D:U cocrystal hydrate and D:T cocrystal hydrate were solved from the singlecrystal X-ray diffraction data. The crystal structures of D:T cocrystal anhydrate, A:H cocrystal hydrate, and A:H cocrystal anhydrate were solved from the laboratory powder X-ray diffraction data. Interestingly, to self-assemble with A in both A:H cocrystal hydrate and anhydrate, H changes its tautomeric form (from naturally occurring 9H tautomer to 7H tautomer). Contrary to the literature fact that canonical nucleobase pairs (A:T, A:U, and G:C) do not self-assemble in water, we show that water molecules are enabling molecular recognition of D:U, D:T, and A:H nucleobase pairs. The latter is true even in the presence of competing nucleobases, e.g., U and T self-assemble with D in the presence of A, and A self-assembles with H in the presence of U and T. Since prebiotic synthetic pathways usually result in a mixture of nucleobases instead of a clean synthesis of a single nucleobase and multiple nucleobases are found in meteorites, it is reasonable to expect that they would have been in spatial proximity in an early Earth setting. For example, in impact craters where nucleobases would jointly diffuse from extraterrestrial bodies such as meteorites to the newly formed ponds. Therefore, it seems that supramolecular interactions between nucleobases might have been important for their chemical evolution. The influence of supramolecular interactions of organic building blocks of life on their chemical evolution is underrepresented in the literature. Most often, the propensity for the formation of covalent condensation bonds is taken as the criteria to assess whether a compound or a reaction pathway, is regarded as plausibly prebiotic. A synergistic approach based on the formation of covalent and non-covalent interactions may provide new insights into chemical evolution. For example, the existence of different tautomeric forms of nucleobases (or other organic building blocks) in the solid-state might offer different reactivity and selectivity in covalent bondforming reactions such as glycosylation or phosphorylation. Besides, UV irradiation is considered to have been one of the main selection pressures on early Earth. Therefore, supramolecular assemblies of nucleobases might have different photochemical properties when compared to individual nucleobases.

In the second part of the thesis, mechanical activation of amino acids with TiO₂ and SiO₂ enabled their oligomerization. For the analysis of the reaction products, we relied on ionpairing high-performance liquid chromatography and mass spectrometry (Q-TOF, triple quadrupole, MALDI TOF). The longest glycine (Gly) oligomer that was obtained from the starting Gly was Gly₁₁. Milling Gly with TiO₂ (at 100 °C, 30 Hz, three 1 g stainless steel balls, for 18 h) resulted in the highest total yield of linear Gly oligomers, 10.2 %. Liquid-assisted grinding experiments with H₂O showed that mechanochemical peptide bond formation tolerates the presence of moisture. Reactions that started from Gly₂, Gly₃, and DKP showed that mechanochemical peptide bond formation is a dynamic process with simultaneous breaking and making of peptide bonds. Importantly, milling Gly and L-alanine (Ala) resulted in the formation of their hetero oligopeptides thus demonstrating that other amino acids can be incorporated into a growing peptide chain. It was shown that time, temperature, atmosphere, and mechanical loading are all important parameters for mechanochemical prebiotic peptide synthesis. Even milling at low frequencies and using stones as milling media resulted in the formation of peptides. It was also shown that glycolic acid (GA) oligomerizes under mechanochemical conditions. Milling GA with SiO₂ (at room temperature, 30 Hz, two 1.4 g stainless steel balls, for 16 h) resulted in the formation of an oligo-ester containing four residues of GA. Furthermore, milling Gly and GA with SiO₂ (at room temperature, 30 Hz, two 1.4 g stainless steel balls, for 16 h) resulted in the formation of at least five hetero oligomers containing peptide and ester bonds. This work demonstrated that the formation of peptides is operational under different geological and environmental scenarios in a solid-state reaction medium simulating dry prebiotic environment. These results may have implications for astrobiology research and reconsideration of extra-terrestrial conditions that can give rise to peptides. For example, weathering of amino acids in dry environments might result in the formation of peptides on mineral surfaces. Once formed, peptides might have catalyzed other important prebiotic reactions such as sugar syntheses. On the other hand, mechanochemical prebiotic peptide bond formation opens the possibility for the sustainable syntheses of modern peptides and amides by mechanochemistry.

§ 6. REFERENCES

- 1. L. E. Orgel, Crit. Rev. Biochem. Mol. Biol. 39 (2004) 99–123.
- 2. A. I. Oparin, *Proischogdenie Zhizni*, Moscovsky Robotchii, Moscow, 1924.
- 3. Eschenmoser, *Tetrahedron* **63** (2007) 12821–12844.
- 4. G. Wächtershäuser, *Microbiol Rev.* **52** (1988) 452–484.
- 5. Damer, D. Deamer, *Astrobiology* **20** (2020) 429–452.
- 6. S. Ross, D. Deamer, *Life* **6** (2016) 28.
- 7. M. Marshall, *Nature* **588** (2020) 210–213.
- 8. B. Damer, D. Deamer, *Life* **5** (2015) 872–887.
- B. K. Pearce, R. E. Pudritz, D. A. Semenov, T. K. Henning, *Proc. Natl. Acad. Sci. U.S.A.* 114 (2017) 11327–11332.
- M. J. V. Kranendonk, R. Baumgartner, T. Djokic, T. Ota, L. Steller, U. Garbe, E. Nakamura, *Astrobiology* 21 (2021) 39–59.
- H. J. Cleaves II, A. Michalkova Scott, F. C. Hill, J. Leszczynski, N. Sahai, R. Hazen, *Chem. Soc. Rev.* 41 (2012) 5502–5525.
- K. Kvenvolden, J. Lawless, K. Pering, E. Peterson, J. Flores, C. Ponnamperuma, I. R. Kaplan, C. Moore, *Nature* 228 (1970) 923–926.
- M. P. Callahan, K. E. Smith, H. J. Cleaves II, J. Ruzicka, J. C. Stern, D. P. Glavin, C. H. House, J. P. Dworkin, *Proc. Natl. Acad. Sci. U.S.A.* 108 (2011) 13995–13998.
- Y. Furukawa, Y. Chikaraishi, N. Ohkouchi, N. O. Ogawa, D. P. Glavin, J. P. Dworkin,
 C. Abe, T. Nakamura, *Proc. Natl. Acad. Sci. U.S.A.* 116 (2019) 24440–24445.
- P. Schmitt-Kopplin, Z. Gabelica, R. D. Gougeon, A. Fekete, B. Kanawati, M. Harir, I. Gebefuegi, G. Eckel, N. Hertkorn, *Proc. Natl. Acad. Sci. U.S.A.* 107 (2010) 2763–2768.
- 16. Chyba, C. Sagan, *Nature* **355** (1992) 125–132.
- 17. A. J. Brearley, *Meteor. Early Sol. Syst. II* (2006) 587–624.
- 18. G. R. Huss, A. E. Rubin, J. N. Grossman, Meteor. Early Sol. Syst. II (2006) 567–586.
- 19. V. E. Hamilton, C. A. Goodrich, A. H. Treiman, H. C. Connolly Jr, M. E. Zolensky, M. H. Shaddad, *Nat Astron* 5 (2020) 350–355.
- 20. C. J. Bennett, C. Pirim, T. M. Orlando, *Chem. Rev.* **113** (2013) 9086–9150.
- 21. T. Okada et al., Nature **579** (2020) 518–522.

- 22. A. A. Simon *et al.*, *Science* **370** (2020) eabc3522.
- T. Stolar, S. Grubešić, N. Cindro, E. Meštrović, K. Užarević, J. G. Hernández, Angew. Chem. Int. Ed. 60 (2021) 12727–12731.
- 24. S. A. Benner, H.-J. Kim, M. A. Carrigan, Acc. Chem. Res. 45 (2012) 2025–2034.
- 25. J. D. Sutherland, *Nat Rev Chem* **1** (2017) 1–7.
- 26. M. Yadav, R. Kumar, R. Krishnamurthy, Chem. Rev. 120 (2020) 4766–4805.
- 27. M. A. Pasek, Chem. Rev. 120 (2020) 4690–4706.
- 28. Preiner, M. et al. *Life* **10** (2020) 1–25.
- 29. A. C. Rios, Y. Tor, Isr. J. Chem. 53 (2013) 1–15.
- 30. M. Fialho, T. P. Roche, N. V. Hud, *Chem. Rev.* **120** (2020) 4806–4830.
- 31. S. Miyakawa, H. J. Cleaves, S. L. Miller, *Orig Life Evol Biosph* **32** (2002) 209–218.
- R. Saladino, C. Crestini, F. Ciciriello, G. Costanzo, E. Di Mauro, *Chem. Biodivers.* 4 (2007) 694–720.
- 33. C. Menor-Salván, M. R. Marín-Yaseli, Chem. Soc. Rev. 41 (2012) 5404–5415.
- 34. C. Menor-Salvan, M. R. Marín-Yaseli, *Chem. Eur. J.* **19** (2013) 6488–6497.
- 35. I. Jiménez, C. Gibard, R. Krishnamurthy, Angew. Chem. Int. Ed. 60 (2021) 2–11.
- 36. Z. Martins, et al. Earth Planet. Sci. Lett. 270 (2008) 130–136.
- 37. C. K. Materese, M. Nuevo, S. A. Sandford, *Astrobiology* **17** (2017) 761–770.
- Y. Oba, Y. Takano, H. Naraoka, N. Watanabe, A. Kouchi, *Nat. Commun.* 10 (2019) 4413.
- N. V. Hud, B. J. Cafferty, R. Krishnamurthy, L. D. Williams, *Chem. & Biol.* 20 (2013) 466–474.
- 40. P. O. P. Ts'o, New York: Acad. Press. I (1974) 453–584.
- 41. P. Cieplak, P. A. Kollman, J. Am. Chem. Soc. 110 (1988) 3734–3739.
- 42. E. Stofer, C. Chipot, R. Lavery, J. Am. Chem. Soc. 121 (1999) 9503–9508.
- 43. B. A. Anderson, K. Fauché, S. C. Karunakaran, J. R. Yerabolu, N. V. Hud, R. Krishnamurthy, *Chem. Eur. J.* **27** (2021) 4033-4042.
- 44. M. D. Kirnos, I. Y. Khudyakov, N. I. Alexandrushkina, B. F. Vanyushin, *Nature* **270** (1977) 369–370.
- 45. Y. Khudyakov, M. D. Kirnos, N. I. Alexandrushkina, B. F. Vanyushin, *Virology* **18** (1978) 8–18.

- V. Pezo, F. Jaziri, P.-Y. Bourguignon, D. Louis, D. Jacobs-Sera, J. Rozenski, S. Pochet,
 P. Herdewijn, G. F. Hatfull, P.-A. Kaminski, P. Marliere, *Science* 372 (2021) 520–524.
- D. Sleiman, P. S. Garcia, M. Lagune, J. Loc'h, A. Haouz, N. Taib, P. Röthlisberger, S. Gribaldo, P. Marlière, P. A. Kaminski, *Science* 372 (2021) 516–520.
- 48. Y. Zhou, X. Xu, Y. Wei, Y. Cheng, Y. Guo, I. Khudyakov, F. Liu, P. He, Z. Song, Z. Li, Y. Gao, E. L. Ang, H. Zhao, Y. Zhang, S. Zhao, *Science* 372 (2021) 512–516.
- 49. S. Reader, G. F. Joyce, *Nature* **420** (2002) 841–844.
- 50. F. H. C. Crick, J. Mol. Biol. 38 (1968) 367–379.
- 51. E. Orgel, J. Mol. Biol. 38 (1968) 381–393.
- 52. Groebke, J. Hunziker, W. Fraser, L. Peng, U. Diederichsen, K. Zimmermann, A. Holzner, C. Leumann, A. Eschenmoser, *Helv. Chim. Acta* **81** (1998) 375–474.
- 53. B. D. Heuberger, C. Switzer, *ChemBioChem* **9** (2008) 2779–2783.
- 54. T. R. Battersby, M. Albalos, M. J. Friesenhahn, Chem. & Biol. 14 (2007) 525–531.
- 55. F. H. C. Crick, J. Mol. Biol. 19 (1966) 548–555.
- 56. J. M. Berg, J. L. Tymoczko, G. J. Gatto, L. Stryer, Biochemistry 8th Ed. 893–924 (2015), W. H. Freeman and Company, New York.
- 57. Danger, R. Plasson, R. Pascal, Chem. Soc. Rev. 41 (2012) 5416–5429.
- Frenkel-Pinter, M. Samanta, G. Ashkenasy, L. J. Leman, *Chem. Rev.* 120 (2020) 4707–4765.
- 59. R. Cronin, C. B. Moore, *Science* **172** (1971) 1327–1329.
- 60. Altwegget *et al.*, *Sci. Adv.* **2** (2016) e160028.
- S. Ioppolo, G. Fedoseev, K. J. Chuang, H. M. Cuppen, A. R. Clements, M. Jin, R. T. Garrod, D. Qasim, V. Kofman, E. F. van Dishoeck, H. Linnartz, *Nat Astron* 5 (2021)197–205.
- Lange, F. Djago, B. Eddhif, Q. B. Remaury, A. Ruf, N. K. V. Leitner, L. L. S. Hendecourt, G. Danger, C. G. Rodier, S. Papot, P. Poinot, *J. Proteome Res.* 20 (2021) 1444–1450.
- 63. A. Shimoyama, R. Ogasawara, Orig. Life Evol. Biosph. 32 (2002) 165–179.
- 64. S. W. Fox, K. Harada *Science* **128** (1958) 1214.
- 65. G. Martra, C. Deiana, Y. Sakhno, I. Barberis, M. Fabbiani, M. Pazzi, M. Vincenti, *Angew. Chem. Int. Ed.* **53** (2014) 4671–4674.

- 66. J. G. Forsythe, S.-S. Yu, I. Mamajanov, M. A. Grover, R. Krishnamurthy, F. M. Fernández, N. V. Hud, *Angew. Chem. Int. Ed.* **54** (2015) 9871–9875.
- Rodriguez-Garcia, A. J. Surman, G. J.T. Cooper, I. Suárez-Marina, Z. Hosni, M. P. Lee,
 L. Cronin, *Nat. Commun.* 6 (2015) 8385.
- T. D. Campbell, R. Febrian, J. T. McCarthy, H. E. Kleinschmidt, J. G. Forsythe, P. J. Bracher, *Nat. Commun.* 10 (2019) 4508.
- 69. G. Schwendinger, B. M. Rode, Anal Sci 5 (1989) 411.
- F. Sauer, M. Haas, C. Sydow, A. F. Siegle, C. A. Lauer, O. Trapp, *Nat Commun* 12 (2021) 7182.
- 71. J. Greenwald, M. P. Friedmann, R. Riek, Angew. Chem. Int. Ed. 55 (2016) 11609– 11613.
- 72. P. Canavelli, S. Islam, M. W. Powner, *Nature* **571** (2019) 546–549.
- 73. J. P. Ferris, R. A. Hill Jr, R. Liu, L. E. Orgel, *Nature* **381** (1996) 59–61.
- J. G. Blank, G. H. Miller, M. J. Ahrens, R. E. Winans, Orig Life Evol Biosph 31 (2001) 15–51.
- 75. H. Sugahara, K. Mimura, *Geochem. J.* **48** (2014) 51–62.
- B. A. Steele, N. Goldman, I.-F. W. Kuo, M. P. Kroonblawd, *Chem. Sci.* 11 (2020) 7760– 7771.
- 77. H. G. Hansma, J. Theor. Biol. 266 (2010) 175–188.
- A. S. Burton, J. C. Stern, J. E. Elsila, D. P. Glavin, J. P. Dworkin, *Chem. Soc. Rev.* 41 (2012) 5459–5472.
- 79. S. Pizzarello, Acc. Chem. Res. **39** (2006) 231–237.
- 80. H. Sugahara, K. Mimura, *Icarus* **257** (2015) 103–112.
- Bertrand, S. van der Gaast, F. Vilas, F. Hçrz, G. Haynes, A. Chabin, A. Brack, F. Westall, *Astrobiology* 9 (2009) 943–951.
- 82. T. D. Campbell, R. Febrian, H. E. Kleinschmidt, K. A. Smith, P. J. Bracher, *ACS Omega* 4 (2019) 12745–12752.
- 83. L. Takacs, Chem. Soc. Rev. 42 (2013) 7649–7659.
- S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D.
 M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin,
 W. C. Shearouse, J. W. Steed, D. C. Waddell, *Chem. Soc. Rev.* 41 (2012) 413–447.

- Baláž, M. Achimovičová, M. Baláž, P. Billik, A. Cherkezova-Zheleva, J. M. Criado, F. Delogu, E. Dutková, E. Gaffet, F. L. Gotor, R. Kumar, I. Mitov, T. Rojac, M. Senna, A. Streletskii, K. Wieczorek-Ciurowa, *Chem. Soc. Rev.* 42 (2013) 7571–7637.
- 86. T. Friščić, C. Mottillo, H. M. Titi, Angew. Chem. Int. Ed. 59 (2020) 1018–1029.
- A. A. L. Michalchuk, E. V. Boldyreva, A. M. Belenguer, F. Emmerling, V. V. Boldyrev, *Front. Chem.* 9 (2021) 359.
- S. Lukin, M. Tireli, T. Stolar, D. Barišić, M. V. Blanco, M. di Michiel, K. Užarević, I. Halasz, J. Am. Chem. Soc. 141 (2019) 1212–1216.
- 89. K. J. Ardila-Fierro, J. G. Hernández, ChemSusChem 14 (2021) 2145–2162.
- 90. D. Braga, L. Maini, F. Grepioni, Chem. Soc. Rev. 42 (2013) 7638–7648.
- 91. K. Užarević, I. Halasz, T. Friščić, J. Phys. Chem. Lett. 6 (2015) 4129-4140.
- 92. D. E. Crawford, J. Casaban, Adv. Mater. 28 (2016) 5747–5754.
- 93. R. Rightmire, T. P. Hanusa, *Dalton Trans.* **45** (2016) 2352–2362.
- 94. S. Garcia-Manyes, A. E. M. Beedle, *Nat. Rev. Chem.* **1** (2017) 0083.
- 95. B. H. Bowser, S. L. Craig, *Polym. Chem.* **9** (2018) 3583–3593.
- 96. J. L. Howard, Q. Cao, D. L. Browne, *Chem. Sci.* 9 (2018) 3080–3094.
- 97. J. Andersen, J. Mack, Green Chem. 20 (2018) 1435–1443.
- 98. C. Bolm, J. G. Hernández, ChemSusChem 11 (2018) 1410–1420.
- 99. V. Štrukil, Synlett **29** (2018) 1281–1288.
- Eguaogie, J. S. Vyle, P. F. Conlon, M. A. Gîlea, Y. Liang, *Beilstein J. Org. Chem.* 14 (2018) 955–970.
- 101. D. Tan, F. García, Chem. Soc. Rev. 48 (2019) 2274–2292.
- A. Beillard, X. Bantreil, T.-X. Métro, J. Martinez, F. Lamaty, *Chem. Rev.* 119 (2019) 7529–7609.
- 103. Perona, P. Hoyos, A. Farrán, M. J. Hernaiz, Green Chem. 20 (2020) 5559–5583.
- 104. Porcheddu, E. Colacino, L. De Luca, F. Delogu, ACS Catal. 10 (2020) 8344-8394.
- 105. M. Pérez-Venegas, E. Juaristi, ACS Sustainable Chem. Eng. 8 (2020) 8881–8893.
- 106. T. Stolar, K. Užarević, CrystEngComm 22 (2020) 4511-4525.
- 107. S. Kaabel, T. Friščić, K. Auclair, ChemBioChem 21 (2020) 742-758.
- 108. W. Pickhardt, S. Grätz, L. Borchardt, Chem. Eur. J. 26 (2020) 12903–12911.
- 109. K. Kubota, H. Ito, *Trends Chem.* **2** (2020) 1066–1081.
- 110. R.T. O'Neil, R. Boulatov, Nat. Rev. Chem. 5 (2021) 148–167.

- A. P. Amrute, J. D. Bellis, M. Felderhoff, F. Schüth, *Chem. Eur. J.* 27 (2021) 6819– 6847.
- 112. F. Hajiali, T. Jin, G. Yang, M. Santos, E. Lam, A. Moores, *ChemSusChem* **15** (2022) e202102535.
- Bento, F. Luttringer, T. M. El Dine, N. Pétry, X. Bantreil, F. Lamaty, *Eur. J. Org. Chem.* (2022) e202101516.
- 114. N. Shan, F. Toda, W. Jones, Chem. Commun. (2002) 2372–2373.
- T. Friščić, A. V. Trask, W. Jones, W. D. S. Motherwell, Angew. Chem. Int. Ed. 45 (2006) 7546–7550.
- K. Užarević, V. Štrukil, C. Mottillo, P. A. Julien, A. Puškarić, T. Friščić, I. Halasz, Cryst. Growth Des. 16 (2016) 2342–2347.
- H. Kulla, M. Wilke, F. Fischer, M. Rollig, C. Maierhofer, F. Emmerling, *Chem. Commun.* 53 (2017) 1664–1667.
- S. Lukin, T. Stolar, I. Lončarić, I. Milanović, N. Biliškov, M. di Michiel, T. Friščić, I. Halasz *Inorg. Chem.* 59 (2020) 12200–12208.
- K. Užarević, N. Ferdelji, T. Mrla, P. A. Julien, B. Halasz, T. Friščić, I. Halasz, *Chem. Sci.* 9 (2018) 2525–2532.
- 120. J. Andersen, J. Brunemann, J. Mack, React. Chem. Eng. 4 (2019) 1229–1236.
- 121. N. Cindro, M. Tireli, B. Karadeniz, T. Mrla, K. Užarević, ACS Sustainable Chem. Eng. 7 (2019) 16301–16309.
- 122. T. Seo, N. Toyoshima, K. Kubota, H. Ito, J. Am. Chem. Soc. 143 (2021) 6165–6175.
- Takahashi, A. Hu, P. Gao, Y. Gao, Y. Pang, T. Seo, J. Jiang, S. Maeda, H. Takaya, K. Kubota, H. Ito, *Nat Commun* 12 (2021) 6691.
- 124. V. Štrukil, I. Sajko, Chem. Commun. 53 (2017) 9101–9104.
- 125. J. G. Hernández, Beilstein J. Org. Chem. 13 (2017) 1463–1469.
- 126. Bolm, J. G. Hernández, Angew. Chem. Int. Ed. 58 (2019) 3285–3299.
- 127. G.-F. Han, F. Li, Z.-W. Chen, C. Coppex, S.-J. Kim, H.-J. Noh, Z. Fu, Y. Lu, C. V. Singh, S. Siahrostami, Q. Jiang, J.-B. Baek, *Nat. Nanotechnol.* 16 (2021) 325–330.
- T. Friščić, I. Halasz, P. J. Beldon, A. M. Belenguer, F. Adams, S. A. J. Kimber, V. Honkimäki, R. E. Dinnebier, *Nature Chem* 5 (2013) 66–73.
- V. Ban, Y. Sadikin, M. Lange, N. Tumanov, Y. Filinchuk, R. Černý, N. Casati, *Anal. Chem.* 89 (2017) 13176–13181.

- G. I. Lampronti, A. A. L. Michalchuk, P. P. Mazzeo, A. M. Belenguer, J. K. M. Sanders, A. Bacchi, F. Emmerling, *Nat Commun* 12 (2021) 6134.
- Gracin, V. Štrukil, T. Friščić, I. Halasz, K. Užarević, Angew. Chem. Int. Ed. 53 (2014) 6193–6197.
- 132. Lukin, K. Užarević, I. Halasz, Nat Protoc 16 (2021) 3492–3521.
- 133. A. A. L. Michalchuk, F. Emmerling, Angew. Chem. Int. Ed. 61 (2022) e202117270.
- 134. K. Kubota, Y. Pang, A. Miura, H. Ito, *Science* **366** (2019) 1500–1504.
- 135. Schumacher, J. G. Hernández, C. Bolm, Angew. Chem. Int. Ed. 59 (2020) 16357–16360.
- 136. Huskić, C. B. Lennox, T. Friščić, Green Chem. 22 (2020) 5881–5901.
- Bolm, R. Mocci, C. Schumacher, M. Turberg, F. Puccetti, J. G. Hernández, Angew. Chem. Int. Ed. 57 (2018) 2423–2426.
- J. E. Elsila, N. M. Johnson, D. P. Glavin, J. C. Aponte, J. P. Dworkin, *Meteorit. Planet.* Sci. 56 (2021) 586–600.
- 139. S. Lamour, S. Pallman, M. Hass, O. Trapp, Life 9 (2019) 52.
- 140. M. Hass, S. Lamour, S. B. Christ, O. Trapp, Commun. Chem. 3 (2020) 140.
- Becker, J. Feldmann, S. Wiedemann, H. Okamura, C. Schneider, K. Iwan, A. Crisp, M. Rossa, T. Amatov, T. Carell, *Science* 366 (2019) 76–82.
- 142. TOPAS, version 4.2; Bruker-AXS: Karlsruhe, Germany, 2009.
- Stolar, S. Lukin, M. Etter, M. Rajić Linarić, K. Užarević, E. Meštrović, I. Halasz, *Chem. Commun.* 56 (2020) 13524–13527.
- 144. A. M. Atria, M. T. Garland, R. Baggio, Acta Cryst. C66 (2010) 0547.
- 145. E. Colacino, M. Carta, G. Pia, A. Porcheddu, P. C. Ricci, F. Delogu, ACS Omega 3 (2018) 9196 – 9209.
- A.W. Tricker, G. Samaras, K. L. Hebisch, M. J. Realff, C. Sievers, *Chem. Eng. J.* 382 (2020) 122954.
- 147. P.A. Cawood, C.J. Hawkesworth, B. Dhuime, GSA Bulletin 125 (2013) 14–32.
- L. Vugrin, M. Carta, S. Lukin, E. Meštrović, F. Delogu, I. Halasz, *ChemRxiv* (2022), DOI: 10.26434/chemrxiv-2022-cbp87
- 149. S. Lherminier, R. Planet, V. L. Vehel, G. Simon, L. Vanel, K. J. Måløy, O. Ramos, *Phys. Rev. Lett.* **122** (2019) 218501.
- 150. C. Catling, K. J. Zahnle, *Sci. Adv.* **6** (2020) eaax1420.

- 151. J. G. Forsythe, S.-S. Yu, I. Mamajanov, M. A. Grover, R. Krishnamurthy, F. M. Fernandez, N. V. Hud, *Angew. Chem. Int. Ed.* **54** (2015) 9871–9875.
- 152. A. D. McKee, M. Solano, A. Saydjari, C. J. Bennett, N. V. Hud, T. M. Orlando, *ChemBioChem* 19 (2018) 1913–1917.
- 153. Ferraris, D.W. Jones, J. Yerkess, Z. Kristallogr. 135 (1972) 240-252.
- 154. J. M. Hughes, M. Cameron, K. D. Crowley, American Mineralogist 74 (1989) 870–876.
- 155. L. Graf, American Mineralogist 46 (1961) 1283–1316.
- 156. K. Koyama, Y. Takeuchi, Z. Kristallogr. 145 (1977) 216–239.
- 157. J. C. A. Boeyens, V. V. H. Ichharam, Z. Kristallogr. 217 (2002) 9-10.
- 158. S. M. Richardson, J. W. Richardson, American Mineralogist 67 (1982) 69–75.
- 159. Bayliss, American Mineralogist 62 (1977) 1168–1172.
- 160. J. W. Gruner, Z. Kristallogr. 88 (1934) 412–419.
- 161. L. Levien, C. T. Prewitt, D. J. Weidner, American Mineralogist 65 (1980) 920–930.

§ 7. APPENDIX

```
macro PS {
macro BGR {
macro RBR { @ }
macro RBT { @ }
 macro REF {
no LIMIT warnings
 'do_errors
Auto T(100)
 iters 1000000000
           "C:\Documents and Settings\Tomislav\Desktop\Stolar\Self-assembly\Experimental data\D starting\PMF\D-NG-240_hires_merged.xy*
 xdd
 r_exp 4.267 r_exp_dash 6.560 r_wp 13.448 r_wp_dash 20.676 r_p 10.494 r_p_dash 18.728 weighted_Durbin_Watson 0.120 gof 3.152
bkg BGR 228.4923695 -19.69729777 -55.15388878 24.77688085 31.84849651 5.964291148
start_X 20
 LP_Factor ( 27)
Zero_Error (PS, 0.01515337979)
Rp 217.5
 Rs 217.5
 Simple_Axial_Model(PS, 3.827129369)
 ymin_on_ymax 0.001
la 0.6533 lo 0.709319 lh 0.2695
la 0.3467 lo 0.713574 lh 0.2795
x_calculation_step 0.002
str
CS_G(PS, 141.300175)
Strain_L(PS, 0.3083608399)
Strain_G(PS, 0.9368756804)
a PS 14.94544671
b PS 12.357385
c PS 6.755097821
 r_bragg 4.30247054
phase_name "hkl_Phase"
space_group Pbca
MVW( 1201.146, 1247.57631, 100.000`)
 scale @ 4.42742893e-005` max 0.1
'view_structure
Out_CIF_STR(D_anhydrate_PMF.cif)
 '2,6-diaminopurine
 rigid
point_for_site
                                                                                 uy 5.955 uz 2.034
uy 4.377 uz 1.168
uy 2.424 uz 0.392
                                         N3 ux 5.648
N4 ux 4.231
N5 ux 3.030
point_for_site
                                         H10 ux 2.395
N6 ux 4.280
N7 ux 6.149
                                                                                  uy 2.796
uy 0.812
uy 3.770
                                                                                                           uz -0.053
uz 1.287
                                                                                                           uz 2.518
                                                                                  uy 1.582
uy 4.656
uy 3.054
                                          N8 ux 6.764
                                                                                                           uz 2.928
                                          C5
C6
C7
                                                      ux 5.320
ux 4.029
                                                                                                           uz 1.908
uz 1.066
point_for_site
point_for_site
point_for_site
point_for_site
point_for_site
point_for_site
point_for_site
point_for_site
point_for_site
                                                       ux 3.230
                                                                                  uy 1.086
                                                                                                           uz 0.560
                                                   ux 2.676
ux 4.807
ux 5.920
                                                                                 uy 0.435
uy 2.053
uy 2.455
uy 6.189
                                                                                                           uz 0.194
uz 1.615
                                          H9
                                          C8
C9
                                                                                                          uz 2.374
                                       H6 ux 6.248
H5 ux 5.077
H7 ux 7.365
                                                                                                          uz 2.619
                                                                                                          uz 1.760
uz 3.478
                                                                                  uy 6.568
uy 1.853
point_for_site
                                        H8 ux 6.634
                                                                                 uy 0.733 uz 2.937
Rotate about axies( RBR 186.49150', RBR -32.56154', RBR -278.10057')
Translate(RBT 51416.47372', RBT 1108.52729', RBT -9028.24471')
prm Bl 1.60395' min 0 max 12
 '2,6-diaminopurine
'2,6-diaminopurine
site N3 num_posns 8 x 51416.90901 y 1108.95993 z
site N4 num_posns 8 x 51416.79509 y 1108.33624 z
site N5 num_posns 8 x 51416.6776 y 1108.63624 z
site H10 num_posns 8 x 51416.6776 y 1108.66860 z
site N6 num_posns 8 x 51416.76675 y 1109.02924 z
site N7 num_posns 8 x 51416.62581 y 1108.66784 z
site N7 num_posns 8 x 51416.62581 y 1108.066784 z
site N6 num_posns 8 x 51416.62587 y 1109.10300 z
site C5 num_posns 8 x 51416.62587 y 1109.02924 z
site C6 num_posns 8 x 51416.62307 y 1108.94062 z
site C6 num_posns 8 x 51416.57012 y 1108.76308 z
site C7 num_posns 8 x 51416.62339 y 1108.76308 z
site C9 num_posns 8 x 51416.64797 y 1108.71373 z
site C9 num_posns 8 x 51416.64797 y 1108.9097 z
site C9 num_posns 8 x 51416.67797 y 1109.01592 z
site H6 num_posns 8 x 51416.94584 y 1109.02475 z
                                                                                                                                       -9028.13074 occ N 1 beq
-9028.11405 occ N 1 beq
-9028.08603 occ N 1 beq
-9028.08670 occ H 1 beq
-9028.07469 occ N 1 beq
-9028.12005 occ N 1 beq
-9028.09106 occ N 1 beq
                                                                                                                                                                                                          =B1:
                                                                                                                                                                                                          =B1;
=B1;
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                                          =B1 :
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                                           =B1;
                                                                                                                                       -9028.09106 ccc N
-9028.1295 ccc C
-9028.09887 ccc C
-9028.07208 ccc C
-9028.06168 ccc H
-9028.09084 ccc C
-9028.10192 ccc C
-9028.15875 ccc H
                                                                                                                                                                                         1 beg
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                         l beq
l beq
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                         1 beq
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                        l beq
l beq
                                                                                                                                                                                                          =B1:
                                                                                                                                                                                                          =B1;
                                                                                                                                                                                         1 beq
                                                                                                                                                                                                           =B1;
 site H5 num_posns
site H7 num_posns
                                                8 x
8 x
8 x
                                                                51416.94584 y 1108.90533 z
51416.64680 y 1109.16610 z
51416.56795 y 1109.10011 z
                                                                                                                                       -9028.15090 occ H
-9028.11518 occ H
-9028.09058 occ H
                                                                                                                                                                                         l beg
                                                                                                                                                                                                          =B1;
 site H7 num_posns
site H8 num_posns
                                                                                                                                                                                         1 beq
                                                                                                                                                                                                          =B1 :
                                                                                                                                                                                        l beq
                                                                                                                                                                                                          =B1;
```

Figure 7.1. An example of input file that was used for the crystal structure solution *via* simulated annealing.

```
macro PS { @ }
macro BGR ( @ )
macro RBR { }
macro RBT {
macro REF { @ }
no_LIMIT_warnings
do_errors
'Auto_T(100)
iters 1000000000
xdd "C:\Documents and Settings\Tomislav\Desktop\Stolar\Self-assembly\Experimental data\D starting\PMF\D-NG-240 hires_merged.xy"
r_exp 5.806 r_exp_dash 10.879 r_wp 9.784 r_wp_dash 18.333 r_p 7.564 r_p_dash 17.718 weighted_Durbin_Watson 0.413 gof 1.685
bkg BGR 162.16077~_0.304336639 -70.2477652~_0.385290409 -20.8492304~_0.332689231 34.3419832~_0.318142462 -32.8780192~_0.306585502
'start_X 4
'finish_X 26
LP_Factor (90)
Zero_Error(PS, 0.01906`_0.00027)
Rp 217.5
Rs 217.5
Simple_Axial_Model(PS, 4.41109`_0.01691)
lam
ymin_on_ymax 0.001
la 0.6533 lo 0.709319 lh 0.2695
la 0.3467 lo 0.713574 lh 0.2795
x calculation step 0.002
str
CS_G(PS, 312.36483`_39.53892)
Strain_L(PS, 0.02753`_0.01404)
Strain_G(PS, 0.42073`_0.00812)
a PS 14.97675 0.00064
b PS 12.38205 0.00035
c PS 6.76696 0.00015
r_bragg 4.42506285
phase_name "DAP anhydrate"
 space_group Pbca
MVW ( 1201.146, 1254.88447`_0.0697825878, 100.000`_0.000)
scale @ 5.10703352e-005'_1.19e-007 max 0.1
  view_structure
Out CIF STR(DAP anhydrate.cif)
prm Bl 2.12341`_0.04694 min 0 max 12
'2,6-diaminopurine
                              8 x REF 0.90672' 0.00002 y REF 0.95805' 0.00004 z REF -0.14595' 0.00015 occ N
8 x REF 0.79403' 0.00006 y REF 0.83551' 0.00005 z REF -0.11532' 0.00025 occ N
8 x REF 0.65725' 0.00003 y REF 0.73391' 0.00004 z REF -0.08965' 0.00016 occ N
8 x REF 0.67734' 0.00012 y REF 0.66519' 0.00015 z REF -0.09221' 0.00040 occ H
site N3 num_posns
                                                                                                                                                                     1 beg
                                                                                                                                                                                =B1;
site N4 num_posns
                                                                                                                                                                     1 beq
                                                                                                                                                                                 =B1;
site N5
             num_posns
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1;
site H10 num_posns
                                                                                                                                                                     1
                                                                                                                                                                        beg
                                                                                                                                                                                 =B1;
site N6 num_posns
                                8 x REF
                                               0.56131
                                                            0.00004 y REF
                                                                                      0.86672 0.00006 z REF -0.07076
1.02736 0.00004 z REF -0.11758
                                                                                                                                            0.00031 occ N
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1:
site N7
              num_posns
                                8 x REF
                                               0.77056
                                                             0.00005
                                                                           y REF
                                                                                                                                            0.00036 occ N
                                                                                                                                                                                 =B1;
                                                                                                                                                                     1
                                                                                                                                                                        beq
                                                                                      1.09950 0.00004 z REF -0.09279
0.93907 0.00004 z REF -0.11915
0.82450 0.00004 z REF -0.09994
0.76386 0.00004 z REF -0.07225
site N8
              num posns
                                8 x REF
                                               0.62669
                                                             0.00003 V REF
                                                                                                                                            0.00013 occ N
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1:
                                8 x REF
                                               0.82212 0.00004 y REF
                                                                                                                                           0.00028 occ C
                                                                                                                                                                     1
                                                                                                                                                                        beq
site C5
              num_posns
                                                                                                                                                                                 =B1;
site C6
              num_posns
                                8 x REF
                                               0.70634 0.00003 V REF
                                                                                                                                            0.00012 occ C
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1;
site C7
                                 8 x REF
                                               0.56994
                                                            _0.00004 y REF
                                                                                                                                            0.00019 occ C
              num_posns
                                                                                                                                                                     1
                                                                                                                                                                        beq
                                                                                                                                                                                 =B1;
                                                                                      0.71412 0.00017 z REF -0.06151
0.90552 0.00004 z REF -0.08865
site H9
              num_posns
                                8 x REF
                                               0.51933
                                                            0.00011 y REF
                                                                                                                                            0.00048 occ H
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1;
                                               0.64639'_0.00003 y REF
0.68174'_0.00004 y REF
0.92940'_0.00013 y REF
                                                                                                                                            0.00019 occ C
site C8
              num_posns
                                8 x REF
                                                                                                                                                                     1
                                                                                                                                                                        beg
                                                                                                                                                                                 =B1;
                               8 x REF 0.68174 _0.00004 y REF 1.01193 _0.00014 z REF -0.09736 _0.00014 dcc C
8 x REF 0.92940 _0.00013 y REF 1.02325 _0.00016 z REF -0.13167 _0.00061 dcc H
8 x REF 0.94762 _0.00013 y REF 0.90384 _0.00015 z REF -0.17103 _0.00193 dcc H
8 x REF 0.64608 _0.00012 y REF 1.16376 _0.00014 z REF -0.08288 _0.00062 dcc H
8 x REF 0.56731 _0.00012 y REF 1.09463 _0.00015 z REF -0.07462 _0.00069 dcc H
site C9
              num_posns
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1:
site H6
                                                                                                                                                                                 =B1;
              num_posns
                                                                                                                                                                     1 beq
site H5
              num posns
                                                                                                                                                                     1 beg
                                                                                                                                                                                 =B1;
              num_posns
                                                                                                                                                                     1
site H7
                                                                                                                                                                                 =B1;
                                                                                                                                                                        beg
site H8
              num_posns
                                                                                                                                                                    1 beg
                                                                                                                                                                                =B1;
```

Figure 7.2. An example of input file that was used for final Rietveld refinement (part 1/2).

penalties_weighting_Kl 100

	37.0		1 0 0 0 1 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0
Distance_Restlain(N3	CS	, 1.3689, 1.30134 _0.00077 , 0.02, 1)
Distance Restrain(N3	H6	, 0.8791, 0.88117`_0.00213 , 0.02, 1)
Distance Restrain(N3	H5	, 0.9076, 0.92443 0.00260 , 0.02, 1)
Distance Restrain(N4	C5	1.3647 1.34978 0.00084 0.02 1)
Distance Resolutin(NT.4	00	1 242 1 224462 0 00005 0 02 1
Distance_Restrain(N4	00	, 1.342, 1.32446 0.00095 , 0.02, 1)
Distance_Restrain(N5	H10	, 0.8585, 0.85617`_0.00189 , 0.02, 1)
Distance Restrain(N5	C6	, 1.3759, 1.34297 0.00064 , 0.02, 1)
Distance Restrain(N5	C7	. 1.348. 1.36428 0.00068 . 0.02. 1)
Distance Destroin (MC	07	1 2061 1 20020 0 00002 0 02 1)
Distance_Restrain(140	01	, 1.2961, 1.28020 _0.00093 , 0.02, 1)
Distance_Restrain(NG	C8	, 1.388, 1.36715`_0.00081 , 0.02, 1)
Distance Restrain(N7	C5	, 1.3218, 1.33848` 0.00082 , 0.02, 1)
Distance Restrain(N7	C9	. 1.3625, 1.35083 0.00099 . 0.02, 1)
Distance Destroin(NO	00	1 2126 1 262501 0 00072 0 02 11
Distance_Restlain(NO	0.9	, 1.5156, 1.56250 _0.00075 , 0.02, 1)
Distance_Restrain(N8	H7	, 0.8509, 0.84966`_0.00191 , 0.02, 1)
Distance Restrain(N8	H8	, 0.8991, 0.89980` 0.00204 , 0.02, 1)
Distance Restrain(C6	C8	. 1.388. 1.34847 0.00066 . 0.02. 1)
Distance Pastrain (C7	HO	0 9736 0 97935 0 00190 0 02 1)
Distance Restrain(~		, 0.5/50, 0.5/555 _0.00150 , 0.02, 1)
Distance_Restrain(C8	69	, 1.4441, 1.42119 0.00074 , 0.02, 1)
Angle Restrain(C5	N3	H6	,120.07, 121.70099 0.14371 , 3, 0.001)
Angle Restrain (C5	N3	HS	121 93 122 65714 0 14584 3 0.001)
Deale Deatherin (NC	110		
Angle_Restlain(no	NS	15	,114.05, 115.40404_0.10500 , 5, 0.001)
Angle_Restrain(C5	N4	C6	,115.25, 114.10088`_0.06237 , 3, 0.001)
Angle Restrain (H10	N5	C6	,126.42, 126.03873 0.12397 , 3, 0.001)
Angle Restrain (H10	N5	C7	.125.76. 126.37069 0.12585 . 3. 0.001)
Angla Pastrain (C6	NE	07	107 81 107 58702 0 04357 3 0 0011
Angle_Restlain(Co	NO	-	,107.51, 107.35702 0.04337 , 3, 0.001)
Angle_Restrain(C7	N6	C8	,106.84, 104.76191 _0.05851 , 3, 0.001)
Angle_Restrain(C5	N7	C9	,117.66, 116.96296`_0.05517 , 3, 0.001)
Angle Restrain(C9	NS	H7	,122.67, 122.69860 0.13027 , 3, 0.001)
Angle Pastrain (C9	NR	HR	122 38 123 21866 0 12766 3 0 0011
Angle_Restlain(CS	NO	110	,122.30, 123.21009 _0.12750 , 3, 0.001)
Angle_Restrain(H/	NS	HS	,112.92, 112.94041 _0.18514 , 3, 0.001)
Angle_Restrain(N3	C5	N4	,119.19, 118.53509`_0.06267 , 3, 0.001)
Angle Restrain (N3	C5	N7	,114.61, 114.53959 0.05926 , 3, 0.001)
Angle Restrain (N4	C5	N7	126.17. 126.58024 0.06519 3. 0.001)
Angle Destroin (N4	00	NTE	100 07 100 06005 0 04004 0 0 0011
Angle_Restlain (N4	0	NS	,129.07, 129.26325 _0.04934 , 3, 0.001)
Angle_Restrain(N4	C6	C8	,126.53, 126.02251`_0.05036 , 3, 0.001)
	00	CB	105 683 104 70867: 0 03887 3 0 0011
Angle Restrain (N5	60	00	,105.005, 104./090/ 0.0599/ , 5, 0.001)
Angle_Restrain(N5 Angle_Restrain(N5	C7	NG	111.92. 111.58448 0.05317 . 3. 0.001)
Angle_Restrain(N5 Angle_Restrain(N5	C7	N6	,111.92, 111.58448_0.05397, 3, 0.001)
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5	C7 C7	N6 H9	,111.92, 111.58448°_0.05317 , 3, 0.001) ,125.15, 125.25939°_0.13129 , 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6	C7 C7 C7	N6 H9 H9	,111.92, 111.58488 _0.05317 , 3, 0.001) ,125.15, 125.25939 ^0.13129 , 3, 0.001) ,122.93, 123.15567 ^0.13368 , 3, 0.001)
Angle_Restrain (NS Angle_Restrain (NS Angle_Restrain (NS Angle_Restrain (N6 Angle_Restrain (N6	C7 C7 C7 C8	N6 H9 H9 C6	,111.92, 111.58448 _0.05317 , 3, 0.001) ,125.15, 125.25939 0.13129 , 3, 0.001) ,122.93, 123.15567 0.13368 , 3, 0.001) ,112.20, 111.35608 0.04904 , 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6	C7 C7 C7 C8 C8	N6 H9 H9 C6 C9	,101.92, 111.58448`_0.05397', 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332``0.05282, 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6	C7 C7 C7 C8 C8 C8	N6 H9 H9 C6 C9	,101.92, 111.58448 _0.05397 , 3, 0.001) ,111.92, 111.58448 _0.05317 , 3, 0.001) ,125.15, 125.25939 _0.13129 , 3, 0.001) ,122.93, 123.15567 _0.13368 , 3, 0.001) ,112.20, 111.35608 _0.04904 , 3, 0.001) ,132.067, 132.58332 _0.05282 , 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (C6	C7 C7 C7 C8 C8 C8 C8	N6 H9 H9 C6 C9 C9	,103,104,70367_0.05357, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (C6 Angle_Restrain (N7	C7 C7 C7 C8 C8 C8 C8 C9	N6 H9 H9 C6 C9 C9 N8	,111.92, 111.58448 _0.05397, 3, 0.001) ,125.15, 125.25939 0.13129, 3, 0.001) ,122.93, 123.15567 _0.13368, 3, 0.001) ,112.20, 111.35608 _0.04904, 3, 0.001) ,132.067, 132.58332 _0.05282, 3, 0.001) ,116.757, 116.06057 _0.04651, 3, 0.001) ,118.15, 119.05389 _0.05441, 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (C6 Angle_Restrain (N7 Angle_Restrain (N7	C7 C7 C7 C8 C8 C8 C8 C9 C9	N6 H9 H9 C6 C9 C9 N8 C8	,101.92, 111.58448`_0.05397', 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,122.0, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,118.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8	C7 C7 C7 C8 C8 C8 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8	,103,104,7036,_0.05357, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,118.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8	C7 C7 C7 C8 C8 C8 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8	,103,104,103,00,00,00,00,00,00,00,00,00,00,00,00,0
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8 Flatten (C7 H9 N5 H	C0 C7 C7 C8 C8 C8 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8	<pre>,103,104,7036*_0.03937, 3, 0.001) ,111.92,111.58448*_0.05317, 3, 0.001) ,125.15, 125.25939*_0.13129, 3, 0.001) ,122.93, 123.15567*_0.13368, 3, 0.001) ,112.20, 111.35608*_0.04904, 3, 0.001) ,112.067, 132.58332*_0.05282, 3, 0.001) ,116.757, 116.06057*_0.04651, 3, 0.001) ,116.15, 119.05389*_0.05441, 3, 0.001) ,121.47, 120.14365*_0.05211, 3, 0.001) ,120.37, 120.75929*_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448825*_0.000529990844, 10)</pre>
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8 Flatten (C7 H9 N5 H	C6 C7 C7 C8 C8 C8 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8	,103.083, 104.76348 _0.05357 , 3, 0.001) ,111.92, 111.58448 _0.05317 , 3, 0.001) ,125.15, 125.25939 ^0.13129 , 3, 0.001) ,122.93, 123.15567 _0.13368 , 3, 0.001) ,112.20, 111.35608 ^0.04904 , 3, 0.001) ,132.067, 132.58332 ^0.05282 , 3, 0.001) ,116.757, 116.06057 _0.04651 , 3, 0.001) ,118.15, 119.05389 ^0.05441 , 3, 0.001) ,121.47, 120.14365 ^0.05211 , 3, 0.001) ,120.37, 120.75929 ^0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 ^0.000529990844, 10)
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8 Flatten (C7 H9 N5 H.	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8	,103.063, 104.76365 _0.03937 , 3, 0.001) ,111.92, 111.58448 `0.05317 , 3, 0.001) ,125.15, 125.25939 `0.13129 , 3, 0.001) ,122.93, 123.15567 `0.13368 , 3, 0.001) ,112.20, 111.35608 `0.04904 , 3, 0.001) ,132.067, 132.58332 `0.05282 , 3, 0.001) ,132.067, 132.58332 `0.05282 , 3, 0.001) ,116.757, 116.06057 `0.04651 , 3, 0.001) ,118.15, 119.05389 `0.05441 , 3, 0.001) ,121.47, 120.14365 `0.05211 , 3, 0.001) ,120.37, 120.75929 `0.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925 `0.000529990844, 10)
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 10 N6	N6 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8	<pre>,103,104,7036*_0.03937, 3, 0.001) ,111.92,111.58448*_0.05317, 3, 0.001) ,125.15, 125.25939*_0.13129, 3, 0.001) ,122.93, 123.15567*_0.13368, 3, 0.001) ,112.20, 111.35608*_0.04904, 3, 0.001) ,122.067, 132.58332*_0.05282, 3, 0.001) ,116.757, 116.06057*_0.04651, 3, 0.001) ,116.15, 119.05389*_0.05441, 3, 0.001) ,121.47, 120.14365*_0.05211, 3, 0.001) ,120.37, 120.75929*_0.04930, 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925*_0.000529990844, 10) ate.pks)</pre>
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (C6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8 Flatten (C7 H9 N5 H. Create_2Th_Ip_file	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 (DAPan	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8	<pre>,103,104,70367_0.05357, 3, 0.001) ,111.92,111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,112.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,116.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks)</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`0	C6 C7 C7 C8 C8 C9 C9 C9 C9 C9 10 N6 (DAPan .00130	N6 H9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103,104,70367_0.05397, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,116.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 0</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical barmonic	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003, 104.7030*_0.03937, 3, 0.001) ,111.92, 111.58448*_0.05317, 3, 0.001) ,125.15, 125.25939*_0.13129, 3, 0.001) ,122.93, 123.15567*_0.13368, 3, 0.001) ,112.20, 111.35608*_0.04904, 3, 0.001) ,122.067, 132.58332*_0.05282, 3, 0.001) ,116.757, 116.06057*_0.04651, 3, 0.001) ,116.757, 116.06057*_0.04651, 3, 0.001) ,121.47, 120.14365*_0.05211, 3, 0.001) ,120.37, 120.75929*_0.04930, 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925*_0.000529990844, 10) ate.pks) 0</pre>
Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N5 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N6 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N7 Angle_Restrain (N8 Flatten (C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic.	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C6 C6 C0 hydr sh5	<pre>,103.003, 104.703480.05357 , 3, 0.001) ,111.92, 111.584480.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,112.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04651 , 3, 0.001) ,116.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 40 </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic.	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 Mhydr sh5 prm	<pre>,103.003, 104.703480.05357 , 3, 0.001) ,111.92, 111.584880.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,132.067, 132.583320.05282 , 3, 0.001) ,132.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04651 , 3, 0.001) ,118.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 4 0 {</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c00 1.0	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 Mydr hydr) min sh5 prm	<pre>,103.003, 104.05397, 0.0397, 3, 0.001) ,111.92, 111.58486`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,116.15, 119.05389`_0.05411, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 0 {</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c00 l.' y20 sh5 c20 0.'	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 C6 C9 C9 N8 C8 C8 C8 C8 C6 C6 C6 C0 hydr sh5 prm 0.0	<pre>,103.003,104.703480.05357 , 3, 0.001) ,111.92,111.584480.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,122.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04551 , 3, 0.001) ,116.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 4.0 { 2834 </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c00 1. y20 sh5_c20 0.	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 H9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C6 C 0 min sh5 prm 00 0_00	<pre>,103.003, 104.70348 _0.05357 , 3, 0.001) ,111.92, 111.58486 _0.05317 , 3, 0.001) ,125.15, 125.25939 ^0.13129 , 3, 0.001) ,122.93, 123.15567 _0.13368 , 3, 0.001) ,112.20, 111.35608 _0.04904 , 3, 0.001) ,122.07, 132.58332 ^0.05282 , 3, 0.001) ,116.757, 116.06057 _0.04651 , 3, 0.001) ,118.15, 119.05389 _0.05441 , 3, 0.001) ,121.47, 120.14365 _0.05211 , 3, 0.001) ,120.37, 120.75929 ^0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 ^0.000529990844, 10) ate.pks) 40 { 2834 2834 3383 </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c20 0.1 y20 sh5_c22 0.1	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	0.00 N6 H9 C6 C9 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003,104.703480.05357 , 3, 0.001) ,111.92,111.584480.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,122.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04651 , 3, 0.001) ,118.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 4 0 { 2834 3383 3161 </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic; sh_order 4 load si y00 !sh5_c00 1.; y20 sh5_c20 0.; y22p sh5_c22p -0.; y40 sh5_c40 -0.;	C7 C7 C7 C8 C8 C9 C9 C9 C9 10 N6 (DAPan .00130 s_hk1 h_C1j_ 00000 81573` 42726` 39430`	N6 N6 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003,104.703480.05357 , 3, 0.001) ,111.92,111.584880.05317 , 3, 0.001) ,125.15,125.259390.13129 , 3, 0.001) ,122.93,123.155670.13368 , 3, 0.001) ,112.20,111.356080.04904 , 3, 0.001) ,122.067,132.583320.05282 , 3, 0.001) ,116.757,116.060570.04651 , 3, 0.001) ,116.15,119.053890.05441 , 3, 0.001) ,121.47,120.143650.05211 , 3, 0.001) ,120.37,120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 0 { 2834 3383 3161</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	C6 C6 C M6 H9 H9 C6 C9 C9 C9 C8 C8 C8 C6 C min sh5 prm _0.0 _0.0 _0.0 _0.0	<pre>,103,104,70346 _0.03357 , 3, 0.001) ,111.92,111.58486 _0.05317 , 3, 0.001) ,125.15, 125.25939 ^0.13129 , 3, 0.001) ,122.93, 123.15567 _0.13368 , 3, 0.001) ,112.20, 111.35608 ^0.04904 , 3, 0.001) ,132.067, 132.58332 ^0.05282 , 3, 0.001) ,132.067, 132.58332 ^0.05441 , 3, 0.001) ,116.15, 119.05389 ^0.05411 , 3, 0.001) ,121.47, 120.14365 ^0.05211 , 3, 0.001) ,120.37, 120.75929 ^0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 ^0.000529990844, 10) ate.pks) 0 { 2884 3383 3161 5395 </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	0.00 H9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C9 Min sh5 prm -0.00 -0.00 -0.00	<pre>,103.003, 104.703480.05357 , 3, 0.001) ,111.92, 111.584880.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,122.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04651 , 3, 0.001) ,118.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 4 0 { 2834 3383 3161 5395 1108}</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	N6 N6 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003,104.703480.05357 , 3, 0.001) ,111.92,111.584880.05317 , 3, 0.001) ,125.15,125.259390.13129 , 3, 0.001) ,122.93,123.155670.13368 , 3, 0.001) ,112.20,111.356080.04904 , 3, 0.001) ,112.067,132.583320.05282 , 3, 0.001) ,116.757,116.060570.04651 , 3, 0.001) ,116.757,116.060570.04651 , 3, 0.001) ,121.47,120.143650.05211 , 3, 0.001) ,120.37,120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 0 { 2834 3383 3161 5395 1108} </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 10 N6 (DAPan .00130 s_hkl h_Cij_ 00000 81573` 42726` 39430` 74314` 12781` / C8	C6 C9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C9 N8 C8 C9 N8 C8 C9 N8 C9 C9 N8 C9 C9 N8 C9 C9 N8 C9 C9 N8 C9 C9 N8 C6 C9 C9 N8 C9 C9 C9 N8 C6 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 N8 C8 C8 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 C9 N8 C8 C9 C9 N8 C8 C9 C9 N8 C8 C8 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	<pre>,103.003, 104.70346 _0.03357 , 3, 0.001) ,111.92, 111.58446 _0.05317 , 3, 0.001) ,125.15, 125.25939 _0.13129 , 3, 0.001) ,122.93, 123.15567 _0.13368 , 3, 0.001) ,112.20, 111.35608 _0.04904 , 3, 0.001) ,112.20, 111.35608 _0.05282 , 3, 0.001) ,116.757, 116.06057 _0.04651 , 3, 0.001) ,116.15, 119.05389 _0.05441 , 3, 0.001) ,118.15, 119.05389 _0.05441 , 3, 0.001) ,120.37, 120.75929 _0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 _0.000529990844, 10) ate.pks) 0 { 28234 3383 3161 5395 1108} </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load s1 y00 !sh5_c00 1. y20 sh5_c20 0. y20 sh5_c22p -0. y40 sh5_c42p -0. y44p sh5_c44p -0.	C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	CG C9 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003, 104.70307_0.03937, 3, 0.001) ,111.92, 111.58486`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,132.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,118.15, 119.05389`_0.05411, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 4 0 { 2834 3363 3161 5395 1108} </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 10 N6 (DAPan .00130 s_hkl h_Cij_ 00000 81573` 42726` 39430` 74314` 12781` / C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	C C C C C C C C C C C C C C C C C C C	<pre>,103.003, 104.703480.05317 , 3, 0.001) ,111.92, 111.584880.05317 , 3, 0.001) ,125.15, 125.259390.13129 , 3, 0.001) ,122.93, 123.155670.13368 , 3, 0.001) ,112.20, 111.356080.04904 , 3, 0.001) ,122.067, 132.583320.05282 , 3, 0.001) ,116.757, 116.060570.04651 , 3, 0.001) ,118.15, 119.053890.05441 , 3, 0.001) ,121.47, 120.143650.05211 , 3, 0.001) ,120.37, 120.759290.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.0007294489250.000529990844, 10) ate.pks) 4 0 { 2834 3383 3161 5395 1108} </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c20 0.1 y20 sh5_c20 0.1 y20 sh5_c20 0.1 y22p sh5_c22p -0. y40 sh5_c40 -0.1 y42p sh5_c44p -0.1 lor_fwhm = sh5 p5 1 xdd_out Plot-data-1	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 10 N6 (DAPan h_Cij_ 000130 s_hkl h_Cij_ 000000 81573` 42726` 39430` 74314` 12781` / Cos(DAPanh	C C C C C C C C C C C C C C C C C C C	<pre>,103.003, 104.7030; 0.03337, 3, 0.001) ,111.92, 111.58486`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 0 { 2834 3383 3161 5395 1108} te.txt load out_record out_fmt out_eqn</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	CG C9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C9 MN8 C8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C6 C9 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C8 C9 MN8 C9 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C9 MN8 C8 C9 M10 M10 M10 M10 M10 M10 M10 M10 M10 M10	<pre>,103.003, 104.70307_0.03337, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,122.07, 132.58332`_0.05282, 3, 0.001) ,118.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 00 { 2834 3383 3161 5395 1108} te.txt load out_record out_fmt out_eqn</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	CG C9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003, 104.70307_0.05337, 3, 0.001) ,111.92, 111.58486`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,118.15, 119.05389`_0.05441, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 4 0 {</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046`_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c00 1. y20 sh5_c20 0. y22p sh5_c22p -0. y40 sh5_c40 -0. y42p sh5_c42p -0. y44p sh5_c44p -0. lor_fwhm = sh5 p5 xdd_out Plot-data-1 { "%11.5f" = X; "%11.5f" = Ych	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 10 N6 (DAPan h_Cij_ 00000 81573 42726 39430 74314 12781 / Cos(DAPanh s;	C C C C C C C C C C C C C C C C C C C	<pre>,103.003, 104.7030*0.03937, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,112.20, 111.35608`_0.04904, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 0 { 2834 3383 3161 5395 1108 te.txt load out_record out_fmt out_eqn </pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	C6 C9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C9 MN8 C8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C8 C9 M10 M10 M10 M10 M10 M10 M10 M10 M10 M10	<pre>,103.003, 104.70300.0337 , 3, 0.001) ,111.92, 111.58486 `_0.05317 , 3, 0.001) ,125.15, 125.25939 `_0.13129 , 3, 0.001) ,122.93, 123.15567 `_0.13368 , 3, 0.001) ,112.20, 111.35608 `_0.04904 , 3, 0.001) ,122.07, 116.06057 `_0.04651 , 3, 0.001) ,118.15, 119.05389 `_0.05441 , 3, 0.001) ,121.47, 120.14365 `_0.05211 , 3, 0.001) ,120.37, 120.75929 `_0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 `_0.000529990844, 10) ate.pks) 00 { 2834 3383 3161 5395 1108} te.txt load out_record out_fmt out_eqn</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046°_0 spherical_harmonic. sh_order 4 load si y00 !sh5_c00 1.0 y20 sh5_c20 0.0 y22p sh5_c22p -0.0 y40p sh5_c42p -0.0 y44p sh5_c42p -0.0 y44p sh5_c44p -0.0 lor_fwhm = sh5 p5 0 xdd_out Plot-data-0 { " %11.5f" = X; " %11.5f" = Yca	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	CG C9 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003, 104.70300.03377, 3, 0.001) ,111.92, 111.58448 `_0.05317 , 3, 0.001) ,125.15, 125.25939 `_0.13129 , 3, 0.001) ,122.93, 123.15567 `_0.13368 , 3, 0.001) ,112.20, 111.35608 `_0.04904 , 3, 0.001) ,122.067, 132.58332 `_0.05282 , 3, 0.001) ,116.757, 116.06057 `_0.04651 , 3, 0.001) ,121.47, 120.14365 `_0.05211 , 3, 0.001) ,120.37, 120.75929 `_0.04930 , 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925 `_0.000529990844, 10) ate.pks) 4 0 {</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restra	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 10 N6 (DAPan h_Cij_ 000000 81573 42726 39430 74314 12781 / Cos(DAPanh s; 1c; s-Ycal	C C C C C C C C C C C C C C C C C C C	<pre>,103.003, 104.7030^_0.03377, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,132.067, 132.58332`_0.04651, 3, 0.001) ,116.757, 116.06057`_0.04651, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 10 { te.txt load out_record out_fmt out_eqn } te.txt load out_record out_fmt out_eqn </pre>
<pre>Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H Create_2Th_Ip_file prm p5 0.03046[*]_0 spherical_harmonic: sh_order 4 load si y00 !sh5_c00 1. y20 sh5_c20 0. y22p sh5_c22p -0. y40 sh5_c40 -0. y42p sh5_c42p -0. y44p sh5_c44p -0. lor_fwhm = sh5 p5 . xdd_out Plot-data-1 { " %11.5f" = X; " %11.5f" = Yob: " %11.</pre>	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	C6 C9 H9 C6 C9 C9 N8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C9 MN8 C8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C8 C9 MN8 C9 C9 MN8 C8 C9 MN8 C9 C9 MN8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C8 C8 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C8 C8 C9 MN8 C8 C8 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C9 C9 MN8 C9 C9 MN8 C9 C9 MN8 C8 C8 C9 MN8 C8 C8 C9 M10 M10 M10 M10 M10 M10 M10 M10 M10 M10	<pre>,103.003, 104.70300.0337 , 3, 0.001) ,111.92, 111.58446 `_0.05317 , 3, 0.001) ,125.15, 125.25939 `_0.13129 , 3, 0.001) ,122.93, 123.15567 `_0.13368 , 3, 0.001) ,112.20, 111.35608 `_0.04904 , 3, 0.001) ,112.20, 111.35608 `_0.04932 , 3, 0.001) ,116.757, 116.06057 `_0.04651 , 3, 0.001) ,118.15, 119.05389 `_0.05441 , 3, 0.001) ,121.47, 120.14365 `_0.05211 , 3, 0.001) ,120.37, 120.75929 `_0.04930 , 3, 0.001) 38 C9 N4 N7 C5 N3, 0, 0.000729448925 `_0.000529990844, 10) ate.pks) 00 { [</pre>
Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N5 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N6 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N7 Angle_Restrain(N8 Flatten(C7 H9 N5 H. Create_2Th_Ip_file prm p5 0.03046°_0 spherical_harmonic. sh_order 4 load s1 y00 !sh5_c00 l. y20 sh5_c20 0. y20 sh5_c20 0. y22p sh5_c22p -0. y40 sh5_c42p -0. y44p sh5_c42p -0. lor_fwhm = sh5 p5 xdd_out Plot-data-1 { " %ll.5f" = X; " %ll.5f" = Yca. " %ll.5f\n" = Ycb. }	C7 C7 C7 C7 C8 C8 C9 C9 C9 C9 C9 C9 C9 C9 C9 C9	CG C9 H9 C6 C9 C9 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8 C8	<pre>,103.003, 104.70307_0.05337, 3, 0.001) ,111.92, 111.58448`_0.05317, 3, 0.001) ,125.15, 125.25939`_0.13129, 3, 0.001) ,122.93, 123.15567`_0.13368, 3, 0.001) ,122.067, 132.58332`_0.05282, 3, 0.001) ,118.15, 119.05389`_0.04651, 3, 0.001) ,121.47, 120.14365`_0.05211, 3, 0.001) ,120.37, 120.75929`_0.04930, 3, 0.001) 8 C9 N4 N7 C5 N3, 0, 0.000729448925`_0.000529990844, 10) ate.pks) 4 0 {</pre>

Figure 7.3. An example of input file that was used for final Rietveld refinement (part 2/2).



Figure 7.4. ORTEP-3 drawing of asymmetric unit of **2** (D:U cocrystal hydrate) with atom numbering scheme. Displacement ellipsoids are drawn for the probability of 50 % and hydrogen atoms are shown as spheres of arbitrary radii. Disordered water molecule has been omitted.

Compound	D:U cocrystal hydrate
Empirical formula	$C_9H_{12}N_8O_3$
Formula wt. / g mol ⁻¹	280.27
Colour	colourless
Crystal dimensions / mm	0.18 x 0.12 x 0.07
Space group	C 2/c
<i>a</i> / Å	10.6851(8)
b / Å	16.5477(10)
<i>c</i> / Å	14.2161(12)
α / °	90
β / °	102.930(8)
γ/°	90
Z	8
$V/\text{\AA}^3$	2449.9(3)
D_{calc} / g cm ⁻³	1.520
λ/Å	1.54719 (CuKα)
μ / mm ⁻¹	1.017
Θ range / °	5.02 - 76.84
T / \mathbf{K}	293(2)
Diffractometer type	Xcalibur Nova
	11 < h < 13;
Range of h, k, l	-20 < k < 17;
	-16 < <i>l</i> < 17
Reflections collected	5096
Independent reflections	2498
Observed reflections	1705
$(I \ge 2\sigma)$	1/85
Absorption correction	Multi-scan
T_{\min}, T_{\max}	0.4930; 1.0000
R _{int}	0.0380
R (F)	0.0673
$R_w (F^2)$	0.2133
Goodness of fit	1.012
H atom treatment	Mixed
No. of parameters	210
No. of restraints	12
$\Delta \rho_{\text{max}}$, $\Delta \rho_{\text{min}}$ (eÅ ⁻³)	0.278; - 0.244

Table 7.1. Crystallographic, data collection and refinement data for single crystal X-ray diffraction data of 2 (D:U cocrystal hydrate).



Figure 7.5. TGA curve of 2 (D:U cocrystal hydrate).



Figure 7.6. ORTEP-3 drawing of asymmetric unit of **10** (D:T 1:2 cocrystal hydrate). Displacement ellipsoids are drawn for the probability of 50 % and hydrogen atoms are shown as spheres of arbitrary radii.

Table 7.2. Crystallographic, data collection and refinement data for single crystal X-ray diffraction data of **10** (D:T 1:2 cocrystal hydrate).

Empirical formula	C ₃₀ H ₄₄ N ₂₀ O ₁₂
Formula weight	876.85
Temperature/K	169.98(10)
Crystal system	monoclinic
Space group	Pc
a/Å	17.3694(5)
b/Å	16.9996(4)
c/Å	6.6652(2)
α/°	90
β/°	93.240(3)
$\gamma^{/\circ}$	90
Volume/Å ³	1964.91(9)
Z	2
$\rho_{calc}g/cm^3$	1.482
µ/mm ⁻¹	1.001
F(000)	920.0
Crystal size/mm ³	$? \times ? \times ?$
Radiation	Cu Ka ($\lambda = 1.54184$)
2O range for data collection/	° 7.282 to 132.944
Index ranges	-20 $\leq h \leq$ 20, 0 $\leq k \leq$ 20, -7 $\leq l \leq$ 7
Reflections collected	6826
Independent reflections	6826 [R _{int} = ?, R _{sigma} = 0.1163]
Data/restraints/parameters	6826/0/574
Goodness-of-fit on F ²	1.059
Final R indexes [I>=2σ (I)]	$R_1 = 0.0783, wR_2 = 0.2256$
Final R indexes [all data]	$R_1 = 0.1039, wR_2 = 0.2405$
Largest diff. peak/hole / e Å	30.57/-0.29
Flack parameter	0.1(4)

	r
molecular formula	$C_{10} H_{12} N_8 O_2$
formula weight (g mol ⁻¹)	276.28
space group	ΡĪ
<i>a</i> (Å)	8.77213 (67)
<i>b</i> (Å)	11.07778 (52)
<i>c</i> (Å)	12.6355 (11)
α (deg)	95.7660 (37)
β (deg)	82.2760 (41)
γ (deg)	89.5282 (32)
$V(\text{\AA}^3)$	1210.27 (15)
Z	4
temperature (K)	295
radiation type	Cu Ka
wavelength (Å)	1.5406
R_p	0.032
R_{wp}	0.047
GOF	7.61
$R(F^2)$	0.020

 Table 7.3. Crystallographic data for 6 (D:T cocrystal anhydrate).



Figure 7.7. PXRD patterns of slurrying A and H in water using different reaction conditions.



Figure 7.8. PXRD patterns of NG and LAG A and H at different temperatures.



Figure 7.9. PXRD patterns of A and H system including DSC heating at 300 °C.

molecular formula	C ₁₀ H ₉ N ₉ O
formula weight (g mol ⁻¹)	271.26
space group	$P2_{1}/n$
<i>a</i> (Å)	12.00132 (53)
<i>b</i> (Å)	25.7944 (12)
<i>c</i> (Å)	3.64521 (11)
α (deg)	90
β (deg)	85.3997 (38)
γ (deg)	90
$V(\text{\AA}^3)$	1124.800 (79)
Z	4
temperature (K)	295
radiation type	Cu Ka
wavelength (Å)	1.5406
R_p	0.034
R_{wp}	0.045
GOF	7.87
$R(F^2)$	0.019

 Table 7.4. Crystallographic data for 8 (A:H cocrystal anhydrate).
molecular formula	$C_{10}H_{13}N_9O_3$
formula weight (g mol ⁻¹)	307.29
space group	$P2_{1}/n$
<i>a</i> (Å)	22.8501 (16)
<i>b</i> (Å)	16.8210 (11)
<i>c</i> (Å)	3.60152 (19)
α (deg)	90
β (deg)	86.7812 (70)
γ (deg)	90
$V(\text{\AA}^3)$	1382.10 (15)
Z	4
temperature (K)	295
radiation type	Μο Κα
wavelength (Å)	0.71073
R_p	0.052
R_{wp}	0.069
GOF	1.81
$R(F^2)$	0.017

 Table 7.5. Crystallographic data for 9 (A:H cocrystal hydrate).



Figure 7.10. PXRD patterns for D and X system.



Figure 7.11. PXRD patterns for H and U system.



Figure 7.12. PXRD patterns for H and C system.



Figure 7.13. PXRD patterns for G and U system.



Figure 7.14. PXRD patterns for D, U, and A system.



Figure 7.15. PXRD patterns for D, T, and A system.



Figure 7.16. PXRD patterns for D, U, and T system.



Figure 7.17. PXRD patterns for A, H, and T system.



Figure 7.18. PXRD patterns for A, H, and U system.



Figure 7.19. IP-HPLC chromatogram for milling Gly with SiO₂ at 100 °C.

XXXIX



Figure 7.20. IP-HPLC chromatogram for milling Gly with montmorillonite at 100 °C.



Figure 7.21. IP-HPLC chromatogram for milling Gly with mica at 100 °C.

XL



Figure 7.22. IP-HPLC chromatogram for milling Gly with TiO₂ at RT in the presence of water.



Figure 7.23. IP-HPLC chromatogram for milling Gly with TiO_2 at 130 °C in the presence of water.

Tomislav Stolar



Figure 7.24. IP-HPLC chromatogram for milling Ala with TiO₂ at 130 °C.



Figure 7.25. IP-HPLC chromatogram for milling Gly and Ala with TiO₂ at 100 °C.



Figure 7.26. IP-HPLC chromatogram for milling Gly with SiO₂ at room temperature for 96 h and at 30 Hz.



Figure 7.27. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 96 h and at 30 Hz.



Figure 7.28. IP-HPLC chromatogram for milling Gly with SiO₂ at room temperature for 48 h and at 30 Hz.



Figure 7.29. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 48 h and at 30 Hz.



Figure 7.30. IP-HPLC chromatogram for milling Gly with SiO₂ at room temperature for 96 h and at 15 Hz.



Figure 7.31. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 96 h and at 15 Hz.



Figure 7.32. IP-HPLC chromatogram for milling Gly with SiO₂ at room temperature for 48 h and at 15 Hz.



Figure 7.33. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 48 h and at 15 Hz.



Figure 7.34. IP-HPLC chromatogram for milling Gly with SiO_2 at room temperature for 96 h and at 5 Hz.



Figure 7.35. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 96 h and at 5 Hz.



Figure 7.36. IP-HPLC chromatogram for milling Gly with SiO₂ at room temperature for 48 h and at 5 Hz.



Figure 7.37. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ at room temperature for 48 h and at 5 Hz.



Figure 7.38. Photograph showing three stones picked up in front of Ruđer Bošković Institute by Prof. José G. Hernández. These stones were used as a milling media for milling Gly with TiO_2 for 18h at 20 Hz and ambient temperature. Gly oligomers up to Gly₄ were formed in this way.



Figure 7.39. IP-HPLC chromatogram for milling Gly with SiO_2 in N_2 atmosphere at room temperature for 16 h and at 30 Hz. Milling under N_2 atmosphere was achieved by loading the jar in glovebox under N_2 atmosphere, closing it tightly and opening only before the chromatographic and mass spectrometry analysis.



Figure 7.40. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO_2 in N_2 atmosphere at room temperature for 16 h and at 30 Hz. Milling under N_2 atmosphere was achieved by loading the jar in glovebox under N_2 atmosphere, closing it tightly and opening only before the chromatographic and mass spectrometry analysis.



Figure 7.41. IP-HPLC chromatogram for milling Gly with SiO_2 in CO_2 atmosphere at room temperature for 16 h and at 30 Hz. Milling under CO_2 atmosphere was achieved by loading the jar containing the reaction mixture in glovebox with dry ice (solid CO_2) to the top, waiting for 30 s and closing it tightly. The jar was opened only before the chromatographic and mass spectrometry analysis. Note: due to high pressure inside, the jar had to be opened with special equipment and slowly vented.



Figure 7.42. LC/MS (Triple quadrupole) chromatograms for milling Gly with SiO₂ in CO₂ atmosphere at room temperature for 16 h and at 30 Hz.



Figure 7.43. Rietveld refinement of analcime PXRD pattern and analcime CIF taken from the literature.¹⁵³



Figure 7.44. PXRD comparison of apatite sample and simulated from the CIF.¹⁵⁴



Figure 7.45. Rietveld refinement of calcite PXRD pattern and calcite CIF taken from the literature.¹⁵⁵



Figure 7.46. Rietveld refinement of clinoptilolite PXRD pattern and clinoptilolite CIF taken from the literature.¹⁵⁶



Figure 7.47. Rietveld refinement of gypsum PXRD pattern and gypsum CIF taken from the literature.¹⁵⁷



Figure 7.48. Rietveld refinement of magnesite PXRD pattern and magnesite CIF taken from the literature.¹⁵⁵



Figure 7.49. Rietveld refinement of muskovite PXRD pattern and muskovite CIF taken from the literature.¹⁵⁸



Figure 7.50. PXRD comparison of pyrite sample and simulated from the CIF.¹⁵⁹



Figure 7.51. Rietveld refinement of talc PXRD pattern and talc CIF taken from the literature.¹⁶⁰



Figure 7.52. Rietveld refinement of quartz PXRD pattern and quartz CIF taken from the literature.¹⁶¹



Figure 7.53. IP-HPLC chromatogram for milling Gly with pyrite at room temperature for 16 h and at 30 Hz.



Figure 7.54. LC/MS (Triple quadrupole) chromatograms for milling Gly with pyrite at room temperature for 16 h and at 30 Hz.



Figure 7.55. IP-HPLC chromatogram for milling Gly with gypsum at room temperature for 16 h and at 30 Hz.



Figure 7.56. LC/MS (Triple quadrupole) chromatograms for milling Gly with gypsum at room temperature for 16 h and at 30 Hz.



Figure 7.57. IP-HPLC chromatogram for milling Gly with talc at room temperature for 16 h and at 30 Hz.



Figure 7.58. LC/MS (Triple quadrupole) chromatograms for milling Gly with talc at room temperature for 16 h and at 30 Hz.


Figure 7.59. IP-HPLC chromatogram for milling Gly with magnesite at room temperature for 16 h and at 30 Hz.



Figure 7.60. LC/MS (Triple quadrupole) chromatograms for milling Gly with magnesite at room temperature for 16 h and at 30 Hz.

LXXI



Figure 7.61. IP-HPLC chromatogram for milling Gly with quartz at room temperature for 16 h and at 30 Hz.



Figure 7.62. IP-HPLC chromatogram for milling Gly with clinoptilolite at room temperature for 16 h and at 30 Hz.



Figure 7.63. LC/MS (Triple quadrupole) chromatograms for milling Gly with clinoptilolite at room temperature for 16 h and at 30 Hz.



Figure 7.64. IP-HPLC chromatogram for milling Gly with apatite at room temperature for 16 h and at 30 Hz.



Figure 7.65. LC/MS (Triple quadrupole) chromatograms for milling Gly with apatite at room temperature for 16 h and at 30 Hz.

0.014



LXXVII



Figure 7.66. IP-HPLC chromatogram for milling Gly with analcime at room temperature for 16 h and at 30 Hz.



Figure 7.67. LC/MS (Triple quadrupole) chromatograms for milling Gly with analcime at room temperature for 16 h and at 30 Hz.



Figure 7.68. IP-HPLC chromatogram for milling Gly with calcite at room temperature for 16 h and at 30 Hz.

10.00 Minutes

12.00

14.00

16.00

18.00

20.00

8.00

6.00

2.00

0.00

4.00

LXXIX



Figure 7.69. LC/MS (Triple quadrupole) chromatograms for milling Gly with calcite at room temperature for 16 h and at 30 Hz.



Figure 7.70. IP-HPLC chromatogram for milling Gly with muskovite at room temperature for 16 h and at 30 Hz.



Figure 7.71. LC/MS (Triple quadrupole) chromatograms for milling Gly with muskovite at room temperature for 16 h and at 30 Hz.



Figure 7.72. IP-HPLC chromatogram for milling glycolic acid with SiO₂ at room temperature for 16 h and at 30 Hz.

3.487

0.60

0.50-

0.40

₽ 0.30-





Figure 7.73. IP-HPLC chromatogram for milling glycine and glycolic acid with SiO₂ at room temperature for 16 h and at 30 Hz.

§ 8. CURRICULUM VITAE

tomislav.stolar@gmail.com
0000-0002-9824-4462
Tomislav Stolar/@StolarTomislav
Research assistant, Laboratory for Applied and
Sustainable Chemistry, Ruđer Bošković Institute,
Zagreb (Croatia)
External associate, PLIVA Croatia, R&D – Teva
Active Pharmaceutical Ingredients Physical
Characterization Department, Zagreb (Croatia)
Scientific research on a voluntary basis, Ruđer
Bošković Institute, Zagreb (Croatia)

Education:

Tomislav obtained his Bachelor's (2014.) and Master's degrees (2016.) from the Department of Chemistry, Faculty of Science, University of Zagreb (Croatia). He is currently enrolled in PhD studies at the same institution.

<u>**Publications**</u> (five selected publications, 16 in total, eight as first author, one as corresponding author, 486 citations on Google Scholar):

- 1) T. Stolar et al., Angew. Chem. Int. Ed. 2021, 60, 12727–12731.
- 2) T. Stolar et al., ACS Appl. Mater. Interfaces 2021, 13, 3070–3077.
- 3) T. Stolar et al., Chem. Commun. 2020, 56, 13524–13527.
- 4) T. Stolar, K. Užarević, CrystEngComm 2020, 22, 4511-4525.
- 5) T. Stolar et al., ACS Sustainable Chem. Eng. 2019, 7, 7102–7110.

Selected conferences & meetings

3) T. Stolar *et al.*, "The Effects of Geological and Environmental Conditions on the Rise of Prebiotic Peptides on Mineral Surfaces", 2022 Astrobiology Science Conference –**oral online presentation** (**11** + **4 min**)

2) T. Stolar *et al.*, "Mechanochemical prebiotic peptide bond formation in the absence of water", AbGradCon online event, 2021 -oral presentation (12 + 3 min)

1) T. Stolar *et al.*, "Importance of solid state in prebiotic chemistry studies", Solid State Science & Research 2019, Zagreb, Croatia - **award for distinguished oral presentation for young researcher (15 min)**

Professional services, independence, and leadership

Peer-reviewing for ChemSusChem and ACS Sustainable Chemistry & Engineering journals.

Projects as PI: CERIC-ERIC project 20212028 'SS-NMR for probing hydrogen-bonding interactions in nucleobase cocrystals'. 168 hours collection at the 600 MHz solid-state NMR spectrometer MAGIC, the Slovenian NMR Center, Ljubljana, Slovenia.

Voted representative of the Department of Physical Chemistry graduate students and post-docs (35 employees) at Ruđer Bošković Institute's "Council of graduate students and post-docs" as well as "Council of the Department of Physical Chemistry" from 2019.–2021.