

62Ni(n, g) and 63Ni(n, g) cross sections measured at the n_TOF facility at CERN

(n_TOF Collaboration) Lederer, C.; Massimi, C.; Berthoumieux, E.; Colonna, N.; Dressler, R.; Guerrero, C.; Günsing, F.; Käppeler, F.; Kível, N.; Pignatari, M.; ...

Source / Izvornik: **Physical Review C - Nuclear Physics, 2014, 89**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1103/PhysRevC.89.025810>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:217:412823>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom](#).

Download date / Datum preuzimanja: **2024-07-11**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



$^{62}\text{Ni}(n,\gamma)$ and $^{63}\text{Ni}(n,\gamma)$ cross sections measured at the n_TOF facility at CERN

C. Lederer,^{1,2,*} C. Massimi,³ E. Berthoumieux,⁴ N. Colonna,⁵ R. Dressler,⁶ C. Guerrero,⁷ F. Gunsing,⁴ F. Käppeler,⁸ N. Kivel,⁶ M. Pignatari,⁹ R. Reifarth,² D. Schumann,⁶ A. Wallner,¹ S. Altstadt,² S. Andriamonje,⁷ J. Andrzejewski,¹⁰ L. Audouin,¹¹ M. Barbagallo,⁵ V. Bécères,¹² F. Bečvář,¹³ F. Belloni,¹⁴ B. Berthier,¹¹ J. Billowes,¹⁵ V. Boccone,⁷ D. Bosnar,¹⁶ M. Brugger,⁷ M. Calviani,⁷ F. Calviño,¹⁷ D. Cano-Ott,¹² C. Carrapiço,¹⁸ F. Cerutti,⁷ E. Chiaveri,⁷ M. Chin,⁷ G. Cortés,¹⁷ M. A. Cortés-Giraldo,¹⁹ I. Dillmann,²⁰ C. Domingo-Pardo,²¹ I. Duran,²² N. Dzysiuk,²³ C. Eleftheriadis,²⁴ M. Fernández-Ordóñez,¹² A. Ferrari,⁷ K. Fraval,⁴ S. Ganesan,²⁵ A. R. García,²⁶ G. Giubrone,²⁷ M. B. Gómez-Hornillos,¹⁷ I. F. Gonçalves,¹⁸ E. González-Romero,¹² F. Gramegna,²³ E. Griesmayer,²⁸ P. Gurusamy,²⁵ S. Harrisopulos,²⁹ M. Heil,²¹ K. Ioannides,³⁰ D. G. Jenkins,³¹ E. Jericha,²⁸ Y. Kadi,⁷ D. Karadimos,³⁰ G. Korschinek,³² M. Krčička,¹³ J. Kroll,¹³ C. Langer,² E. Lebbos,⁷ H. Leeb,²⁸ L. S. Leong,¹¹ R. Losito,⁷ M. Lozano,¹⁹ A. Manousos,²⁴ J. Marganec,¹⁰ S. Marrone,⁵ T. Martinez,¹² P. F. Mastinu,²³ M. Mastromarco,⁵ M. Meaze,⁵ E. Mendoza,¹² A. Mengoni,³³ P. M. Milazzo,¹⁴ F. Mingrone,³ M. Mirea,³⁴ W. Mondalaers,³⁵ C. Paradela,²² A. Pavlik,¹ J. Perkowski,¹⁰ R. Plag,²¹ A. Plompen,³⁵ J. Praena,¹⁹ J. M. Quesada,¹⁹ T. Rauscher,³⁶ A. Riego,¹⁷ F. Roman,^{7,34} C. Rubbia,^{7,37} R. Sarmiento,¹⁸ P. Schillebeeckx,³⁵ S. Schmidt,² G. Tagliente,⁵ J. L. Tain,²⁷ D. Tarrío,²² L. Tassan-Got,¹¹ A. Tsinganis,⁷ L. Tlustos,⁷ S. Valenta,¹³ G. Vannini,⁵ V. Variale,⁵ P. Vaz,¹⁸ A. Ventura,³³ M. J. Vermeulen,³¹ R. Versaci,⁷ V. Vlachoudis,⁷ R. Vlastou,³⁸ T. Ware,¹⁵ M. Weigand,² C. Weiß,²⁸ T. J. Wright,¹⁵ and P. Žugec¹⁶

(n_TOF Collaboration)

¹University of Vienna, Faculty of Physics, Austria²Johann-Wolfgang-Goethe Universität, Frankfurt, Germany³Dipartimento di Fisica, Università di Bologna, and Sezione INFN di Bologna, Italy⁴Commissariat à l'Énergie Atomique (CEA) Saclay - Irfu, Gif-sur-Yvette, France⁵Istituto Nazionale di Fisica Nucleare, Bari, Italy⁶Paul Scherrer Institut, Villigen PSI, Switzerland⁷European Organization for Nuclear Research (CERN), Geneva, Switzerland⁸Karlsruhe Institute of Technology, Campus Nord, Institut für Kernphysik, Karlsruhe, Germany⁹Department of Physics, University of Basel, Klingelbergstrasse 82, CH-4056 Basel, Switzerland¹⁰Uniwersytet Łódzki, Lodz, Poland¹¹Centre National de la Recherche Scientifique/IN2P3 - IPN, Orsay, France¹²Centro de Investigaciones Energeticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain¹³Charles University, Prague, Czech Republic¹⁴Istituto Nazionale di Fisica Nucleare, Trieste, Italy¹⁵University of Manchester, Oxford Road, Manchester, United Kingdom¹⁶Department of Physics, Faculty of Science, University of Zagreb, Croatia¹⁷Universitat Politècnica de Catalunya, Barcelona, Spain¹⁸Instituto Tecnológico e Nuclear (ITN), Lisbon, Portugal¹⁹Universidad de Sevilla, Spain²⁰Physik Department E12 and Excellence Cluster Universe, Technische Universität München, Munich, Germany²¹GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany²²Universidade de Santiago de Compostela, Spain²³Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Legnaro, Italy²⁴Aristotle University of Thessaloniki, Thessaloniki, Greece²⁵Bhabha Atomic Research Centre (BARC), Mumbai, India²⁶Centro de Investigaciones Energeticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain²⁷Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Spain²⁸Atominstytut, Technische Universität Wien, Austria²⁹National Centre of Scientific Research (NCSR), Demokritos, Greece³⁰University of Ioannina, Greece³¹University of York, Heslington, York, United Kingdom³²Technical University of Munich, Munich, Germany³³Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile (ENEA), Bologna, Italy³⁴Horia Hulubei National Institute of Physics and Nuclear Engineering - IFIN HH, Bucharest - Magurele, Romania³⁵European Commission JRC, Institute for Reference Materials and Measurements, Retieseweg 111, B-2440 Geel, Belgium³⁶Department of Physics and Astronomy, University of Basel, Basel, Switzerland³⁷Laboratori Nazionali del Gran Sasso dell'INFN, Assergi (AQ), Italy³⁸National Technical University of Athens (NTUA), Greece

(Received 6 January 2014; revised manuscript received 10 February 2014; published 28 February 2014)

The cross section of the $^{62}\text{Ni}(n,\gamma)$ reaction was measured with the time-of-flight technique at the neutron time-of-flight facility n_TOF at CERN. Capture kernels of 42 resonances were analyzed up to 200 keV neutron energy and Maxwellian averaged cross sections (MACS) from $kT = 5\text{--}100$ keV were calculated. With a total uncertainty of 4.5%, the stellar cross section is in excellent agreement with the the KADoNiS compilation at $kT = 30$ keV, while being systematically lower up to a factor of 1.6 at higher stellar temperatures. The cross section of the $^{63}\text{Ni}(n,\gamma)$ reaction was measured for the first time at n_TOF. We determined unresolved cross sections from 10 to 270 keV with a systematic uncertainty of 17%. These results provide fundamental constraints on s -process production of heavier species, especially the production of Cu in massive stars, which serve as the dominant source of Cu in the solar system.

DOI: [10.1103/PhysRevC.89.025810](https://doi.org/10.1103/PhysRevC.89.025810)

PACS number(s): 25.40.Lw, 25.40.Ny, 26.20.Kn, 27.50.+e

I. MOTIVATION

The astrophysical slow neutron capture process (s process) in stars produces about half of the elemental abundances between Fe and Bi. The s process is attributed to environments of neutron densities of typically $10^6\text{--}10^{12}$ cm^{-3} , resulting in neutron capture time scales of the order of years. When an unstable nucleus is produced by neutron capture, β decays are usually faster than subsequent neutron capture, so the reaction path follows the valley of stability. The s process takes place in different stellar sites. In particular, the s -process abundances in the solar system are made by contributions from different generations of stars, resulting in three major components, a *main*, a *weak*, and a *strong* component (see, e.g., [1]). The main component dominates in the s contributions between Zr and the Pb region and is mainly associated with thermally pulsing asymptotic giant branch (AGB) stars of $1\text{--}3 M_{\odot}$ with an initial metal content close to solar [2]. During the AGB phase, He burning takes place in a shell surrounding the inert C/O core of the star. Thermal pulses are caused by He shell flashes which occur because He burning cannot sustain hydrostatic equilibrium within a thin shell. As a consequence of the mixing processes and the temperature peaks induced by the thermonuclear flashes, neutrons are released in $^{13}\text{C}(\alpha,n)$ and the $^{22}\text{Ne}(\alpha,n)$ reactions, respectively [3]. The strong component also originates in AGB stars but with much lower metallicities than solar [4]. It is responsible for about half of the solar ^{208}Pb abundances and for the full s -process contribution to Bi. The weak s process takes place in massive stars ($>8M_{\odot}$) which later explode as supernova (e.g., [5]), and is producing most of the s abundances in the mass region between Fe and Zr [6–10]. In these stars, neutrons are mostly produced at the end of convective He Core burning and during the later convective carbon shell burning phase via $^{22}\text{Ne}(\alpha,n)$ reactions.

The resulting s -process abundances N_s depend strongly on cross sections averaged over the stellar neutron spectrum. These Maxwellian averaged cross sections (MACS) are defined as

$$\langle\sigma\rangle = \frac{2}{\sqrt{\pi}} \frac{1}{(k_B T)^2} \int_0^{\infty} \sigma(E) E \exp\left(-\frac{E}{k_B T}\right) dE, \quad (1)$$

where k_B is the Boltzmann constant, T the stellar temperature, and $\sigma(E)$ the cross section as a function of energy E . The temperatures in s -process environments range from 0.09 to 1 GK (GigaKelvin), corresponding to kT values between 5 and 90 keV. For an accurate determination of MACSs, $\sigma(E)$ should be known up to a few hundred keV. Accurate cross sections are particularly important between Fe and Zr and for the light neutron poisons. The uncertainty of a single cross section may be propagated to the abundances of the following isotopes on the s -process path, or over the complete s -process distribution in the case of neutron poisons (see, e.g., [11]). This propagation effect is a peculiar feature of the the weak s process [12,13].

To have accurate s -process abundances N_s derived from precise neutron capture measurements is also of great importance for r -process studies because solar r -process abundances N_r are computed as residuals of the total solar abundances N_{\odot} after subtracting N_s :

$$N_r = N_{\odot} - N_s. \quad (2)$$

Because current stellar cross sections in the Fe/Ni mass region exhibit fairly large uncertainties, a campaign was started at the neutron time of flight facility n_TOF at CERN to measure the neutron capture cross sections of all stable isotopes of Fe and Ni with improved accuracy. Additionally, the (n,γ) cross section of the long-lived radionuclide ^{63}Ni ($t_{1/2} = 101.2 \pm 1.5$ yr [14]) was studied at n_TOF [15]. This paper describes the measurement and data analysis of the (n,γ) experiments on ^{62}Ni and ^{63}Ni .

Current data on $^{62}\text{Ni}(n,\gamma)$ include time-of-flight measurements [16–19] as well as activation measurements to directly determine the MACS at $kT = 25$ keV [20–22]. Neutron capture resonances have been analyzed over a large energy range ($E_n < 200$ keV) by Beer and Spencer [17], while there are a few other measurements investigating only the first strong $\ell = 0$ resonance at 4.6 keV [23–25]. Different results for this first s -wave resonance ($\ell = 0$) lead to severe differences in the low neutron energy part of evaluated cross sections, listed in libraries such as ENDF/B-VII.1 [26], JENDL-4.0 [27], and JEFF-3.1.1 [28]. The n_TOF data allowed us to determine resonance capture kernels up to 200 keV neutron energy; Maxwellian averaged cross sections were determined from $kT = 5$ to 100 keV with uncertainties between 4.5 and 10.4%.

*Present address: School of Physics and Astronomy, University of Edinburgh, UK; claudia.lederer@ph.ed.ac.uk

We also measured the $^{63}\text{Ni}(n,\gamma)$ cross section above thermal neutron energies (25 meV). Results for the resonance capture kernels and Maxwellian averaged cross sections are already published in [15]. In this paper, we present results of the unresolved capture cross section between 10 and 270 keV.

II. MEASUREMENT

A. Facility

The measurements were performed at the neutron time-of-flight facility n_TOF (Phase2) at CERN. At n_TOF, a highly intense, pulsed neutron beam is produced by spallation reactions of a pulsed 20-GeV proton beam from the CERN Proton Synchrotron on a massive lead target. The initially very energetic neutrons are moderated by a water layer surrounding the spallation target. The resulting neutron flux approximates an energy dependence proportional to $1/E_n$ and ranges from thermal (25 meV) up to a few GeV. Because of the long flight path of 185 m and a pulse width of 7 ns, a high resolution in neutron energy of $\Delta E/E \approx 3 \times 10^{-4}$ and $\Delta E/E \approx 5 \times 10^{-3}$ can be achieved at 1 eV, and at 1 MeV, respectively [29]. For a detailed description of the n_TOF facility, see Ref. [29].

The (n,γ) reactions on ^{62}Ni and ^{63}Ni were studied in separated campaigns. During the ^{63}Ni campaign, additional data were taken again with the ^{62}Ni sample, because ^{62}Ni represented the most abundant impurity in the ^{63}Ni sample. For the final $^{62}\text{Ni}(n,\gamma)$ cross section, results from both campaigns were combined.

B. Detection setup

The prompt γ -ray cascade that is emitted after each neutron capture event was detected using a pair of C_6D_6 scintillation detectors where the housing was made of carbon fiber [30] to reduce their sensitivity to neutrons to the minimum possible value. This feature is important because neutrons scattered from the sample can be captured in the detector material and produce γ rays which are not distinguishable from neutron capture in the sample of interest. The C_6D_6 detector system is installed perpendicular to the neutron beam and about 9 cm upstream from the capture sample. In this configuration, background from in-beam photons, produced at the spallation target and scattered by the sample, is minimized. Additionally, angular distribution effects of γ rays from $\ell > 0$ resonances can be neglected in this position. The C_6D_6 detectors were calibrated at 0.662, 0.898, 1.836, and 4.44 MeV using standard ^{137}Cs , ^{88}Y , and AmBe γ sources. Calibration runs were repeated every week during the measurement to monitor the detector stability. The data acquisition system records the full

pulse shape using Flash ADCs at a sampling rate of 500 MHz, corresponding to a time resolution of 2 ns. A trigger signal from the Proton Synchrotron, shortly before the proton bunch hits the neutron target, starts the data acquisition. Data are recorded for 16 ms in the 8 MB on-board buffer memory of the digitizers, covering the neutron energy range down to 0.7 eV. In the second campaign, the data acquisition system was adjusted to a recording time of 80 ms, thus extending the minimum measurable neutron energy to 27 meV.

C. Samples

The ^{62}Ni sample consisted of 2 g metal powder, which was pressed into a stable pellet 20 mm in diameter and about 1 mm in thickness. The ^{63}Ni sample was produced about 30 years ago by breeding a highly enriched ^{62}Ni sample in the ILL high flux reactor at Grenoble [31–33]. A first analysis of this material confirmed that it was free of any detectable impurities apart from the ingrown Cu component. After a chemical separation of the Cu, the remaining Ni fraction was converted into NiO grains typically 1–2 mm in size and with a total mass of 1156 mg. Finally, the grains were sealed in a light cylindrical container made from polyether-ether-ketone ($[\text{C}_{20}\text{H}_{12}\text{O}_3]_n$, PEEK, wall thickness 0.15 mm), with a total weight of 180 mg. Mass spectroscopic analysis of the sample yielded a ^{63}Ni to ^{62}Ni ratio of 0.123 ± 0.001 . This sample was used for measuring the $^{63}\text{Ni}(n,\gamma)$ cross section [15] and for fitting the first large $^{62}\text{Ni}(n,\gamma)$ resonance at 4.6 keV because of its smaller thickness (see Sec. IV A 1 for details). Additionally to the Ni samples, a Au sample of the same geometry as the Ni samples was used to normalize the cross section. A summary of the samples is shown in Table I.

III. DATA ANALYSIS

A. Determination of the capture yield

All time-of-flight spectra were corrected for dead-time effects, which never exceeded 1%. The count rate C measured in a capture experiment is related to the capture yield Y_c via

$$C(E_n) = Y_c(E_n)\phi_n(E_n)\varepsilon_c + B(E_n), \quad (3)$$

with $\phi_n(E_n)$ being the number of neutrons hitting the sample, ε_c the detection efficiency for capture events, and $B(E_n)$ the background reactions. To obtain the detection efficiency which is independent of the de-excitation path of the compound nucleus, we applied the pulse height weighting technique [34]. Usually, the detection efficiency for a single γ ray depends strongly on its energy, but by subjecting a pulse height dependent weight to each recorded signal, one can achieve a

TABLE I. Sample characteristics. All samples were of cylindrical shape and 2 cm in diameter.

Sample	Mass (mg)	Enrichment (w%)		Thickness (10^{-3} atoms/b)	Chemical form
		^{62}Ni	^{63}Ni		
^{62}Ni	1989	98.0	–	6.20	Metal pellet
^{63}Ni	1156	69.4	8.4	5.68	Oxide grains
^{197}Au	596	–	–	0.584	Metal foil

detection efficiency,

$$\varepsilon_c = k \times E^*, \quad (4)$$

which is a linear function of the excitation energy E^* of the compound nucleus, regardless of the decay pattern of the capture cascade. The excitation energy E^* is the sum of the reaction Q value (6.84 and 9.66 MeV for ^{62}Ni and ^{63}Ni , respectively) and the neutron energy in the center-of-mass system. The weights can be parametrized with a polynomial function of the energy deposited in the detector. Weighting functions were determined by simulating the detector response to monoenergetic γ rays using GEANT-4 [35], implementing a detailed geometry of the experimental setup.

After weighting, the capture yield Y_c can be calculated as

$$Y_c(E_n) = N \frac{C_w(E_n) - B_w(E_n)}{E^* \phi_n(E_n)}, \quad (5)$$

where C_w is the weighted count rate, B_w the weighted background, N a normalization factor, and ϕ_n the neutron flux incident on the sample. We used a neutron flux evaluated using long-term measurements with several detectors and Monte Carlo simulations [36]. The uncertainty in the neutron flux is 2% below 10 keV and above 100 keV, and 4%–5% between 10 and 100 keV neutron energy. To obtain the absolute capture yield, the absolute detection efficiency, and the fraction of the neutron flux incident on the sample (beam interception factor) need to be known. After applying weighting functions, the efficiency to detect a capture event for each isotope only depends on the excitation energy of the compound nucleus. The systematic uncertainty in the capture yield ascribed to the pulse height weighting technique is 2% [37]. The normalization factor for obtaining the absolute detection efficiency N is then the same for all measured isotopes after scaling the weighted counts with the excitation energy E^* . The beam interception, together with the normalization factor N was determined with the saturated resonance technique at the $E_n = 4.9$ eV resonance in Au, using an Au sample of the same diameter as the Ni samples. If the Au sample is chosen sufficiently thick, no neutrons are transmitted through the sample at the resonance energy. Because the capture width Γ_γ is bigger than the neutron width Γ_n for this resonance, almost all neutrons interacting with the sample get captured. It was demonstrated in Ref. [38] that a normalization obtained from this saturated resonance in Au is nearly independent of even large changes in the resonance parameters.

Because the neutron beam profile changes with neutron energy, the beam interception factor depends slightly on neutron energy as well. This effect was determined by Monte Carlo simulations [29]. In the investigated neutron energy range the beam interception factor never changed by more than $\pm 1.5\%$ compared to the value at 4.9 eV. We estimated the systematic uncertainty of the final cross section from the normalization N and the beam interception, including a possible misalignment of the sample which would affect the energy dependence of the beam interception, as 1%. The resulting total systematic uncertainty for determining the absolute capture yield is consequently 3% up to 10 keV and from 100 to 200 keV, and 5.5% from 10 to 100 keV neutron energy.

The effective neutron flight path, and thus the neutron energy calibration, was determined relative to low energy resonances in Au, which have been recently measured at the time-of-flight facility GELINA with high precision [39].

B. Backgrounds

The background for capture measurements at n_TOF consists of a number of different components.

Ambient background is coming from cosmic rays, natural radioactivity, and a possible radioactivity of the sample itself. This background is determined by runs without neutron beam.

Sample-independent background, from reactions of the neutron beam with any structure material, is determined in runs with an empty sample holder.

Sample-dependent background consists of two components. Neutrons, scattered from the sample into the experimental area where they are captured, and photons, which are produced at the spallation target and are scattered from the sample into the detector. The latter background, called in-beam γ background, appears at neutron energies between 10 and 300 keV. It stems mainly from neutron capture on the hydrogen of the moderator and could be significantly improved in the second campaign by using borated water as moderator. This improvement is demonstrated in Fig. 1, which shows a comparison of the ^{62}Ni capture yields from both campaigns, using water in the first, and borated water in the second campaign.

Sample-dependent backgrounds can be investigated using black resonance filters installed about halfway between the spallation target and the sample. These filters are sufficiently thick that the neutron spectrum is left void of neutrons at the energies of certain strong resonances. Accordingly, events in these energy windows are from background reactions. We checked this background for neutron energies below 1 keV by comparing sample spectra with filters with the spectrum of the empty sample holder with filters and found no indication for such a sample related background. For higher neutron energies,




FIG. 1. (Color online) Capture yield of ^{62}Ni using water (black) and borated water (red) as moderator. The addition of boron yields a significant reduction of the photon-induced background in the keV region.

this comparison was not possible because of lack of statistics. Because this background, however, is varying smoothly with neutron energy, it can be assumed as being constant over the width of the resonance and therefore be fitted while fitting the resonance shape. This approach could be cross checked by analyzing the ^{62}Ni data from two different campaigns, each having different backgrounds (for the second campaign borated water was used as moderator, reducing the photon background). The capture kernels of ^{62}Ni resonances mostly agreed within statistical uncertainties for both campaigns. For the few exceptions, the standard deviation of the two fits was used as uncertainty of the capture kernel.

Multiple scattering (MS) is a background that arises when a neutron is captured in the sample after it had been scattered within the sample itself. This background can be large in resonances with high scattering-to-capture ratios and depends also strongly on the sample geometry. The MS corrections are considered by the SAMMY code [40], which was used for analyzing the neutron resonances in ^{62}Ni . For the unresolved




FIG. 2. (Color online) (Top) Capture yield of ^{62}Ni (black, solid line) compared with backgrounds from neutron reactions in surrounding materials (pink, solid line, measured with empty sample holder) and ambient background (blue, shaded line). While the ambient background is 2 orders of magnitude smaller than the signal over the whole energy range, the empty background plays a crucial role in the higher keV range. (Bottom) Zoom into the neutron energy region from 6 to 100 keV.




FIG. 3. (Color online) Capture yield of ^{63}Ni (green) compared with backgrounds from neutron reactions on ^{62}Ni in the sample (black) and with surrounding materials (pink, shaded line, measured with empty sample container), and ambient background (black, shaded line). The spectrum recorded with the ^{62}Ni sample was scaled to the areal density of ^{62}Ni in the ^{63}Ni sample.

cross section of ^{63}Ni no such corrections could be applied because of the unknown scattering cross section. However, the effect is small because the ^{63}Ni sample was relatively thin. A possible overestimation of the cross section from this effect is included in the systematic uncertainty of the cross section.

A further sample related background consists of γ rays originating from inelastic scattering of neutrons. This background can be neglected in this measurement because the first excited state in ^{62}Ni and the first excited state above the detector threshold of 250 keV in ^{63}Ni are above 0.5 MeV. In both cases population of those levels was not possible in the investigated neutron energy range [41].

The capture yields of ^{62}Ni and ^{63}Ni together with the ambient and sample-independent background components are shown in Figs. 2 and 3, respectively.

IV. RESULTS ON $^{62}\text{Ni}(n,\gamma)$

A. Resonance analysis

Neutron resonances up to about 200 keV neutron energy were identified and analyzed using the multilevel R -matrix code SAMMY [40]. The fitting procedure applied in SAMMY to find the “best fit” values of parameters and the associated parameter covariance matrix is based on the Bayes’ theorem. The resonance shapes were fitted using the Reich-Moore approximation, including corrections for self-shielding, multiple scattering, and impurities in the sample, which were mainly other Ni isotopes. Experimental effects, such as Doppler broadening and the resolution of the capture setup, were taken into account. Because the measured resonance widths were in most cases larger than the natural widths due to broadening, only the capture kernel could be determined. It is related to the resonance area via

$$k_\gamma = \frac{2}{\pi\lambda^2} \int_{-\infty}^{+\infty} \sigma(E)dE = g_s \frac{\Gamma_n\Gamma_\gamma}{\Gamma_n + \Gamma_\gamma}, \quad (6)$$

TABLE II. Resonance energies E_R and capture kernels k_γ of the $^{62}\text{Ni}(n,\gamma)$ reaction. When possible, Γ_γ values have been fitted using spin assignments and Γ_n values from Beer and Spencer [17]. Resonances, which were not seen in any previous measurement, are marked by an asterisk.

E_R (eV)	g_s	Γ_n (meV)	Γ_γ (meV)	k_γ (meV)
2128.6 ± 0.2				0.570 ± 0.043
4614.8 ± 6.8	1		2545 ± 143	
8438.4 ± 1.1				11.1 ± 0.5
9540.3 ± 0.7				146.4 ± 6.1
12 225.4 ± 1.7*				15.6 ± 2.5
17 791.5 ± 1.4				52.6 ± 2.2
20 602.3 ± 1.5*				37.6 ± 1.8
24 621.9 ± 0.5				76.9 ± 3.4
28 417.5 ± 3.0				124.4 ± 5.1
29 507.1 ± 3.2				211.2 ± 8.6
29 960.1 ± 2.4*				13.9 ± 2.0
34 473.5 ± 6.4				114 ± 12
38 279.5 ± 1.8				313 ± 17
40 547.8 ± 2.2				56.8 ± 6.6
41 241.6 ± 2.6				59 ± 12
43 023 ± 19	1	340 000	496 ± 45	
45 137.1 ± 2.1				160 ± 9
53 402.4 ± 6.0*				81 ± 14
57 024 ± 15				108 ± 15
57 634 ± 9*				71 ± 10
63 443.6 ± 2.9				90 ± 25
67 911.8 ± 2.6*				75 ± 28
70 892.9 ± 3.2*				61 ± 12
74 419.6 ± 2.6				186 ± 15
77 463 ± 25	1	700 000	265 ± 53	
78 519.3 ± 8.1				130 ± 14
81 469 ± 31*				79 ± 13
93 944 ± 46				114 ± 29
95 038 ± 1033	1	2 500 000	<1200	
104 168 ± 22				371 ± 73
106 550 ± 1460	1	4 600 000	<3300	
113 203.2 ± 6.7				208 ± 44
120 052 ± 47				323 ± 54
131 919 ± 15*				174 ± 36
139 011 ± 45				504 ± 84
144 191 ± 25				488 ± 135
147 713 ± 32*				568 ± 60
149 873 ± 66	1	140 000	584 ± 117	
161 745 ± 19				557 ± 94
170 593 ± 21*				403 ± 70
180 902 ± 21*				445 ± 91
187 175 ± 45	1	90 000	1610 ± 296	

where λ denotes the de Broglie wavelength at the resonance energy, and Γ_n , Γ_γ the neutron and capture widths of the resonance. The statistical spin factor $g_s = (2J + 1)/(2s + 1)(2I + 1)$ is determined by the resonance spin J , the neutron spin $s = 1/2$, and the spin I of the target nucleus. The results obtained from the SAMMY fits with their statistical uncertainties are listed in Table II for resonances up to 200 keV. We used the partial neutron widths Γ_n obtained by Beer and Spencer [17] for $\ell = 0$ resonances to fit the radiative width Γ_γ . For

resonances with $\ell > 0$, no experimental data for partial widths were available, so the capture kernel k_γ is given in the table. Examples for resonance fits are shown in Fig. 4. Table II lists the combined result and propagated statistical uncertainties of both measurement campaigns. The systematic uncertainties from the pulse height weighting (2%), the normalization (1%), and the neutron flux shape (2%–5%) are not included in Table II. This leads to a total systematic uncertainty in the capture kernel of 3% for resonances up to 10 keV and from 100–200 keV, and 5.5% for resonances from 10 to 100 keV.

1. Resonance at $E_R = 4.6$ keV

The shape of the neutron resonance at $E_R = 4.6$ keV is highly affected by background from multiple scattering, because of its very high scattering-to-capture ratio of ≈ 800 . It was found impossible to fit this resonance with the relatively thick ^{62}Ni sample, therefore data measured with the thinner ^{63}Ni sample, where multiple scattering is less important, were used to analyze this resonance. Because the estimated multiple scattering background varies with the neutron width Γ_n , the resonance was fitted while keeping Γ_n constant. The resonance was assigned as $\ell = 0$ because of its shape. Using two previously measured values for the neutron width, $\Gamma_n = 1.822$ keV [25] and $\Gamma_n = 2.075$ keV [24], Γ_γ values of 2.4 and 2.7 meV were obtained in the SAMMY fits, respectively. A resonance fit was not possible using a third experimental value for Γ_n of 1.3 keV [42]. Because this resonance is an s wave, the two possible options for the fit yield different contributions to the cross section at lower energies. In fact, the thermal cross section obtained with the two choices is 16.2 barn for $\Gamma_\gamma = 2.7$ meV, but only 12.8 barn for $\Gamma_\gamma = 2.4$ meV. Previous measurements of the thermal cross section result in reported values between 14.0 and 21 barn [43–50], with the majority of values grouped around 14.5 barn [43,45,46,48,49]. Because of this large spread, these previous measurements cannot give us a hint on the correct Γ_γ value. A new measurement of this resonance using a much thinner sample would be desirable in the future, especially because this resonance contributes about 50% to the Maxwellian averaged cross section (MACS) at $kT = 30$ keV.

2. Level spacing

It is expected that the average level density of the compound nucleus is constant over the investigated energy range. Figure 5 shows that the accumulated number of observed levels as a function of neutron energy follows the expected linear behavior up to about 80 keV. The increasing number of missing levels is due to the weakening signal-to-background ratio combined with the decreasing energy resolution of the n_TOF setup. We find an average level spacing of roughly 28 keV for s -wave resonances and 3.4 keV for $\ell > 0$ resonances. The consequences of missing resonances for the Maxwellian averaged cross sections are discussed in Sec. IV B.

B. Maxwellian averaged cross sections

We calculated Maxwellian averaged cross section from $kT = 5$ –100 keV using the resonance parameters obtained




FIG. 4. (Color online) (a)–(d) Examples for resonances fitted with the program SAMMY [40]. The dots are the measured data; the line represents the result of the resonance fit. Panel (a) shows the fit of the 4.6-keV resonance which was analyzed using the spectra obtained with the ^{63}Ni sample. The data in panels (b)–(d) are from the first ^{62}Ni campaign.




FIG. 5. (Color online) Accumulated number of levels as a function of neutron energy. The black dots represent the data; the red line is a linear fit from 0 to 80 keV.

from the SAMMY fits. Resonances parameters from 200 keV onwards were taken from the JENDL-4.0 library [27]. The MACS values from $kT = 5$ –100 keV together with their statistical and systematic uncertainties are listed in Table III, and Table IV details the individual uncertainties for three typical values of kT . Systematic uncertainties include the pulse height weighting technique, the normalization, and the neutron flux. The impact of the two different fits for the 4.6-keV resonance according to the different multiple scattering corrections was included as separate systematic uncertainty (called “MS at $E_R = 4.6$ keV” in Table IV).

To investigate the effect of missing levels on the MACS values an average cross section was calculated from our data in the energy range from 81 to 200 keV, using simulated self-shielding and multiple scattering corrections. These corrections were obtained by means of Monte Carlo simulations taking into account the sample geometry and neutron capture and scattering cross sections. The MACS values of this approach were between 3% and 7% higher in the range $kT = 40$ –100 keV than the results calculated from resonance

TABLE III. Maxwellian averaged cross sections of the $^{62}\text{Ni}(n,\gamma)$ reaction from 5 to 100 keV together with statistical and systematic uncertainties.

kT (keV)	MACS (mb)	Uncertainty (%)	
		Statistical	Systematic
5	181.2	0.6	5.2
10	83.2	0.6	4.9
15	50.8	0.6	4.8
20	35.8	0.7	4.4
25	27.4	1.0	4.3
30	22.2	1.5	4.2
40	16.0	2.7	−4.1/ + 5.1
50	12.5	3.8	−4.1/ + 6.7
60	10.2	4.7	−4.0/ + 7.2
80	7.44	6.0	−3.9/ + 8.0
100	5.75	6.7	−3.8/ + 8.0

TABLE IV. Contributions to the uncertainties (in %) for the stellar $^{62}\text{Ni}(n,\gamma)$ cross sections (see text for details).

kT (keV)	5	30	100
Weighting functions	2	2	2
Normalization	1	1	1
Neutron flux shape	2.0	2.7	2.9
MS at $E_R = 4.6$ keV	4.2	2.3	0.9
Missing levels	–	–	+7
Counting statistics	0.6	1.5	6.7
Total	5.2	4.5	–7.7/+10.4

data only. We included this difference as additional systematic uncertainty in Table IV (missing levels).

A comparison of our results to previous measurements and evaluations is shown in Fig. 6. For $kT < 30$ keV, the n_{TOF} data are in agreement with the results of Alpizar-Vicente *et al.* [19]. At 25 keV and 30 keV, our MACS is in excellent agreement with activation measurements of Nassar *et al.* [20] and Dillmann *et al.* [22], while being significantly lower than a previous time-of-flight measurement by Tomyo *et al.* [18]. Towards higher kT values, our data start to deviate from the results of Alpizar-Vicente *et al.* [19], being systematically lower up to a factor of 1.8. As investigated by Monte Carlo simulations, missing levels because of high background at high neutron energies cannot account for that difference. For $kT > 50$, our data are in fair agreement with MACS calculated using resonance parameters of the JENDL-4.0 evaluation [27], which is mainly based on a measurement by Beer and Spencer [17].

V. RESULTS FOR $^{63}\text{Ni}(n,\gamma)$

The resonance analysis for the $^{63}\text{Ni}(n,\gamma)$ reaction was already described and published in Ref. [15]. In this section,




FIG. 6. (Color online) Maxwellian averaged cross sections from 5 to 100 keV compared to previous measurements (Alpizar-Vicente *et al.* [19], Nassar *et al.* [20], Dillmann *et al.* [22], and Tomyo *et al.* [18]). The results obtained with data from the JENDL-4.0 evaluation (dashed line, [27]) and the recommended MACS values of the KADONIS compilation (solid line, [51]) are included as well.

TABLE V. Average $^{63}\text{Ni}(n,\gamma)$ cross sections between 10 and 270 keV laboratory neutron energy with statistical uncertainties. The total systematic uncertainty is 17%.

Neutron Energy (eV)		Cross section (mb)
E_{low}	E_{high}	
10 104	12 136	87 ± 3
12 136	14 577	142 ± 29
14 577	17 506	160 ± 26
17 506	21 023	111 ± 25
21 023	30 304	77 ± 15
30 304	43 664	57 ± 22
43 664	62 871	50 ± 12
62 871	90 456	37 ± 8
90 456	130 027	22 ± 7
130 027	186 705	18 ± 8
186 705	267 743	7.0 ± 5.8

the results for the unresolved cross section of $^{63}\text{Ni}(n,\gamma)$ are presented. From 10 keV onwards, we calculated an averaged cross section, because the high background, mainly from reactions of neutrons with ^{62}Ni and with the sample container prevented us from analyzing more resonances. The $^{63}\text{Ni}(n,\gamma)$ capture yield was calculated by subtracting the background from $^{62}\text{Ni}(n,\gamma)$ reactions using the spectra recorded with the ^{62}Ni sample and the known ^{62}Ni abundance in the ^{63}Ni sample. Background from reactions on oxygen is negligible, because of the low reaction cross section. The average cross section was calculated using the thin target approximation,

$$\sigma = \frac{Y_c}{n}, \quad (7)$$

where n is the areal density of the sample and Y_c the neutron capture yield. As for ^{62}Ni , systematic uncertainties are coming from the pulse height weighting technique (2%), the neutron flux (2%–5%), and the normalization (1%). Additionally, the $^{62}\text{Ni}:^{63}\text{Ni}$ ratio in the sample contributed an uncertainty of $\leq 2\%$. The background subtraction from reactions on ^{62}Ni in the sample introduces the largest systematic uncertainty, which we estimated as 15% based on different ways to treat the background at ^{62}Ni resonances. Assuming a high multiple scattering correction of 5%, the total systematic uncertainty of this measurement amounts to 17%. The cross sections from 10 to 270 keV, along with statistical uncertainties, are listed in Table V. MACS values and the implications of the ^{63}Ni cross section for stellar nucleosynthesis were published in [15].

VI. ASTROPHYSICAL IMPLICATIONS

In addition to the cross sections of the target nuclei in their ground states, as measured here, reactions on thermally excited states have to be considered in the determination of stellar reaction rates to be used in astrophysical models. For $^{62}\text{Ni}(n,\gamma)$, the population of excited states is negligible across the full energy range of s -process temperatures. Thus, the measured laboratory cross sections directly allow one to derive the stellar rates. Because of the higher nuclear level density




FIG. 7. (Color online) (Top) Final isotopic s -process abundances between Fe and Ga normalized to solar system abundances. The red circles represent the abundances using the $^{62}\text{Ni}(n,\gamma)$ MACS of this work and the $^{63}\text{Ni}(n,\gamma)$ MACS reported in Ref. [15]. This distribution is compared to the results using the recommended MACS of ^{62}Ni and ^{63}Ni of the KADONIS compilation [51]. As a reference, the overabundance of ^{16}O is shown as a black continuous line, divided and multiplied by 2 (black dashed lines). (Bottom) Ratio between the abundances using the new cross sections and the abundances using KADONIS cross sections.

of ^{63}Ni , only a fraction of the stellar rate can be constrained by a measurement of $^{63}\text{Ni}(n,\gamma)$ cross sections and theoretical corrections have to be applied as described in [15].

The impact of our new results on $^{62}\text{Ni}(n,\gamma)$ and $^{63}\text{Ni}(n,\gamma)$ on the weak s process in massive stars was investigated for a full stellar model for a $25 M_{\odot}$ star with an initial metal content of $Z = 0.02$ [52]. The complete nucleosynthesis was followed with the postprocessing NuGrid code MPPNP [53]. Figure 7 shows the s -process abundance distribution in the mass region from Fe to Ga after the convective core He and the convective C shell burning phase. Although the solar system s -process abundances in the Ni-Cu-Zn region may be partially affected by the following core-collapse supernova event (e.g., [13,54]), the pre-explosive s -process distribution is relevant as it serves as seed for the later explosive nucleosynthesis. The abundance distribution calculated with the MACSs of ^{62}Ni and ^{63}Ni from this work and Ref. [15] is compared to the abundances calculated with the recommended MACS data of the KADONIS compilation v0.3 [51]. Because the ^{62}Ni MACS of this work is smaller than the value in KADONIS for $kT > 50$ keV, neutron capture rates of ^{62}Ni in the C shell burning phase, where temperatures correspond to $kT \gtrsim 90$ keV, are

smaller and the final abundance of ^{62}Ni increases by 20%. The corresponding lower production of ^{63}Ni results in lower abundances of ^{63}Cu and ^{64}Zn . This decrease is compensated for ^{65}Cu and above ^{66}Zn by the fact that the ^{63}Ni MACS itself is a factor of 2 higher than the MACS value in KADONIS, resulting in a stellar rate that is about 40% larger than the KADONIS rate at typical shell C burning temperatures after considering the contribution from neutron capture on excited states in ^{63}Ni [55]. Accordingly, the s abundances in this region differ only by 1%–2%. For ^{63}Cu , which is mainly produced by the radiogenic decay of ^{63}Ni after C shell burning, the effect of a smaller ^{62}Ni MACS and a higher ^{63}Ni MACS causes a 40% lower abundance of ^{63}Cu . Because the ^{65}Cu abundance remains essentially unchanged, the isotopic ratio $^{63}\text{Cu} : ^{65}\text{Cu}$ is reduced after C shell burning. These results will allow one to better define the following explosive contribution to the copper inventory of the solar system.

VII. SUMMARY

We measured the cross sections of the $^{62}\text{Ni}(n,\gamma)$ and $^{63}\text{Ni}(n,\gamma)$ reactions at the neutron time-of-flight facility n_TOF at CERN.

For $^{62}\text{Ni}(n,\gamma)$, the resonance analysis was performed up to 200 keV neutron energy. In total, 42 levels could be identified, of which 12 were not seen in previous experiments. The Maxwellian averaged cross sections, calculated from resonance parameters is in good agreement with previous measurements for kT values up to 50 keV. At higher energies our results are systematically lower than the data by Alpizar-Vicente *et al.* [19], but in good agreement with the evaluations in the data libraries JENDL [27] and ENDF/B-VII [26], which are mainly based on a measurement by Beer and Spencer [17]. Our MACS at 100 keV is also a factor of 1.6 lower than the currently recommended value of the KADONIS compilation.

For the $^{63}\text{Ni}(n,\gamma)$ reaction, the neutron resonance analysis together with the stellar cross sections are published elsewhere [15]. We determined averaged cross sections between 10 and 270 keV with systematic uncertainties of 17%.

The impact of the new stellar (n,γ) cross sections of ^{62}Ni and ^{63}Ni was studied with a stellar model for a $25 M_{\odot}$ star with $Z = 0.02$. We find significant changes in the s abundances of ^{62}Ni (+20%) and ^{63}Cu (−40%), whereas the changes for heavier s -process isotopes are less than 2%. These results are particularly important to understand the solar system abundances of Cu, which is dominantly produced in massive stars.

ACKNOWLEDGMENTS

The authors would like to thank H. Danninger and C. Gierl of the Technical University of Vienna for their help preparing the ^{62}Ni sample. This work was partly supported by the Austrian Science Fund (FWF), Projects No. P20434 and No. I428 and by the Federal Ministry of Education and Research of Germany, Project No. 05P12RFFN6. M.P. acknowledges support from the Ambizione grant of the SNSF (Switzerland), from NuGrid thanks to the EU MIRG-CT-2006-046520,

from the NSF Grants No. PHY 02-16783 and No. PHY 09-22648 (Joint Institute for Nuclear Astrophysics, JINA), and from EuroGenesis (MASCHE). T.R. acknowledges the

Swiss NSF, the European Research Council, and the THEXO Collaboration within the 7th Framework Program ENSAR of the EU.

-
- [1] F. Käppeler, R. Gallino, S. Bisterzo, and W. Aoki, *Rev. Mod. Phys.* **83**, 157 (2011).
- [2] M. Busso, R. Gallino, and G. Wasserburg, *Ann. Rev. Astron. Astrophys.* **37**, 239 (1999).
- [3] R. Gallino, C. Arlandini, M. Busso, M. Lugaro, C. Travaglio, O. Straniero, A. Chieffi, and M. Limongi, *Astrophys. J.* **497**, 388 (1998).
- [4] C. Travaglio, R. Gallino, M. Busso, and R. Gratton, *Astrophys. J.* **549**, 346 (2001).
- [5] S. E. Woosley, A. Heger, and T. A. Weaver, *Rev. Mod. Phys.* **74**, 1015 (2002).
- [6] J. G. Peters, *Astrophys. J.* **154**, 225 (1968).
- [7] R. G. Couch, A. B. Schmiedekamp, and W. D. Arnett, *Astrophys. J.* **190**, 95 (1974).
- [8] S. A. Lamb, W. M. Howard, J. W. Truran, and I. Iben, *Astrophys. J.* **217**, 213 (1977).
- [9] C. M. Raiteri, M. Busso, G. Picchio, R. Gallino, and L. Pulone, *Astroph. J.* **367**, 228 (1991).
- [10] C. M. Raiteri, M. Busso, G. Picchio, and R. Gallino, *Astrophys. J.* **371**, 665 (1991).
- [11] C. Massimi *et al.* (n_TOF Collaboration), *Phys. Rev. C* **85**, 044615 (2012).
- [12] M. Busso and R. Gallino, *Astron. Astrophys.* **151**, 205 (1985).
- [13] T. Rauscher, A. Heger, R. Hoffman, and S. Woosley, *Astrophys. J.* **576**, 323 (2002).
- [14] R. Colle, B. Zimmermann, P. Cassette, and L. Laureano-Perez, *Applied Radiation and Isotopes* **66**, 60 (2008).
- [15] C. Lederer *et al.* (n_TOF Collaboration), *Phys. Rev. Lett.* **110**, 022501 (2013).
- [16] H. Beer, R. R. Spencer, and A. Ernst, *Astron. Astrophys.* **37**, 197 (1974).
- [17] H. Beer and R. R. Spencer, *Nucl. Phys. A* **240**, 29 (1975).
- [18] A. Tomyo, Y. Temma, M. Segawa, Y. Nagai, H. Makii, T. Ohsaki, and M. Igashira, *Astrophys. J.* **623**, L153 (2005).
- [19] A. M. Alpizar-Vicente, T. A. Bredeweg, E. I. Esch, U. Greife, R. C. Haight, R. Hatarik, J. M. O'Donnell, R. Reifarth, R. S. Rundberg, J. L. Ullmann, D. J. Vieira, and J. M. Wouters, *Phys. Rev. C* **77**, 015806 (2008).
- [20] H. Nassar, M. Paul, I. Ahmad, D. Berkovits, M. Bettan, P. Collon, S. Dababneh, S. Ghelberg, J. P. Greene, A. Heger, M. Heil, D. J. Henderson, C. L. Jiang, F. Kappeler, H. Koivisto, S. O'Brien, R. C. Pardo, N. Patronis, T. Pennington, R. Plag, K. E. Rehm, R. Reifarth, R. Scott, S. Sinha, X. Tang, and R. Vondrasek, *Phys. Rev. Lett.* **94**, 092504 (2005).
- [21] S. Walter, Ph.D thesis, University of Karlsruhe, 2008.
- [22] I. Dillmann, T. Faestermann, G. Korschinek, J. Lachner, M. Maiti, M. Poutivtsev, G. Rugel, S. Walter, F. Kaeppler, M. Erhard, A. R. Junghans, C. Nair, R. Schwengner, and A. Wagner, *Nucl. Instrum. Methods Phys. Res., Sect. B* **268**, 1283 (2010); *11th International Conference on Accelerator Mass Spectrometry*, Rome, Italy, Sept. 14–19, 2008.
- [23] R. W. Hockenbury, Z. M. Bartolome, J. R. Tatarczuk, W. R. Moyer, and R. C. Block, *Phys. Rev.* **178**, 1746 (1969).
- [24] H. P. Axmann, D. A. J. Endacott, J. E. Jolly, and M. C. Moxon, *Nickel Isotope Resonance Parameters*, Progress Report 18 (Atomic Energy Research Establishment, Harwell Reports, Harwell, 1972).
- [25] L. Litvinskiy, P. Vorona, V. Krivenko, V. Libman, A. Murzin, G. Novoselov, N. Trofimova, and O. Chervonnaya, *Vop. At. Nauki i Tekhn., Ser. Yadernye Konstanty* **1990**, 27 (1990).
- [26] M. Chadwick, M. Herman, P. Obložinský, M. Dunn, Y. Danon, A. Kahler *et al.*, *Nucl. Data Sheets* **112**, 2887 (2011).
- [27] K. Shibata, O. Iwamoto, T. Nakagawa, N. Iwamoto, A. Ichihara, S. Kunieda, S. Chiba, K. Furutaka, N. Otuka, T. Ohsawa, T. Murata, H. Matsunobu, A. Zukeran, S. Kamada, and J. Katakura, *J. Nucl. Sci. Technol.* **48**, 1 (2011).
- [28] A. Koning, R. Forrest, M. Kellett, R. Mills, H. Henriksson, and Y. Rugama, JEFF Report 21 (OECD/NEA, Paris, 2006).
- [29] C. Guerrero *et al.* (n_TOF Collaboration), *Eur. Phys. J. A* **49**, 27 (2013).
- [30] R. Plag, M. Heil, F. Käppeler, P. Pavlopoulos, R. Reifarth, and K. Wisshak, *Nucl. Instrum. Methods Phys. Res., Sect. A* **496**, 425 (2003).
- [31] A. Harder, S. Michaelsen, A. Jungclaus, A. Williams, H. Börner, and M. Trautmannsheimer, *Z. Phys. A* **343**, 7 (1992).
- [32] H. Muthig, Ph.D thesis, Technical University of Munich, 1984.
- [33] M. Trautmannsheimer, Ph.D thesis, Technical University of Munich, 1992.
- [34] R. Macklin and R. Gibbons, *Phys. Rev.* **159**, 1007 (1967).
- [35] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [36] M. Barbagallo *et al.* (n_TOF Collaboration), *Eur. Phys. J. A* **49**, 156 (2013).
- [37] U. Abbondanno *et al.* (n_TOF Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **521**, 454 (2004).
- [38] A. Borella, G. Aerts, F. Gunsing, M. Moxon, P. Schillebeeckx, and R. Wynants, *Nucl. Instrum. Methods Phys. Res., Sect. A* **577**, 626 (2007).
- [39] C. Massimi, A. Borella, S. Kopecky, C. Lampoudis, P. Schillebeeckx, M. Moxon, and G. Vannini, *J. Korean Phys. Soc.* **59**, 1689 (2011).
- [40] N. Larson, *Updated Users Guide for SAMMY: Multi-level R-matrix Fits to Neutron Data Using Bayes Equations*, Report No. ORNL/TM-9179/R7 (Oak Ridge National Laboratory, Oak Ridge, 2003).
- [41] B. Singh, *Nucl. Data Sheets* **108**, 197 (2007).
- [42] J. Garg, J. Rainwater, and W. Havens, *Phys. Rev. C* **3**, 2447 (1971).
- [43] H. Pomerance, *Phys. Rev.* **88**, 412 (1952).
- [44] C. McMullen, B. Pate, R. Tomlinson, and L. Yaffe, *Can. J. Chem.* **34**, 1742 (1956).
- [45] D. Horrocks and A. Harkness, *Phys. Rev.* **125**, 1619 (1962).
- [46] G. Sims and D. Juhnke, *J. Inorg. Nucl. Chem.* **32**, 2839 (1970).
- [47] H. Ishikawa, *Nucl. Instrum. Methods Phys. Res.* **109**, 493 (1973).

- [48] H. Michael, A. Neubert, and H. Nickel, *Applied Radiation and Isotopes* **25**, 183 (1974).
- [49] A. Ishaq, A. Robertson, W. Prestwich, and T. Kennett, *Z. Phys. A* **281**, 365 (1977).
- [50] L. Venturini and B. Pecequilo, *Applied Radiation and Isotopes* **48**, 493 (1997).
- [51] I. Dillmann, R. Plag, F. Käppeler, and T. Rauscher, in *Proceedings of the Workshop “EFNUDAT Fast Neutrons–Scientific Workshop on Neutron Measurements, Theory & Applications,”* April 28–30, 2009, Geel, Belgium, p. 55.
- [52] M. Pignatari, F. Herwig, R. Hirschi, M. Bennett, G. Rockefeller, C. Fryer, F. X. Timmes, A. Heger, S. Jones, U. Battino, C. Ritter, A. Dotter, R. Trappitsch, S. Diehl, U. Frischknecht, A. Hungerford, G. Magkotsios, C. Travaglio, and P. Young, [arXiv:1307.6961](https://arxiv.org/abs/1307.6961).
- [53] M. Pignatari and F. Herwig, *Nucl. Phys. News* **22**, 18 (2012).
- [54] M. Pignatari, R. Gallino, M. Heil, M. Wiescher, F. Käppeler, F. Herwig, and S. Bisterzo, *Astrophys. J.* **710**, 1557 (2010).
- [55] T. Rauscher, *Astrophys. J.* **755**, L10 (2012).