

Manifestations of the electron-phonon interaction range in angle-resolved photoemission spectra

Krsnik, J.; Strocov, V. N.; Nagaosa, N.; Barišić, Osor Slaven; Rukelj, Zoran; Yakubenya, S. M.; Mishchenko, A. S.

Source / Izvornik: **Physical Review B**, 2020, 102

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1103/PhysRevB.102.121108>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:217:876445>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-01-31**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



Manifestations of the electron-phonon interaction range in angle-resolved photoemission spectraJ. Krsnik^{1,*}, V. N. Strocov,² N. Nagaosa,^{3,4} O. S. Barišić,¹ Z. Rukelj,⁵ S. M. Yakubenko,⁶ and A. S. Mishchenko^{3,6}¹*Institute of Physics, HR-10000 Zagreb, Croatia*²*Swiss Light Source, Paul Scherrer Institute, Villigen CH-5232, Switzerland*³*RIKEN Center for Emergent Matter Science (CEMS), Wako, Saitama 351-0198, Japan*⁴*Department of Applied Physics, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan*⁵*Department of Physics, Faculty of Science, University of Zagreb, HR-10000 Zagreb, Croatia*⁶*NRC “Kurchatov Institute,” 123182 Moscow, Russia*

(Received 4 December 2019; revised 2 September 2020; accepted 3 September 2020; published 22 September 2020)

Numerous angle-resolved photoemission spectroscopy (ARPES) studies of a wide class of low-density metallic systems, ranging from doped transition metal oxides to quasi-two-dimensional interfaces between insulators, exhibit phonon sidebands below the quasiparticle peak as a unique hallmark of polaronic correlations. Here, we single out properties of ARPES spectra that can provide a robust estimate of the effective range (screening length) of the electron-phonon interaction, regardless of the limited experimental resolution, dimensionality, and particular features of the electronic structure, facilitating a general methodology for an analysis of a whole class of materials.

DOI: [10.1103/PhysRevB.102.121108](https://doi.org/10.1103/PhysRevB.102.121108)

Introduction. It has been well established, both experimentally and theoretically, that for low concentrations of itinerant charge carriers the electron-phonon interaction (EPI) produces phonon sidebands, appearing as satellites in angle-resolved photoemission spectroscopy (ARPES) spectra below the quasiparticle peak (QP) [1]. It has been argued [2,3] that energy and momentum intensity distributions of these satellites depend on the spatial range of the EPI, the unscreened polar Fröhlich interaction having the longest and the Holstein on-site interaction the shortest range. However, despite extensive ARPES data available [4–13], we are not aware of any systematic theoretical study which would consider the dependence of these very unique manifestations of the EPI on the screening length. We can mention only a study [2] of the screened Fröhlich interaction for a few particular cases in specific materials and the theoretical study [3] of the EPI with a hypothetical forward scattering. Hence, an accurate theoretical description of ARPES spectra in dependence on the screening length is of primary importance for a knowledge of the effective interaction that governs polaronic correlations in a plethora of materials [4–13], exhibiting metal-insulator transitions, colossal magnetoresistance, commensurate-incommensurate transition, quantum Hall effect, etc.

Recently, first-principle calculations [14] have been used, considering the leading-order expansion of the electron self-energy [15,16] and the cumulant expansion [10,12,17], to simulate measured ARPES spectra of some specific polaronic materials. In their implementation of the cumulant expansion, the latter works are restricted to self-energy diagrams with electron Green's functions propagating in one direction of time only [18], giving approximate results for finite electron

densities, including the lowest-order case. For the higher-order corrections, additional approximations are introduced by treating higher phonon processes as uncorrelated [19]. On the other hand, our calculations are based on a direct evaluation of self-energy diagrams, considering the exact momentum and frequency dependence of the leading corrections to the phonon sidebands, including the exact leading (second-order) vertex correction to the second sideband. Instead of focusing on some particular system, our analysis identifies different behaviors across all ranges of EPI and provides a general procedure for the estimation of the EPI range from experimental data, prior to material-specific calculations.

Modeling. We consider the standard EPI model for a D -dimensional lattice, describing the interaction between bare electrons in a band, $\hat{H}_{el} = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}$, and dispersionless optical phonons with frequency ω_0 , $\hat{H}_{ph} = \omega_0 \sum_{\mathbf{q}} b_{\mathbf{q}}^{\dagger} b_{\mathbf{q}}$. The screened Fröhlich interaction is given by

$$\hat{H}_{el-ph} = \sum_{\mathbf{k}, \mathbf{q}} V_{\mathbf{q}}^d a_{\mathbf{k}+\mathbf{q}}^{\dagger} a_{\mathbf{k}} (b_{-\mathbf{q}}^{\dagger} + b_{\mathbf{q}}), \quad (1)$$

where $|V_{\mathbf{q}}^d|^2 = a_d / (|\mathbf{q}|^{d-1} + q_{TF}^{d-1})$, with $a_{d=3} = 2\sqrt{2}\pi\alpha$ for three-dimensional (3D) and $a_{d=2} = \sqrt{2}\pi\alpha$ for two-dimensional (2D) systems [20–24], and α characterizing the strength of the EPI [23]. We assume a static screening characterized by the Thomas-Fermi wave number q_{TF} . $r_{TF} = \pi/q_{TF}$ roughly gives the screening range in the lattice constant units, with $q_{TF} \rightarrow 0$ ($q_{TF} \rightarrow \infty$) for the unscreened Fröhlich (Holstein) EPI.

The spectral function $A(\mathbf{k}, \omega)$, measured by ARPES for $\omega < 0$ (electron removal processes), is determined by the electron Green's function $G(\mathbf{k}, \omega)$, $A(\mathbf{k}, \omega) = |\text{Im}G(\mathbf{k}, \omega)|/\pi$. In terms of the self-energy $\Sigma(\mathbf{k}, \omega)$, appearing due to the EPI in

*Corresponding author: jkrsnik@ifs.hr

Eq. (1), it is given by

$$A(\mathbf{k}, \omega) = \frac{|\text{Im}\Sigma(\mathbf{k}, \omega)|}{[\omega - \xi_{\mathbf{k}} - \text{Re}\Sigma(\mathbf{k}, \omega)]^2 + [\text{Im}\Sigma(\mathbf{k}, \omega)]^2}, \quad (2)$$

with $\xi_{\mathbf{k}} = \epsilon_{\mathbf{k}} - \mu < 0$, the hole energy measured from the Fermi level μ . For a weak EPI, the self-energy contributions in the denominator of Eq. (2) may be neglected. For a stronger EPI, the real part of the self-energy may be absorbed [25] into a renormalized dispersion $\tilde{\epsilon}_{\mathbf{k}}$, $\tilde{\epsilon}_{\mathbf{k}} = \epsilon_{\mathbf{k}} + \text{Re}\Sigma(\mathbf{k}, \omega)$. This facilitates experimental analysis, because $\tilde{\epsilon}_{\mathbf{k}}$ is a measured quantity. We neglect the imaginary part $\text{Im}\Sigma(\mathbf{k}, \omega)$ in the denominator of Eq. (2) since it barely affects the \mathbf{k} and ω dependence of the spectral density $A(\mathbf{k}, \omega)$.

Our attention is on recent ARPES studies of polaronic materials, when only a small part of the conduction band $\epsilon_{\mathbf{k}}$ is filled. The EPI leads to a complex structure of $A(\mathbf{k}, \omega)$, where in addition to the QP peak one observes phonon sidebands, with the n th sideband being shifted downward from the Fermi level by $n\omega_0$ [2,3]. Since the ARPES seldom show more than two sidebands [6,8,10,12], we concentrate our analysis on these. In particular, we consider the $\mu < \omega_0$ case that ensures a pattern of spectrally separated phonon sidebands, experimentally reported for many different systems [4–8,10–12]. Within the zero-temperature diagrammatic expansion, the first-order contribution in α to $\text{Im}\Sigma$ [26],

$$\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega) = \pi \sum_{\mathbf{q}}^{\text{BZ}} |V_{\mathbf{q}}^d|^2 \times \Theta(-\xi_{\mathbf{k}+\mathbf{q}}) \delta(\omega - \xi_{\mathbf{k}+\mathbf{q}} + \omega_0), \quad (3)$$

corresponds to the leading contribution to the first phonon sideband, restricted to the frequency window $[-\omega_0 - \mu, -\omega_0]$. The second sideband is found in the window $[-2\omega_0 - \mu, -2\omega_0]$, with the leading contribution given by second-order α^2 terms derived here (for details, see the Supplemental Material [27]),

$$\begin{aligned} \text{Im}\Sigma^{(2)}(\mathbf{k}, \omega) &= \pi \sum_{\mathbf{q}}^{\text{BZ}} \Theta(-\xi_{\mathbf{q}}) \delta(\omega - \xi_{\mathbf{q}} + 2\omega_0) \sum_{\mathbf{q}'}^{\text{BZ}} \frac{|V_{\mathbf{q}'}^d|^2 |V_{\mathbf{q}-\mathbf{q}'}^d|^2}{(\xi_{\mathbf{q}} - \xi_{\mathbf{k}+\mathbf{q}'} - \omega_0)} \\ &\times [(\xi_{\mathbf{q}} - \xi_{\mathbf{k}+\mathbf{q}'} - \omega_0)^{-1} + (\xi_{\mathbf{q}} - \xi_{\mathbf{q}-\mathbf{q}'} - \omega_0)^{-1}]. \end{aligned} \quad (4)$$

The first term in the last line of Eq. (4) is given by the non-crossing two-phonon diagram, whereas the crossing diagram with the leading vertex correction gives the second term.

We set $\omega_0 = 1$ and consider the dispersion for a cubic lattice in 3D/2D, $\epsilon_{\mathbf{k}} = 2t \sum_{i=1}^D [1 - \cos(k_i)]$, where t is the nearest-neighbor hopping. With the lattice constant $a = 1$, the effective mass of the noninteracting electron at the bottom of the band is $m_0 = 1/2t$. To search for general properties of ARPES sideband components in various systems, we use two very different sets of parameters for the broad and the narrow electron bands: S1 (S2) denotes $\mu = 0.5$ ($\mu = 0.1$) and $t = 1$ ($t = 1/24$), with mass $m_0 = 0.5$ ($m_0 = 12$). In both these cases, only a small fraction of the lowest band states is occupied.

Phonon sidebands. A typical example of the ARPES intensity and of the corresponding $\text{Im}\Sigma$ intensity for the first




FIG. 1. Contour plot of the (a) ARPES component $A^{(1)}(\mathbf{k}, \omega)$ and (b) $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ for the first sideband for the 3D parameter set S1 and strong screening $r_{TF} = 0.1$. Right panels show EDCs for the corresponding cross sections in the left panels. Note, two EDCs for $\text{Im}\Sigma$ coincide.

sideband and strong screening is shown in Fig. 1. In experimental works, it is often concluded that the higher intensity accumulated within the Fermi surface (FS), $k \leq k_F \ll \pi$, identifies the weakly screened Fröhlich interaction. However, a more careful inspection of Eq. (2) leads to a different interpretation. $A^{(1)}(\mathbf{k}, \omega)$ in Fig. 1(a) exhibits a strong momentum dependence, due to $(\omega - \xi_{\mathbf{k}})^{-2}$ in the denominator in Eq. (2), which highlights the area within the FS. In fact, the latter easily camouflages the fact that the EPI might be over-screened. Indeed, as shown in Fig. 1(b), for strong screening $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ is momentum independent and uniformly spread over the whole Brillouin zone (BZ): in the right panel of Fig. 1(b), the two energy distribution curves (EDCs) for $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ are almost indistinguishable. The ω dependence is determined solely by the density of occupied electron states at the bottom of the band. This may easily be derived from Eq. (3), assuming a momentum-independent interaction $V_{\mathbf{q}}^d$. Different behaviors of $A^{(1)}(\mathbf{k}, \omega)$ and $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ in Fig. 1 explain the universality of the experimentally found confinement of the intensity of the ARPES sidebands in the momentum-space region near the band minimum [4–8,10,12]. However, it emphasizes as well that very different confinements may characterize the ARPES spectrum and the corresponding $\text{Im}\Sigma$ intensity, rendering $\text{Im}\Sigma$ as a quantity that is far more suitable for estimation of the range of EPI.

In contrast to strong screenings, for weak screenings ($r_{TF} \gtrsim 20$) the map of $\text{Im}\Sigma$ in Fig. 2 exhibits maxima that almost exactly follow the QP dispersion. In fact, using this property the weak-screening regime can be unambiguously identified from the behavior of the first phonon sideband. On the other hand, in our results the EDC and momentum distribution curves (MDCs) maxima of $\text{Im}\Sigma$ for the second sideband are absent, with its typical intensity shown in Fig. 3(a). By taking the second derivative $d^2\text{Im}\Sigma^{(2)}(\mathbf{k}, \omega)/dk^2$, which is fully in line with the conventional experimental ARPES-data processing that uses the curvature analysis, one recovers a dispersion resembling that of the QP peak. This is a general property of the second sideband, well illustrated by Fig. 3, in which the intensities of $\text{Im}\Sigma^{(2)}(\mathbf{k}, \omega)$ and $d^2\text{Im}\Sigma^{(2)}(\mathbf{k}, \omega)/dk^2$ are compared, together with the intensity dependence along the EDCs and the MDCs. We note that the necessity for the differential analysis of the second sideband follows purely from theoretical results,




FIG. 2. Contour plots of $\text{Im}\Sigma$ for the QP band, the first and the second phonon sideband for the 3D parameter set S1, and for weak screening $r_{TF} = 100$.

rather than being a consequence of a limited quality of particular experimental data.

Anomalous phonon sideband dispersions. Between the strong and the weak screening regimes, exhibiting clearly different and characteristic behaviors of $\text{Im}\Sigma$, another regime sets in, which can be recognized from anomalous phonon sideband dispersions. For intermediate EPI ranges, the structure of $\text{Im}\Sigma$ is very sensitive on parameters, which is illustrated by Fig. 4. The curve that follows the EDC maxima




FIG. 3. Contour plots of (a) $\text{Im}\Sigma^{(2)}(\mathbf{k}, \omega)$ and (b) its second derivative $d^2\text{Im}\Sigma^{(2)}(\mathbf{k}, \omega)/dk^2$ for the 3D parameter set S1 and $r_{TF} = 100$. Right panels in (a) and (b) are the EDCs and the upper panels are the MDCs along the cuts shown in the left-bottom panels of (a) and (b). The dashed curves represent the QP dispersion, shifted by $2\omega_0$.




FIG. 4. (a) Contour plot of $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ for the 3D parameter set S1 and $r_{TF} = 7$. The dashed curve represents the QP dispersion $\epsilon_{\mathbf{k}}$, shifted downward by ω_0 . The circles follow the maxima of EDCs, fitted by a parabola (solid line) that corresponds to the mass $m^* = 0.76m_0$. (b) EDCs for cross sections shown in panel (a). (c) m^* derived from the first sideband for parameters when EDCs maxima do not follow the QP dispersion. (d) Upward shift ϵ_0 characterizing the EDC maxima for the first phonon sideband.

appears shifted upward by ϵ_0 from the lower edge of the first sideband. In comparison to the QP dispersion, parabolic fits of the EDCs maxima give different effective masses, e.g., for the $\text{Im}\Sigma$ intensity shown in Fig. 4 one obtains $m^* = 0.76m_0$. Such anomalous phonon sideband dispersions are found for all eight studied cases. Furthermore, anomalies in 3D and 2D are different: Figs. 4(c) and 4(d) show that m^* is smaller (larger) in 3D (2D) and ϵ_0 varies (being zero) in 3D (2D). While in the 3D cases the EDC maxima follow the parabolic dispersion over the whole first sideband, in the 2D cases the parabolic dispersion at small momenta transforms, exhibiting a jump toward the large momenta.

The anomalous sideband dispersion, which can be obtained either from the EDCs or from the standard curvature analysis, characterizes the ARPES spectra as well. Importantly, as seen from Fig. 5(a), the values of ϵ_0 and m^* obtained from these spectra and from $\text{Im}\Sigma$ are different. For the intermediate




FIG. 5. Contour plot of the (a) ARPES component $A^{(1)}(\mathbf{k}, \omega)$ and (b) $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ for the 3D parameter set S2 and $r_{TF} = 5$. Circles follow maxima of EDCs. These are fitted by parabolas, giving corresponding effective masses m^* and shifts ϵ_0 of the phonon sideband dispersion.

screening length in Fig. 5, $r_{TF} = 5$, the confinement of the ARPES intensity is enhanced in comparison to that obtained from $\text{Im}\Sigma$, being reduced to momenta, $k < k_r$, where k_r is considerably smaller than k_F . This effect is experimentally observed as well [4,8,12]. In general, for all the considered models and dimensionalities, we find that all anomalous behaviors are restricted to the specific screening range, $3 \leq r_{TF} \leq 20$.

Confinement estimators. Depending on the behavior of the first phonon sideband, our analysis identified three different regimes as a function of the range of the EPI. The calculations as well elucidated few features of ARPES sidebands that are independent of specific details of a measurement or material parameters, like the direction $[klm]/[kl]$, the dimensionality 3D/2D or the parameter set S1/S2. In particular, the confinement of the ARPES sidebands in the Brillouin zone is a common property for all three screening regimes. On the other hand, the confinement of the $\text{Im}\Sigma$ intensity depends strongly on r_{TF} . In fact, a procedure based on this confinement may be developed to determine the range of the EPI from experimental ARPES spectra.

Let us start by considering the $\text{Im}\Sigma$ intensity integrated over the energies within the n th phonon sideband for a given \mathbf{k} , $\mathcal{R}_n(\mathbf{k}) = \int_{-n\omega_0 - \mu}^{-n\omega_0} d\omega \text{Im}\Sigma^{(n)}(\mathbf{k}, \omega)$. Then, using $\mathcal{R}_n(\mathbf{k})$, the confinement within the FS may be expressed by an estimator corresponding to the ratio of the intensity within and outside the FS,

$$R_n = \left(\int_0^{k_F} dk \frac{\mathcal{R}_n(\mathbf{k})}{k_F} \right) / \left(\int_{k_F}^{\pi} dk \frac{\mathcal{R}_n(\mathbf{k})}{\pi - k_F} \right). \quad (5)$$

Such estimator of the confinement is particularly suitable for an analysis of the experimental data since it involves averaging over frequency and momenta intervals, reducing the effects of noise in experimental ARPES spectra. Moreover, it is unaffected by the arbitrariness in the normalization of the experimental ARPES intensity, as well as it is independent of the coupling strength α . Furthermore, the regime of strong screening, when $\text{Im}\Sigma$ is \mathbf{k} independent, may be identified directly from R_n , $R_n \rightarrow 1$ when $r_{TF} \rightarrow 0$.

With the exception of the $r_{TF} \rightarrow 0$ limit, R_n is insufficient to estimate r_{TF} . Namely, for different systems the maximal value of R_n may change by an order of magnitude. Instead of R_n , thus, in such situations we consider another estimator of the confinement,

$$C_n(r_{TF}) = [R_n(r_{TF}) - 1][R_n(\infty) - 1]^{-1}. \quad (6)$$

For all eight cases (parameter sets S1/S2, 3D/2D, diagonal $[k'k'k']/[k'k']$, and nondiagonal $[k'00]/[k', 0]$ directions) and for both sidebands, the estimator in Eq. (6) exhibits a fairly universal behavior, which is well illustrated by Fig. 6. $C_n(r_{TF})$ is scaled in Eq. (6) by the unscreened $r_{TF} \rightarrow \infty$ value of R_n . This value may be obtained from measurements of the reference material with unscreened EPI (small dopings), particularly for systems and experimental setups for which the charge density may easily be controlled. Alternatively, for the first sideband, $R_1(\infty)$ may easily be obtained numerically from closed expressions for $\text{Im}\Sigma^{(1)}(\mathbf{k}, \omega)$ for the 2D [27] and 3D [26] systems. Namely, all parameters that define $R_1(\infty)$




FIG. 6. Confinement parameter $C_n(r_{TF})$ (6) for the parameter sets S1 and S2, the first O1 and the second O2 phonon sidebands in (a) 3D and (b) 2D.

(effective mass m^* , Fermi level μ and k_F) may be extracted directly from the experimental QP dispersion.

Experimental data. We end our analysis with an illustrative example, dealing with a particularly complex experimental realization of EPI effects for low electron densities. We consider the ARPES spectrum of oxygen-deficient $\text{LaAlO}_3/\text{SrTiO}_3$ interface [28], involving different phonons and limited experimental resolution (0.04 eV), making a theoretical modeling difficult. Yet, even in such circumstances, our approach may provide valuable insights on the range of the EPI. Figure 7(a) shows the details of the APRES spectrum, after the subtraction of the structureless background. $k_F \approx 0.37 \text{ \AA}^{-1}$ and $\mu \approx 0.07 \text{ eV}$ may easily be estimated from the QP dispersion.

As explained in the discussion that follows Eq. (2), with QP properties known, one may obtain $\text{Im}\Sigma$ in Fig. 7(b) from the ARPES data in Fig. 7(a) (for details, see [27]). The frequency window in Fig. 7(b) dominated by the coupling to the LO3 phonon is highlighted separately, corresponding to the first LO3 sideband ($\omega_{LO3} \approx 0.12 \text{ eV}$ [8]). By averaging $\mathcal{R}_1(\mathbf{k})$ over these frequencies, $-\mu - \omega_{LO3} \leq \omega \leq -\omega_{LO3}$, one obtains $R_1 = 1.03$. This is almost the minimal value that the estimator R_1 can take, which alone clearly indicates the short-range EPI and the strong screening limit. The latter, as analyzed theoretically in Fig. 1, is characterized by the almost k -independent $\text{Im}\Sigma$. For a more complicated structure of $\text{Im}\Sigma$, further considerations based on $C_1(r_{TF})$ in Eq. (6) would be required. We are stressing that the confinement of ARPES spectral weight in Fig. 7(a) is much stronger ($R_1 = 1.85$), because of which one may incorrectly conclude




FIG. 7. (a) Raw ARPES data with the constant background removed. (b) $\text{Im}\Sigma$ corresponding to (a). (c) Measure of confinement $\mathcal{R}_1(k_x)/\text{MAX}\{\mathcal{R}_1(k_x)\}$ for data in (a) (black) and (b) (red) panel.

about the long-range character of the coupling to the LO3 phonons.

Conclusions. Our study provides a means to estimate the range/screening length of the EPI directly from the ARPES spectra, giving important insights into the polaronic correlations present in a wide class of real systems involving many puzzling physical phenomena. The \mathbf{k} and ω dependencies of the first and the second sideband are analyzed in detail as a function of screening, in terms of exact leading corrections obtained from the diagrammatic expansion, including the leading vertex correction. It is shown that the range of the EPI may be extracted from the confinement of $\text{Im}\Sigma$ within the Brillouin zone, even when the experimental resolution is very limited. Our results apply to all recently investigated

low-density metallic systems, characterized by the screened Fröhlich interaction, while our methodology is not restricted to phonons only, since coupling to excitations of other nature (plasmons, magnons, paramagnons, charge-order fluctuations, etc.) may be analyzed in a similar manner.

Acknowledgments. N.N. and A.S.M. acknowledge the support by JST CREST Grant No. JPMJCR1874, Japan. J.K. acknowledges the support by the Croatian Science Foundation Project IP-2016-06-7258. O.S.B. acknowledges the support by the QuantiXLie Center of Excellence, a project co-financed by the Croatian Government and European Union through the European Regional Development Fund - the Competitiveness and Cohesion Operational Programme (Grant No. KK.01.1.1.01.0004).

-
- [1] A. Alexandrov, *Theory of Superconductivity: From Weak to Strong Coupling* (CRC Press, Boca Raton, FL, 2003).
- [2] C. Chen, J. Avila, S. Wang, Y. Wang, M. Mucha-Kruczyński, C. Shen, R. Yang, B. Nosarzewski, T. P. Devereaux, G. Zhang *et al.*, *Nano Lett.* **18**, 1082 (2018).
- [3] Y. Wang, K. Nakatsukasa, L. Rademaker, T. Berlijn, and S. Johnston, *Supercond. Sci. Technol.* **29**, 054009 (2016).
- [4] S. Moser, L. Moreschini, J. Jaćimović, O. S. Barišić, H. Berger, A. Magrez, Y. J. Chang, K. S. Kim, A. Bostwick, E. Rotenberg *et al.*, *Phys. Rev. Lett.* **110**, 196403 (2013).
- [5] C. Chen, J. Avila, E. Frantzeskakis, A. Levy, and M. C. Asensio, *Nat. Commun.* **6**, 8585 (2015).
- [6] S. Moser, S. Fatale, P. Krüger, H. Berger, P. Bugnon, A. Magrez, H. Niwa, J. Miyawaki, Y. Harada, and M. Grioni, *Phys. Rev. Lett.* **115**, 096404 (2015).
- [7] R. Yukawa, K. Ozawa, S. Yamamoto, H. Iwasawa, K. Shimada, E. F. Schwier, K. Yoshimatsu, H. Kumigashira, H. Namatame, M. Taniguchi *et al.*, *Phys. Rev. B* **94**, 165313 (2016).
- [8] C. Cancellieri, A. S. Mishchenko, U. Aschauer, A. Filippetti, C. Faber, O. S. Barišić, V. A. Rogalev, T. Schmitt, N. Nagaosa, and V. N. Strocov, *Nat. Commun.* **7**, 10386 (2016).
- [9] Z. Wang, S. McKeown Walker, A. Tamai, Y. Wang, Z. Ristic, F. Y. Bruno, A. de la Torre, S. Riccò, N. C. Plumb, M. Shi *et al.*, *Nat. Mater.* **15**, 835 (2016).
- [10] C. Verdi, F. Caruso, and F. Giustino, *Nat. Commun.* **8**, 15769 (2017).
- [11] C. Zhang, Z. Liu, Z. Chen, Y. Xie, R. He, S. Tang, J. He, W. Li, T. Jia, S. N. Rebec *et al.*, *Nat. Commun.* **8**, 14468 (2017).
- [12] J. M. Riley, F. Caruso, C. Verdi, L. B. Duffy, M. D. Watson, L. Bawden, K. Volckaert, G. van der Laan, T. Hesjedal, M. Hoesch *et al.*, *Nat. Commun.* **9**, 2305 (2018).
- [13] M.-A. Husanu, L. Vistoli, C. Verdi, A. Sander, V. Garcia, J. Rault, F. Bisti, L. L. Lev, T. Schmitt, F. Giustino *et al.*, *Commun. Phys.* **3**, 62 (2020).
- [14] F. Giustino, *Rev. Mod. Phys.* **89**, 015003 (2017).
- [15] G. Antonius, S. Poncé, E. Lantagne-Hurtubise, G. Auclair, X. Gonze, and M. Côté, *Phys. Rev. B* **92**, 085137 (2015).
- [16] S. Ponc, E. Margine, C. Verdi, and F. Giustino, *Comput. Phys. Commun.* **209**, 116 (2016).
- [17] S. M. Story, J. J. Kas, F. D. Vila, M. J. Verstraete, and J. J. Rehr, *Phys. Rev. B* **90**, 195135 (2014).
- [18] B. Gumhalter, V. Kovač, F. Caruso, H. Lambert, and F. Giustino, *Phys. Rev. B* **94**, 035103 (2016).
- [19] D. Dunn, *Can. J. Phys.* **53**, 321 (1975).
- [20] F. Stern and W. E. Howard, *Phys. Rev.* **163**, 816 (1967).
- [21] M. Portnoi and I. Galbraith, *Solid State Commun.* **103**, 325 (1997).
- [22] H. Ouerdane, *J. Appl. Phys.* **110**, 074905 (2011).
- [23] F. M. Peeters, Wu Xiaoguang, and J. T. Devreese, *Phys. Rev. B* **33**, 3926 (1986).
- [24] J. T. Devreese and A. S. Alexandrov, *Rep. Prog. Phys.* **72**, 066501 (2009).
- [25] A. S. Alexandrov and J. Ranninger, *Phys. Rev. B* **45**, 13109 (1992).
- [26] G. Mahan, *Many-Particle Physics* (Springer US, New York, 2012).
- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.102.121108> for: i) the derivation of the leading self-energy contributions to the first and second phonon sidebands and, ii) a realistic example of data processing from raw ARPES to the imaginary part of the electron-phonon self-energy.
- [28] A. Chikina, F. Lechermann, M.-A. Husanu, M. Caputo, C. Cancellieri, X. Wang, T. Schmitt, M. Radovic, and V. N. Strocov, *ACS Nano* **12**, 7927 (2018).