

# The enigmatic multiple star VV Ori

---

**Budding, Edwin; Southworth, John; Pavlovski, Krešimir; Rhodes, Michael D; Zihao, Wu; Love, Tom; Blackford, Mark G; Banks, Timothy S; Alexander, Murray**

Source / Izvornik: **Monthly Notices of the Royal Astronomical Society, 2024, 527, 6389 - 6405**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

<https://doi.org/10.1093/mnras/stad3586>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:685681>

Rights / Prava: [Attribution 4.0 International](#) / [Imenovanje 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-08-16**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



# The enigmatic multiple star VV Ori

Edwin Budding<sup>1,2</sup>, John Southworth<sup>1,3</sup>★, Krešimir Pavlovski<sup>4</sup>, Michael D. Rhodes<sup>5</sup>, Wu Zihao<sup>6</sup>, Tom Love<sup>7</sup>, Mark G. Blackford<sup>8</sup>, Timothy S. Banks<sup>9,10</sup> and Murray Alexander<sup>11</sup>

<sup>1</sup>*Carter Observatory, 40 Salamanca Road, Kelburn, Wellington 6012, New Zealand*

<sup>2</sup>*School of Chemical & Physical Sciences, Victoria University of Wellington, PO Box 600, Wellington 6140, New Zealand*

<sup>3</sup>*Astrophysics Group, Keele University, Staffordshire, ST5 5BG, UK*

<sup>4</sup>*Department of Physics, Faculty of Science, University of Zagreb, Bijenička cesta 32, 10000 Zagreb, Croatia*

<sup>5</sup>*Brigham Young University, Provo, Utah 84602, USA*

<sup>6</sup>*Department Statistics & Data Science, National University of Singapore, 6 Science Drive 2, 117546, Singapore*

<sup>7</sup>*Variable Stars South, RASNZ, PO Box 3181, Wellington 6011, New Zealand*

<sup>8</sup>*Variable Stars South, Congarinni Observatory, Congarinni, NSW 2447, Australia*

<sup>9</sup>*Nielsen, 675 6th Ave, New York, NY 10011, USA*

<sup>10</sup>*Department of Physical Science & Engineering, Harper College, 1200 W Algonquin Rd, Palatine, IL 60067, USA*

<sup>11</sup>*Physics Department, University of Winnipeg, 515 Portage Avenue, Winnipeg R3B 2E9, Canada*

Accepted 2023 November 10. Received 2023 November 10; in original form 2023 September 12

## ABSTRACT

New photometry, including *TESS* data, have been combined with recent spectroscopic observations of the Orion Ib pulsating triple-star system VV Ori. This yields a revised set of absolute parameters with increased precision. Two different programmes were utilized for the light-curve analysis, with results in predictably close agreement. The agreement promotes confidence in the analysis procedures. The spectra were analysed using the FDBINARY programme. The main parameters are as follows:  $M_1 = 11.56 \pm 0.14$  and  $M_2 = 4.81 \pm 0.06$  ( $M_\odot$ ). We estimate an approximate mass of the wide companion as  $M_3 = 2.0 \pm 0.3 M_\odot$ . Similarly,  $R_1 = 5.11 \pm 0.03$ ,  $R_2 = 2.51 \pm 0.02$ ,  $R_3 = 1.8 \pm 0.1$  ( $R_\odot$ );  $T_{e1} = 26600 \pm 300$ ,  $T_{e2} = 16250 \pm 420$ , and  $T_{e3} = 10000 \pm 1000$  (K). The close binary's orbital separation is  $a = 13.91$  ( $R_\odot$ ); its age is  $8 \pm 2$  (Myr) and its photometric distance is  $396 \pm 7$  pc. The primary's  $\beta$  Cep type oscillations support these properties and confirm our understanding of its evolutionary status. Examination of the well-defined  $\lambda 6678$  He I profiles reveals the primary to have a significantly low projected rotation: some 80 per cent of the synchronous value. This can be explained on the basis of the precession of an unaligned spin axis. This proposal can resolve also observed variations of the apparent inclination and address other longer term irregularities of the system reported in the literature. This topic invites further observations and follow-up theoretical study of the dynamics of this intriguing young multiple star.

**Key words:** binaries: close – stars: early-type – stars: variables:  $\beta$  Cep type – stars: individual: VV Ori.


## 1 INTRODUCTION

VV Ori is a very bright ( $V \sim 5.4$ ,  $M_V \sim -2.8$  mag) and well-known eclipsing binary in the Belt, or  $\epsilon$  Ori, grouping of the Orion Ib association (Blaauw 1964; Wright 2020). Superficially, it resembles the system V Pup, with its period of  $\sim 1.5$  d and B1 + B3–5 type components (Budding et al. 2021). However, numerous studies of the close pair in VV Ori have found a detached arrangement of young early-type main sequence stars (Fig. 1), unlike the semidetached configuration of V Pup. As the binary's orbital plane is close to the line of sight, a succession of complete (or nearly so) eclipses is observed, allowing confidence about the determination of fitting function parameters, including those relating to the distribution of brightness over the stellar surfaces.

Massive young stars like those in VV Ori are frequently found in groups of multiple-star systems (Sana et al. 2012). Such star formation regions are thought to have an important role in determining the long-term behaviour of galaxies (Langer 2012; Zucker et al. 2022). A striking recent discovery about VV Ori, enabled by the long duration and high quality of *TESS* photometry (Ricker et al. 2014), is the existence of low-level  $\beta$  Cep-type pulsations that may be associated with the primary star (Southworth, Bowman & Pavolvski 2021).

It is well known that essential properties of stars can be determined from combining the analysis of photometric data with spectroscopic data – the ‘eclipse method’. We follow that approach for VV Ori. This has been more recently enhanced by the understanding of stellar pulsational behaviour or asteroseismology (Aerts, Christensen-Dalsgaard & Kurtz 2010; Murphy 2018, Bowman 2020). A decomposition of the pulsations into constituent oscillation modes will furnish direct information on the stellar structure, particularly if constraints are imposed by membership of a co-evolutionary group of nearby stars (Lampens 2006).

\* E-mail: [taylorssouthworth@gmail.com](mailto:taylorssouthworth@gmail.com)



**Figure 1.** WEBDA colour–magnitude diagram based on the data of Warren & Hesser (1977). The Padova isochrone for  $\log(\text{age}) = 6.78$  yr is shown with solar composition, reddening  $E(B - V) = 0.05$  and distance modulus 8.16. VV Ori is marked by a red-coloured star.

With regard to early (B type) stellar pulsators, those of the  $\beta$  Cep (Lesh 1978) and slowly pulsating (SPB) kinds (Waelkens 1991) are well known.  $\beta$  Cep variables show low-order gravity ( $g$ ) and pressure ( $p$ ) oscillation modes, with typical brightness variations of order 0.1 mag. They are associated with pulsation periods in the range 2–6 h (Stankov & Handler 2005) and masses of around 8–15  $M_{\odot}$ . SPB pulsations have quite longer periods, generally a few days, and are found in the lower-mass range of 3–9  $M_{\odot}$ . The variability is believed to result from high-order  $g$ -modes. More recent space-based observations, however, have uncovered both  $p$ - and  $g$ -type low-amplitude pulsations in many massive stars outside the foregoing ranges (Pedersen et al. 2019; Bursens et al. 2020).

Combining regular light and radial velocity (RV) curve analyses with astroseismology, especially with recently increased data accuracy, permits stellar masses and radii to be more confidently specified (Ratajczak, Pigulski & Pavlovski 2017; Southworth et al. 2020; Southworth, Bowman & Pavolvski 2021; Salmon et al. 2022). That being said, stellar oscillations in massive eclipsing binaries have only been definitely established in relatively few cases so far (e.g. Clausen 1996, Southworth & Bowman 2022, Erdem et al. 2022). Stellar structural parameters may also be checked from eccentric close binary systems through non-Keplerian (apsidal) motion that relates to the scale of their proximity effects (Sterne 1939; Russell 1942; Kopal 1959). For more recent discussion, see also Welsh et al. (2011); Hambleton et al. (2013); Feiden (2015); Bowman et al. (2019); Handler et al. (2020); Kurtz et al. (2020) and Fuller et al. (2020). A review of the use of space-based photometry for binary-star science can be found in Southworth (2021).

VV Ori contains components of spectral types B1 V and mid-B V in an essentially circular orbit of period 1.485 d. Its eclipsing nature was discovered in 1903 (Barr 1905). The early history of its study was summarized by Wood (1946) and Duerbeck (1975). Two of the more recent published investigations of the system parameters are those of Sarma & Vivekananda (1995) and Terrell, Munari & Siviero (2007).

Multiplicity, beyond the two eclipsing stars, has been discussed in a number of studies over the years. Excess scatter in early-RV data led to cautious suggestions of a third body with an orbital period of  $\sim 120$  d (Daniel 1915; Struve & Luyten 1949), a proposal that has been disfavoured in later papers (Terrell, Munari & Siviero 2007; Van Hamme & Wilson 2007). These latter papers cast doubt on the evidence for a close third body, at least regarding the RV analyses, though various photometric studies (Budding & Najim 1980; Chambliss 1983; Van Hamme & Wilson 2007) admitted a small third light contribution. However, Horch et al. (2017) have resolved a companion at an angular separation of 0.23 arcsec using speckle interferometry. The magnitude differences between this companion and the binary system are 3.88 mag at 692 nm and 3.43 mag at 880 nm. Applying a linear photometric gradient approximation (Golay 1974) would yield a magnitude difference of about 4.45 in the V range, or about 0.017 of the system’s light, which is essentially the same value found by Chambliss (1983). At a distance of  $441 \pm 22$  pc (Gaia Collaboration et al. 2023), the separation corresponds to a projected distance of  $\sim 87$  au, and thus a minimum orbital period of around 200 yr. This resolved companion cannot, therefore, be directly responsible for the putative 120 d orbital variations.

Recently, Southworth, Bowman & Pavolvski (2021) noted that the succession of well-observed minima, over the years, has shown a transition between complete and partial eclipses. This can be associated with a systematic variation of the apparent inclination of the binary orbit to the line of sight. That there may be some variations of shorter period than that of the wide orbit is thus still an open question.

A consensus from RV studies (Daniel 1915; Struve & Luyten 1949; Beltrami & Galeotti 1969; Duerbeck 1975; and Popper 1993) presented the system as mostly single-lined, though Beltrami & Galeotti (1969) were able to model blended line profiles to derive a velocity separation and thence a mass ratio of  $\sim 0.45$ ; similar to that of Duerbeck (1975).

In this study, we bring new evidence to bear on outstanding uncertainties about the system, regarded as important for fixing the properties of massive early-type stars (Eaton 1975; Popper 1993; Terrell, Munari & Siviero 2007; Van Hamme & Wilson 2007; Southworth, Bowman & Pavolvski 2021). We examine new high-quality photometry from the *TESS* satellite, and high-dispersion spectroscopy from the FIES spectrograph of the 2.56-m Nordic Optical Telescope (NOT), Tenerife, Spain; as well as the HERCULES spectrograph of the University of Canterbury Mt John Observatory, New Zealand. These new data yield important revisions in the physical properties of the component stars. We use up-to-date analytical tools to bear on the data analysis, and check on evolutionary models, keeping in mind additional evidence relating to the Orion OB 1b membership and the degree of multiplicity of VV Ori.

## 2 PHOTOMETRY

Table 1 presents a summary of the main parameters derived in previous photometric studies of VV Ori light curves (LCs) at visual (V) wavelengths. Standard notation is used, that is,  $L_n$  for fractional luminosities (where  $n$  indicates a given star in the system),  $r_n$  for relative radii,  $i$  for the orbital inclination, and  $u_n$  for linear limb-darkening coefficients. The sources of these parameters are as follows: Du-75 = Duerbeck (1975); Na-81 = Budding & Najim (1980); Ch-84 = Chambliss (1984); Sa-95 = Sarma & Vivekananda (1995); Te-07 = Terrell, Munari & Siviero (2007); So-21 = Southworth, Bowman & Pavolvski (2021). Some variations of parameter values can be noticed across Table 1, but these are generally in keeping with

**Table 1.** Reference parameters for historic photometric model fits.

| Parameter    | Du-75             | Na-80            | Ch-84            | Sa-95            | Te-07             | So-21             |
|--------------|-------------------|------------------|------------------|------------------|-------------------|-------------------|
| $\lambda$ nm | 540               | 425              | 530              | 530              | (530)             | <i>TESS</i>       |
| $M_2/M_1$    | 0.45              | 0.45             | 0.42             | 0.42             | 0.38              | 0.376             |
| $L_1$        | $0.878 \pm 0.06$  | $0.926 \pm 0.09$ | $0.891 \pm 0.10$ | $0.878 \pm 0.01$ | $0.908 \pm 0.09$  | $0.892 \pm 0.07$  |
| $L_2$        | $0.122 \pm 0.04$  | $0.069 \pm 0.03$ | $0.092 \pm 0.05$ | $0.122 \pm 0.01$ | $0.098 \pm 0.05$  | $0.103 \pm 0.005$ |
| $L_3$        | $0.000 \pm 0.06$  | $0.005 \pm 0.03$ | $0.017 \pm 0.11$ | $0.00 \pm 0.01$  | $0.00 \pm 0.07$   | $0.005 \pm 0.07$  |
| $r_1$        | $0.366 \pm 0.006$ | $0.371 \pm 0.01$ | $0.363 \pm 0.02$ | $0.378 \pm 0.02$ | $0.369 \pm 0.005$ | $0.372 \pm 0.001$ |
| $r_2$        | $0.180 \pm 0.01$  | $0.168 \pm 0.03$ | $0.176 \pm 0.05$ | $0.179 \pm 0.05$ | $0.179 \pm 0.013$ | $0.182 \pm 0.003$ |
| $i$          | $85.6 \pm 1.0$    | $85.8 \pm 1$     | $85.6 \pm 2$     | $86.1 \pm 5$     | $85.9 \pm 0.5$    | $78.3 \pm 0.5$    |
| $T_h$ (K)    | 25 000            | 25 000           | 25 000           | 25 000           | 26 199            | 26 200            |
| $T_c$ (K)    | 15 000            | 16 000           | 15 700           | 15 500           | 16 073            | 16 100            |
| $u_1$        | 0.07              | 0.37             | 0.28             | 0.33             | –                 | 0.63 (bol)        |
| $u_2$        | 0.45              | 0.45             | 0.32             | 0.37             | –                 | 0.71 (bol)        |

**Table 2.** Summary of Congarinni BVR photometry of VV Ori, along with uncertainties, given as standard deviations.


| Date     | Phase | V     | Standard deviation | B – V  | Standard deviation | V – R  | Standard deviation |
|----------|-------|-------|--------------------|--------|--------------------|--------|--------------------|
| 1901 26  | 0.0   | 5.627 | 0.034              | –0.140 | 0.042              | –0.113 | 0.046              |
| 18 12 25 | 0.5   | 5.465 | 0.017              | –0.157 | 0.024              | –0.114 | 0.021              |
| 18 12 28 | 0.5   | 5.467 | 0.016              | –0.166 | 0.021              | –0.113 | 0.023              |
| 18 12 29 | 0.25  | 5.308 | 0.022              | –0.157 | 0.029              | –0.114 | 0.034              |
| 1901 25  | 0.25  | 5.314 | 0.022              | –0.157 | 0.032              | –0.104 | 0.035              |
| 18 12 29 | 0.75  | 5.281 | 0.020              | –0.158 | 0.028              | –0.125 | 0.021              |

the listed uncertainties. This table is not intended to be exhaustive, but it gives an impression of reference quantities derived from data analysis over the last several decades, keeping in mind concomitant increases of data accuracy.

Further multicolour photometry of VV Orionis (see Table 2) was carried out over 10 nights between December 2018 and February 2019 from the Congarinni Observatory, NSW, Australia ( $152^\circ 52' \text{ E}$ ,  $30^\circ 44' \text{ S}$ , 20 m above mean sea level). Images were captured with an ATIK™ One 6.0 CCD camera equipped with Johnson–Cousins BVR filters attached to an 80 mm f6 refractor, which, given the brightness of the stars, was stopped down to 50-mm aperture. MaxIm DL™ software was used for image handling, calibration, and aperture photometry. HD 36779 was used as the main comparison star. Its magnitude and colours were determined as  $V = 6.223$ ,  $B - V = -0.165$ , and  $V - R = -0.067$ , in close agreement with the Johnson 11-colour catalogue (Ducati 2002). The derived LCs are shown in Fig. 2. Optimal parameters from WINFITTER modelling (Rhodes 2022) are given in Table 3. Using the results given in Table 2 as a guide, we derived BVR magnitudes of the three identified photometric components of VV Ori as: 5.23, 8.00, 9.95 (B); 5.41, 8.02, 9.79 (V); 5.65, 8.09, 9.61 (R). Taking into account local variations of the mean reddening ( $\sim 0.09$ , according to Warren & Hesser, 1978) and errors of measurement, these figures are in fair accord with the assigned early- and mid-B main sequence types for the close pair attended by a cooler, A-F type, companion.

## 2.1 Light-curve analysis

We first sought to match the form of the LC with the well-known Wilson–Devinney (WD) code (Wilson & Devinney 1972; Wilson 1979). We used the 2004 version of the code (Wilson & van Hamme 2004) implemented using the JKTWD wrapper (Southworth et al. 2011). This code adopts Roche model geometry for the calculation


**Figure 2.** WINFITTER model fitted to recent BVR LCs from the Congarinni Observatory. The V and R LCs have been vertically offset by 0.15 and 0.30 to allow presentation on a single chart.

of the shapes of binary stars (Kopal 1959), although it becomes computationally expensive to perform on the original complete set of *TESS* observations. We therefore took the data from sector 6, converted them to orbital phase, and averaged them into 400 equally spaced phase bins.

These data were then fitted using WD to obtain the simplest initial model that matched them well. This, our default solution, was obtained by fitting the following model parameters: surface potentials of the two stars, orbital inclination, phase of the primary mid-eclipse, and light contributions of both eclipsing stars together with a possible third light. The third light is expressed as a fraction of the total brightness of the system at phase 0.25. The light contributions from the two stars are expressed in the WD code on a different flux scale, and add to  $4\pi$  minus the third light for zero differential magnitude. The fitting was performed in mode 0 (see Table 4), where the effective temperature ( $T_e$ ) values of the two stars are fixed and their light contributions matched directly. We therefore fixed the  $T_e$  values and


**Table 3.** Parameters for model fits to the BVR photometry shown in Fig. 2. Adopted parameters are listed below the 3rd row horizontal line. The geometric elements  $r_1$ ,  $r_2$ , and  $i$ , are weighted averages from initial fittings at the separate wavelengths. The value of  $L_3$  in the B filter, being expectably less than its uncertainty as an optimal fitting result, is here a fixed hyperparameter.

| Parameter       | B                 | V                 | R                 |
|-----------------|-------------------|-------------------|-------------------|
| $L_1$           | $0.926 \pm 0.012$ | $0.901 \pm 0.013$ | $0.894 \pm 0.013$ |
| $L_2$           | $0.072 \pm 0.003$ | $0.082 \pm 0.003$ | $0.085 \pm 0.003$ |
| $L_3$           | 0.012             | $0.016 \pm 0.014$ | $0.021 \pm 0.013$ |
| $r_1$ (mean)    |                   | $0.359 \pm 0.008$ |                   |
| $r_2$ (mean)    |                   | $0.171 \pm 0.004$ |                   |
| $i$ (deg, mean) |                   | $80.0 \pm 0.8$    |                   |
| $M_2/M_1$       | 0.42              |                   |                   |
| $T_h$ (K)       | 26600             |                   |                   |
| $T_c$ (K)       | 16250             |                   |                   |
| $u_1$           | 0.28              | 0.25              | 0.21              |
| $u_2$           | 0.29              | 0.29              | 0.24              |
| $\chi^2/\nu$    | 1.01              | 1.05              | 0.90              |
| $\Delta l$      | 0.014             | 0.012             | 0.010             |

**Table 4.** Summary of the parameters for the WD fittings of the *TESS* sector 6, phase-binned LC of VV Ori. Detailed descriptions of the control parameters can be found in the WD code user guide (Wilson & van Hamme 2004). The uncertainties have been determined via comparisons through a wide range of model fits.

| Parameter                              | WD2004 name | Value               |
|--|-------------|---------------------|
| <i>Control &amp; fixed parameters:</i> |             |                     |
| WD2004 operation mode                  | MODE        | 0                   |
| Treatment of reflection                | MREF        | 1                   |
| Number of reflections                  | NREF        | 1                   |
| LD law                                 | LD          | 2 (logarithmic)     |
| Numerical grid size (normal)           | N1, N2      | 60                  |
| Numerical grid size (coarse)           | N1L, N2L    | 60                  |
| <i>Fixed parameters:</i>               |             |                     |
| Mass ratio                             | RM          | 0.418               |
| Orbital eccentricity                   | E           | 0.0                 |
| $T_e$ of primary (K)                   | TAVH        | 26,660              |
| $T_e$ of secondary (K)                 | TAVH        | 16,250              |
| Bolometric albedos                     | ALB1, ALB2  | 1.0, 1.0            |
| Rotation rates                         | F1, F2      | 1.0, 1.0            |
| Gravity darkening                      | GR1, GR2    | 1.0, 1.0            |
| Logarithmic LD coefficients            | Y1A, Y2A    | 0.217, 0.205        |
| <i>Fitted parameters:</i>              |             |                     |
| Primary potential                      | PHSV        | $3.220 \pm 0.026$   |
| Secondary potential                    | PHSV        | $3.637 \pm 0.023$   |
| Orbital inclination (deg)              | XINCL       | $79.77 \pm 0.19$    |
| Primary light contribution             | HLUM        | $10.48 \pm 0.10$    |
| Secondary light contribution           | CLUM        | $1.215 \pm 0.012$   |
| Third light                            | EL3         | $0.071 \pm 0.008$   |
| Primary linear LD coefficient          | X1A         | $0.359 \pm 0.049$   |
| Secondary linear LD coefficient        | X2A         | $0.22 \pm 0.15$     |
| Fractional radius of primary           |             | $0.3656 \pm 0.0019$ |
| Fractional radius of secondary         |             | $0.1794 \pm 0.0015$ |

mass ratio at the adopted spectroscopic values, presented in Section 3, and assumed a circular orbit. We also assumed, for the present purpose, synchronous rotation, as well as albedo parameters and gravity-darkening exponents set to unity. The limb darkening (LD) effect was accounted for using a logarithmic law with one coefficient adjustable and the other fixed, for both stars. Maximum numerical precision was used, by setting the quantities N1, N2, N1L, and N2L




**Figure 3.** The adopted optimal WD model (green line) to the *TESS* sector 6 phase-binned LC of VV Ori (red-filled circles). The residuals of the fit are plotted in the lower panel using a greatly enlarged y-axis to bring out the detail.

to 60. For the effective wavelength of observation, the Cousins  $R$  band was adopted as it is the closest available approximation to the *TESS* passband. The ‘simple’ option for the reflection effect was chosen.

With these settings in place, we were able to obtain a satisfactory fit to the phase-binned data as shown in Fig. 3. The residuals are shown on a larger scale to demonstrate the presence of structure remaining in the phase-binned data. Whilst the differences between observed and calculated LCs are relatively small (an rms of 0.3 mmag), the remaining pulsational signature means the residuals are dominated by red noise. Using fitting procedures that allowed for orbital eccentricity, we conclude that the quantity  $e \cos \omega$ , where  $e$  is the eccentricity and  $\omega$  the longitude of periastron, must be smaller than 0.001. A summary of the fitted and fixed parameters is given in Table 4. The fractional stellar radii specified in Table 4 are volume-equivalent values. Table 3-2 in Kopal’s (1959) book shows these radii to be one or two per cent larger than the corresponding unperturbed radii ( $r_0$ ) at the same mass ratio.

Determination of the uncertainties of the fitted parameters is not trivial. We follow the approach of Southworth (2020) on this. The uncertainties derive almost entirely from the model limitations, even taking into account the residual pulsation signature, because the precision of the data is extremely high, showing up systematic effects that are not in the model. We therefore ran WD with selected differences in parameter values to determine their effects on the resulting fits, as follows:

- (i) changing the mass ratio by the uncertainty in the spectroscopic value;
- (ii) fixing  $e \sin \omega = 0.005$ , instead of assuming a circular orbit;
- (iii) decreasing the rotation parameters for the stars by 10 per cent;
- (iv) changing the albedos by 10 per cent;
- (v) changing the gravity-darkening exponents by 10 per cent;
- (vi) lowering the N1, N2 values from 60 to 55 in steps of 1;
- (vii) fitting for  $T_e$  in mode 2, instead of fitting for HLUM and CLUM in mode 0;
- (viii) using the more detailed reflection model;
- (ix) specifying the square root instead of the logarithmic LD law;
- (x) using the Cousins  $I$  filter for the *TESS* pass-band;



**Figure 4.** The upper panel presents an optimal model LC to normalized SAP flux measures from *TESS* for the orbit 13 of Sector 6, TBJD 1487.38930345–1488.87467768, 1070 data points were taken from the original source with no binning. The lower panel reveals the  $\beta$  Cep type behaviour in the residuals.

- (xi) fixing third light to be zero; and
- (xii) changing the number of phase bins from 400 to 300 or 500.

The net result of these tests was a compilation of values for the fitted parameters; each group associated with a different model run. We rejected all runs where the fit was significantly worse (more than 0.35 mmag rms in the residuals), and added all the parameter differences in quadrature to determine the uncertainty for each fitted parameter. These are given in Table 4. The largest contributions to these uncertainties were found to come from the effects of the adopted rotation, albedo, and eccentricity parameters. The derived uncertainty percentages are nonetheless still small, and the fitted parameters have highly consistent values.

For an independent assessment of the LC we used the programme WinFitter (WF), discussed in Budding & Demircan (2022) chapter 7. The algebraic form of its fitting function means that parameter space can be searched rapidly and thoroughly using relatively large data sets. An example of a LC from *TESS* Sector 6 is shown in Fig. 4 together with an optimal WF model. The residuals are shown in the lower panel where the  $\beta$  Cep oscillations of the primary component are clearly evident. The adopted light elements for this data set were Min I = TBJD1487.389303 + 1.48537742E.

LCs from both Sectors 6 and 32 were collected together, phased and binned using this ephemeris. The Sector 6 data were binned with a reduction factor of 14:1 down from 14 847 original data points, while Sector 32 data were binned with the ratio 17:1, from the 17 917 source data. Numerous optimal fitting experiments were performed on these reduced data sets. Starting values were guided by historic findings. The finally adopted parameter sets are given in Table 5, and are in reasonable agreement with the WD model parameters. The separately derived main geometric elements ( $r_1$ ,  $r_2$ ,  $i$ ) of the WD and WF fittings to the *TESS* data sets are within the uncertainty estimates of each other. They concur that the *TESS* data point to a distinctly lower inclination than the values cited in Table 1.

**Table 5.** Optimal WF parametrization of *TESS* photometry from Sectors 6 and 32, modelling the entire sectors. The data sets were phased by the orbital period, thus averaging out the pulsations. The column titled ‘Fig. 4’ lists the parameter estimates for the data period plotted in Fig. 4. The mass ratio, stellar temperatures, and LDs are held constant across these fits.

| Parameter    | Sector 6          | Sector 32         | Fig. 4            |
|--------------|-------------------|-------------------|-------------------|
| $L_1$        | $0.847 \pm 0.010$ | $0.875 \pm 0.006$ | $0.860 \pm 0.013$ |
| $L_2$        | $0.098 \pm 0.003$ | $0.096 \pm 0.002$ | $0.100 \pm 0.004$ |
| $L_3$        | $0.055 \pm 0.005$ | $0.008 \pm 0.003$ | $0.040 \pm 0.005$ |
| $r_1$        | $0.371 \pm 0.002$ | $0.370 \pm 0.002$ | $0.368 \pm 0.002$ |
| $r_2$        | $0.183 \pm 0.003$ | $0.184 \pm 0.003$ | $0.181 \pm 0.001$ |
| $i$ (deg)    | $78.2 \pm 0.4$    | $78.0 \pm 0.4$    | $79.0 \pm 0.5$    |
| $M_2/M_1$    |                   | 0.48              |                   |
| $T_h$ (K)    |                   | 26 600            |                   |
| $T_c$ (K)    |                   | 16 250            |                   |
| $u_1$        |                   | 0.18              |                   |
| $u_2$        |                   | 0.20              |                   |
| $\chi^2/\nu$ | 1.14              | 0.97              | 0.90              |
| $\Delta l$   | 0.0013            | 0.002             | 0.0025            |

## 2.2 Fitting function with unaligned axes

It was mentioned earlier that Southworth, Bowman & Pavlovski (2021) discussed visible LC effects associated with an apparent variation of the inclination parameter. The precession of a spin axis that is unaligned to that of the orbit is feasible for a young multiple system in a process of angular momentum evolution, and, since the flux from the system is dominated by the primary star, such effects would be mainly associated with the behaviour of that star. In this section, we check on the scale of relevant effects and how the WF fitting function may address this topic.

In general, the matching of a theoretical model to a photometric data set (LC) is achieved by optimizing the agreement of a fitting function  $l(\phi; a_i)$ , where  $\phi$  is the orbital phase and  $a_i$  are a set of parameters whose adjustment is implied by the optimization process. This fitting function can be set in the form

$$l(\phi) = \int_{A_1} J_1 dA_1 + \int_{A_2} J_2 dA_2 - \int_{A_e} J_e dA_e, \quad (1)$$

where the suffix 1 refers to the ‘primary’ object, with higher surface temperature, say, having projected surface area  $A_1$ , while suffix 2 denotes the secondary, projecting area  $A_2$  perpendicular to the line of sight. The received flux scales linearly with the integrated locally projected flux  $J$  from either source. The suffix  $e$  relates to an eclipse: the relevant term is zero when there is no eclipse and maximizes during complete eclipse phases. Eclipses of the secondary star are covered by the same form, but with the eclipsing and eclipsed roles reversed,  $J_e$  being suitably re-assigned.

Equation (1) is reducible to a relatively simple form in the ‘spherical model’ (Russell & Shapley 1912). This reduction suggests that modelling can be developed to cater for more realistic situations using the Taylor expansion

$$l(\phi) = l_0(\phi) + \Delta l_{A_1}(\phi) + \Delta l_{A_2}(\phi) - \Delta l_{A_e}(\phi), \quad (2)$$

where  $l_0(\phi)$  in equation (2) would be normalized so that the sum of the first two integrals, in the unperturbed, uneclipsed situation, is unity. In this case, the first term becomes the primary’s fractional luminosity  $L_1$ , the second, correspondingly,  $L_2$ .

We can adopt, without loss of generality, that suffix 1 refers to the star about to be eclipsed and the zero-order LC equation becomes

$$l_0(\phi) = 1 - \alpha\{u_1, k, d(r_1, i, \phi)\}L_1, \quad (3)$$

where  $\alpha$  is Russell's (1912a, b) light loss function, depending here on the renormalized separation  $d = \delta/r_1$ ;  $\delta$  being the separation of the two star centres in units of the mean orbital radius on the tangential sky plane, given by

$$\delta^2 = \sin^2 \phi \sin^2 i + \cos^2 i. \quad (4)$$

Other parameters include the ratio of radii  $r_2/r_1$ , written as  $k$  in equation (3), and a limb-darkening coefficient  $u_1$ .

For a more realistic model we should evaluate the effects of tidal and rotational perturbations of the photosphere, associated with the proximity of the two stars, at given phases, as well as the 'reflection' effects that result from their mutual irradiation. Calculating the surface distortion involves the radial displacements in the key directions of the line of centres and the axis of rotation. Local areal projections for any viewing angle are then resolved into their components in the observer's frame of reference.

Both WD and WF converge to the same approximation for the first-order surface perturbation given as

$$\frac{\Delta' r}{r_0} = q \sum_{j=2}^4 r_0^{j+1} (1 + 2k_j) P_j(\lambda) + nr_0^3 (1 - \nu^2), \quad (5)$$

where  $r$  is the local stellar radius expressed as a fraction of the orbital separation of the components with mean value  $r_0$ , and  $k_j$  are the well-known structural constants reflecting the distribution of matter through the star.  $\lambda$  is the direction cosine of the angle between the radius vector  $\hat{r}$  and the line of centres, and  $\nu$  is the direction cosine of the angle between  $\hat{r}$  and the rotation axis. The coefficients  $k_j$  can be taken from suitable stellar models, for example Inlek, Budding & Demircan (2017), or set to zero in the Roche model (equations 1-11 in ch.1, and 2-6 in ch. 3 of Kopal, 1959).


With the surface perturbations as serial harmonic functions, they are seen (equation 5) to start with terms of order  $r^3$ , so by the time we reach terms of order  $r^6$  the gravitational interaction of tides on tides would be taken into account. WF analysis, in which the mutual effect of perturbations on perturbations is neglected, therefore proceeds up to and including terms of order  $r^5$ . This implies three separate tidal terms in the fitting function.

Rotation with a constant angular velocity, as is usually assumed, requires only one source term. However, location of the rotation axis with respect to the orbit will, in general, involve two additional angular parameters:  $\epsilon$  the obliquity and  $\psi$  the precession angle. In the case of aligned rotation and orbit axes, the surface distortions due to rotation, as they appear to a remote observer, involve integrals factored by orientation-dependent direction cosines. These reduce down to the independent cosine of the angle between the line of sight and the rotation axis  $n_0$ . This becomes replaced, in the unaligned case, by  $n'_0$ , where


$$n'_0 = n_0 \cos \epsilon + \sin i \sin \epsilon \cos \psi \quad (6)$$

(equation (1.13) of Budding & Demircan 2022).

Fig. 5 plots the measured depths of the secondary minima over a succession of LCs from *TESS* sector 6 with mean epoch TJD 1477.5973. This comes from averaging the values of 13 individual flux measures – at phase 0.5 and the 6 points on either side. A trend of order 2 mmag becomes apparent over the 18-d time interval. LC modelling of the partial, but near total, secondary eclipse shows that this would necessitate a change of inclination significantly greater than the  $\sim 0.01^\circ$  (for 18 d) considered by Southworth, Bowman & Pavlovski (2021) after inspection of historic LCs. However, the residuals presented in Fig. 4 allow an expectation of point-to-point variations in Fig. 5 of several mmag, so this apparent short-term




**Figure 5.** A trend across a flux interval of several mmag can be seen in the succession of *TESS* monitoring of secondary minima in the Sector 6 data.



**Figure 6.** The effects of varying the precession angle of the primary star's spin axis, at a  $20^\circ$  obliquity, on the relative flux during secondary minimum.

trend is not separable from effects *not* due to inclination changes. Alternatively, LC modelling shows that variations of several mmag in the central depth of the secondary minimum would result from changes of the apparent inclination, that is,  $\arccos(n'_0)$ , of order  $1^\circ$ . Fig. 6 shows that this could be accounted for by precessional movement of order  $\sim 10^\circ$  on a feasible time-scale of  $\sim 10$  yr.

A further series of *TESS* LCs with mean epoch TJD 2186.0373, that is, 708.44 d later, were collected in sector 32. Unfortunately, this nearly 2-yr time baseline produces only a  $0.4^\circ$  shift in the inclination at the rate presented by Southworth, Bowman & Pavlovski, so comparable to the uncertainty in the estimated inclination values of Tables 4 and 5. The mean relative fluxes at mid-secondary minimum from the two sectors are  $0.8549 \pm 0.0015$  and  $0.8551 \pm 0.0014$ , which does not yet provide convincing support for a secular change in the relative depth of the secondary minimum over the two years separating the two *TESS* sectors involved.



**Figure 7.** Phase-binned versions of sectors 6 and 32 *TESS* observations (top) and their difference on a magnified scale (bottom).

### 2.3 Evaluation of changes between *TESS* sectors

The hypothesis of orbital evolution associated with a third body, or some other cause, leads naturally to the suggestion that the LC may change between the two *TESS* sectors. Southworth, Bowman & Pavolvski (2021) found a change in orbital inclination which in turn may change the eclipse depths. We therefore searched for these effects.


We first fitted the two *TESS* sectors simultaneously to obtain an orbital ephemeris which precisely matched these data sets without influence from other data. We then phase-binned each *TESS* sector into 400 bins (as done earlier). A plot of these sectors and the difference between them can be found in Fig. 7. A decrease in the eclipse depths is clear.

We then fitted both data sets using the WD code (see subsection 2.1). For each fit we adopted one approach to modelling the data sets, and analysed them separately. We performed a range of fits varying the numerical precision of the code, the treatment of rotation, albedo, and LD. The differences in orbital inclination across all fits averaged at close to  $0.17^\circ$  (mean) with a standard deviation of  $0.08^\circ$ . We therefore find a change in inclination to a significance level of  $2\sigma$  over a time period of 706 d. The LC clearly changes, but we are not yet able to quantify this with sufficient statistical significance. It is unfortunate that no further observations of VV Ori by *TESS* are currently scheduled.


An alternative explanation of the change in eclipse depth is imperfections in the data reduction. An obvious possibility is an error in the background subtraction which would shift the LC up or down by a fixed amount of flux. In this case, we would expect to see the difference between the *TESS* sectors (Fig. 7) appear as a scaled-down version of the LC. That is not the case: the differences between sectors occur only during the eclipses. This supports the possibility of changing apparent inclination in the eclipsing pair in the VV Ori system.

### 2.4 Frequency analysis of *TESS* data

Southworth, Bowman & Pavolvski (2021) performed a frequency analysis of pulsations in the Sector 6 *TESS* photometry of VV Ori, recorded in 2018–19. They found 51 significant frequency components between  $1.4$  and  $27.8 \text{ d}^{-1}$ , with two dominant pulsation



**Figure 8.** The first 2.5 d of *TESS* SAP from Sector 32. Out-of-eclipse regions used for frequency analysis are shown in red.




**Figure 9.** Residual pulsations in the inter-minimum phases after removing the eclipsing binary LC.

modes ( $\nu_1 = 9.1766 \pm 0.0001$  and  $\nu_2 = 9.0324 \pm 0.0002 \text{ d}^{-1}$ ). There were also several independent g-mode frequencies below  $3 \text{ d}^{-1}$ . These components were attributed to  $\beta$  Cep oscillations of the primary and SPB variations of the secondary, respectively.

Here, we have analysed Sector 32 pulsations recorded in 2020, using a different approach to removing the close binary contribution to the LC. Fig. 8 shows part of the simple aperture photometry (SAP) LC. For clarity, only the first 2.5 d are presented; however, all 27 d data were analysed. Only the out-of-eclipse observations (shown in red) were used in the frequency analysis, in order to avoid complications to the pre-whitening operation arising from eclipse effects. The proximity effects were removed empirically, by fitting 6th-order polynomials to the out-of-eclipse sections of the phased LCs. The maxima before and after primary eclipses were fitted separately. Residual pulsations after subtracting the polynomial models are shown in Fig. 9. We used the same methodology to analyse the Sector 6 data, allowing direct comparison with the results from Southworth, Bowman & Pavolvski (2021).

Frequency analysis of residual pulsations was performed using PERIOD04 software (Lenz & Breger 2004). Iterative pre-whitening identified 60 significant frequencies with amplitude signal-to-noise  $> 4$ . The frequencies, amplitudes, and phases are listed in Appendix Table A1. Fig. 10 presents the frequency spectra based on the residuals from the LC fittings to the 2018 and 2020 data. General features of the Sectors 6 and 32 amplitude spectra are similar. The two dominant frequencies reported in Southworth, Bowman & Pavolvski (2021) remain and with similar amplitudes. The frequencies we obtain are: (2018)  $9.17680 \pm 0.00007$  and  $9.03131 \pm 0.00016 \text{ d}^{-1}$ ; (2020)  $9.17726 \pm 0.00007$  and  $9.03282 \pm 0.00017 \text{ d}^{-1}$ . The relative amplitudes of peaks between  $9$  and  $14 \text{ d}^{-1}$  are different, however,





**Figure 10.** Frequency spectra of VV Ori using PERIOD04 on the residuals from the 2018 and 2020 binary LC fittings to the *TESS* data sets.

while the cited frequencies differ by more than their formal uncertainty estimates. This may be attributed to the different data extraction methods employed, but slight changes in the pulsation properties in the two years between the *TESS* observation sets cannot be ruled out in view of the low level of the formal errors.

### 3 SPECTROMETRY

Spectroscopic data examined in this study include observations made with the HERCULES spectrograph (Hearnshaw et al. 2002), using the 1-m McLellan telescope at the University of Canterbury Mt John Observatory (UCMJO) ( $\sim 43^{\circ}59'S$ ,  $174^{\circ}27'E$ ) in New Zealand. Around 25 spectral images were obtained, distributed over the period 2010–15. These UCMJO observations provide somewhat incomplete coverage for the first half of the full cycle, but allow a consistent result. The data were collected with a  $4k \times 4k$  Spectral Instruments (SITE) camera (Skuljan 2004). Starlight from the telescope was passed to the spectrograph by a  $100 \mu\text{m}$  effective diameter fibre, corresponding to a theoretical resolution of  $\sim 40\,000$ . Wavelength and relative flux calibration was performed using the latest version of the software package HRSP (Skuljan, private communication) that outputs measurable data in FITS (Wells, Greisen & Harten 1981) formatted files. Fair weather exposures were usually for  $\sim 500$  s.

Some 45 clear orders (80–125) of the échelle were set up for inspection, using the software package VSPEC (Desnoux & Buil 2005).<sup>1</sup> Useful spectra typically have a signal to noise ratio (S/N) of  $\sim 100$  in order 85. Given the high-red sensitivity of the SITE camera, this drops to  $\sim 50$  by order 125.

Appendix Table A2<sup>2</sup> lists individual lines, mainly of the primary, detected in this study. Apart from  $H_{\alpha}$  and  $H_{\beta}$ , the best-defined line is probably the primary He I  $\lambda 6678$  line, where the corresponding weak secondary feature is also seen. A similar situation holds at He I  $\lambda 5875$ ; but the other He lines appear weaker and the secondary is not clearly distinguished at  $\lambda 5047$ . High excitation lines of C, N,

O, and Si are increasingly seen in the higher orders, though their S/N deteriorates and blending can be expected as the primary non-hydrogen lines are  $\sim 6\text{--}8 \text{ \AA}$  wide. The He II feature at  $\lambda 4686$ , together with the high-excitation lines, suggest the type may be a bit earlier than the often cited B1. The interstellar Na II lines reported by Terrell, Munari & Siviero (2007) are confirmed and separated into two main concentrations separated by about  $0.33 \text{ \AA}$  ( $\sim 17 \text{ km s}^{-1}$ ).

The UCMJO data on VV Ori were combined with a set of 45 high-resolution spectra taken in 2006 and 2007 with the FIES spectrograph at the NOT at the Roque de los Muchachos Observatory, La Palma, Canarias, Spain. This facility has been reviewed by Telting et al. (2014). Although some disparities were encountered in combining the FIES and HERCULES data, the sharp interstellar sodium lines are quite prominent and useful for checking wavelength calibration. The greatest deviations show up in the last two spectra obtained with HERCULES. These images from December 2015 are separated by over a year from the main cluster of observations (Table 6). The FIES spectra form a more compact data set, coming from two short observing runs in November 2006, and October 2007. In the context of possible secular changes in RV curves, it would be desirable to complete phase coverage over as short an interval as possible.

#### 3.1 Spectral disentangling: determination of the orbital parameters

Spectral lines in high-mass binary systems, like VV Ori, are usually broadened by high rotational velocities and often become blended over the course of the orbital cycle. Therefore, direct RV measurements are probably uncertain. In such cases, even more modern cross-correlation function (CCF) techniques become ineffective.

Spectral disentangling (Simon & Sturm 1994; Hadrava 1995) overcomes most of these problems, enabling a simultaneous determination of the orbital parameters, together with a reconstruction of the individual spectra of the components. There is no need for template spectra in this technique, which are usually the main source of uncertainty in RV evaluation by the CCF method. This can come from mismatches in spectral types (Hensberge & Pavlovski

<sup>1</sup>This is an MS-Windows™ based package that includes essential data-processing functions.

<sup>2</sup>Placed in the appendix on page 16 to avoid disrupting the flow of the paper.


**Table 6.** RV data of the components of VV Ori derived from the He I lines.

| BJD<br>2450000 + | Orbital<br>phase | RV1<br>km s <sup>-1</sup> | RV2<br>km s <sup>-1</sup> |
|------------------|------------------|---------------------------|---------------------------|
| 5539.9398        | 0.4272           | -26.83                    | -                         |
| 5539.9576        | 0.4391           | -24.28                    | -                         |
| 5544.8758        | 0.7502           | 164.45                    | -302.3                    |
| 5544.8917        | 0.7609           | 163.44                    | -302.3                    |
| 5544.9560        | 0.8040           | 146.60                    | -66.1                     |
| 5545.0213        | 0.8482           | 134.39                    | -192.6                    |
| 5549.9023        | 0.1342           | -97.22                    | -                         |
| 5798.2189        | 0.3083           | -127.32                   | 282.3                     |
| 5798.2600        | 0.3360           | -120.18                   | 276.7                     |
| 5798.2737        | 0.3452           | -120.69                   | 280.8                     |
| 5875.9380        | 0.6311           | 157.24                    | -190.7                    |
| 5875.9391        | 0.6318           | 133.34                    | -221.2                    |
| 5875.9529        | 0.6411           | 142.01                    | -226.8                    |
| 5875.9530        | 0.6411           | 157.22                    | -208.6                    |
| 5876.0580        | 0.7119           | 181.71                    | -58.2                     |
| 5876.0584        | 0.7121           | 153.74                    | -282.9                    |
| 5876.0720        | 0.7213           | 181.67                    | -260.5                    |
| 5876.0724        | 0.7216           | 159.35                    | -266.1                    |
| 5880.0972        | 0.4312           | -15.61                    | -                         |
| 6255.1256        | 0.9114           | 75.7                      | -                         |
| 6258.0204        | 0.8602           | 109.36                    | -231.9                    |
| 6258.0690        | 0.8930           | 101.71                    | -236.0                    |
| 6666.9080        | 0.1355           | -95.60                    | 282.5                     |
| 6673.9112        | 0.8502           | 106.81                    | -240.0                    |
| 6675.0142        | 0.5928           | 107.71                    | -187.0                    |
| 6993.9697        | 0.3231           | -194.65                   | 202.7                     |
| 7350.1629        | 0.1229           | -99.01                    | 261.9                     |
| 7356.1382        | 0.1456           | -109.98                   | 268.8                     |

2007). Precision is thus gained in the relative RV values, but the mean motion of the system  $V_\gamma$  has to be determined separately. With disentangling, the components' spectra are effectively separated, which, in turn, permits useful atmospheric diagnostics for either star. This allows a determination of their metallicity values, or detailed abundance signatures. For recent applications of spectral disentangling in complex high-mass binary systems see Pavlovski, Southworth & Tamajo (2018) and Pavlovski et al. (2023), where the methodology used in the present work is described in detail.

Since the FIES and HERCULES spectra do not cover all the same spectral range, we concentrated on the region where they overlap, that is, between the H  $\gamma$  and H  $\beta$  lines. This spectral segment covers about 400 Å and contains various metal lines, the most prominent of which is the He I 4471 Å line. As seen in Fig. 11 the helium and metal lines are resolved in the spectra of VV Ori obtained at quadrature. However, in spite of the large RV amplitude of the secondary component (about 660 km s<sup>-1</sup>), H  $\beta$  and the other hydrogen lines reflect this only in a changing asymmetry through the course of the orbital cycle. The two main components are never clearly resolved in the Balmer lines. Although spectral disentangling will resolve the hydrogen lines of the primary and secondary, because of the severe blending, precision in determination of the orbital parameters from the H lines is certainly smaller than for the resolved helium and metal lines.

The third body identified through speckle interferometry in the WDS catalogue is approximately 4 mag fainter in V than the primary. It is therefore probably an early-A-type main sequence star (Eker et al. 2018) whose observed spectra would show only relatively weak features apart from the Balmer lines. Admission of a third



**Figure 11.** Disentangled spectra of the primary component (in solid blue colour) and the secondary component (in solid red colour) superimposed on the observed spectrum of VV Ori in the quadrature (in solid grey). Whilst the He I 4920 Å line is clearly resolved, the broad H  $\beta$  line shows only an asymmetric profile, due to unresolved components.


contribution into the modelling at the level of a few per cent of the main component cannot be definitely confirmed from the residuals in the two-component fitting.

The spectral disentangling programme specifically referred to in this paper is FDBINARY (Ilijic et al. 2004). This is based on the formulation of disentangling in Fourier space, after Hadrava (1995). Methodological principles concerning disentangling in the wavelength domain are from Simon & Sturm (1994). In FDBINARY Fast Fourier Transforms (FFTs) are used that enable high flexibility in the selection of suitable spectral segments for disentangling whilst maintaining the original spectral resolution.

The RV curves shown in Fig. 12 are based on the line-centre wavelength values determined by the line-fitting process. The observed data points were optimally matched by the theoretical model curve (the continuous lines shown in Fig. 12) derived using the RV-curve application of WF. The corresponding parameters are listed in Table 7. The velocity semi-amplitudes ( $K_1$  and  $K_2$ ) are both significantly larger than those inferred from the masses given by Terrell, Munari & Siviero (2007). As a result, the masses we derive are also substantially larger than previously thought.

### 3.2 Atmospheric parameters

Once individual components' spectra have been separated they can be used for detailed spectroscopic analysis as single star spectra. The disentangled spectra still refer to the common continuum of the binary, hence they are diluted by the fractional light contribution of either component to the total light of the system. Generally, there are two options at this point: (i) disentangled spectra might be first renormalized to their own continua, using the light ratio determined by other means, that is, LC analysis (Pavlovski & Hensberge 2010), or (ii) the analysis could be performed directly on the disentangled spectra, since these spectra contain information on the light ratio (Tamajo, Pavlovski & Southworth 2011). It has been shown that uncertainties in the determination of the light ratio from disentangled spectra are comparable to the uncertainties achieved in photometric analysis, and are typically on the order of 1 per cent (Pavlovski et al.



**Figure 12.** The relative radial velocities for the components of VV Ori representing the orbital solution obtained with spectral disentangling (solid black lines). The observations obtained with the FIES spectrograph are shown as open blue circles, while open red ones represent the HERCULES data.

**Table 7.** The orbital parameters for VV Ori determined by spectral disentangling. The orbit is essentially circular, hence only the RV semi-amplitudes  $K_1$  and  $K_2$  of the components are determined (see Fig. 12). The derived values of the mass ratio,  $q$ , and masses of the components,  $M_1$  and  $M_2$  multiplied by  $\sin^3 i$ , are given.

| Parameter                        | Value      | Uncertainty |
|----------------------------------|------------|-------------|
| $K_1$ km s <sup>-1</sup>         | 137.20     | ± 0.59      |
| $K_2$ km s <sup>-1</sup>         | 328.58     | ± 1.56      |
| $P$ (d)                          | 1.48537742 | –           |
| $q$                              | 0.418      | ± 0.003     |
| $a \sin i$                       | 13.67      | ± 0.05      |
| $M_1 \sin^3 i M_\odot$           | 10.97      | ± 0.13      |
| $M_2 \sin^3 i M_\odot$           | 4.58       | ± 0.04      |
| $V_\gamma$ (km s <sup>-1</sup> ) | 26.7       | ± 2.0       |

2009; Pavlovski, Southworth & Tamajo 2018; Pavlovski et al. 2022, 2023). In the present work, we decided on the second option, in which the light ratio pertaining to the spectral segment studied would be determined simultaneously with the atmospheric parameters.

The optimal fitting of our disentangled spectra was performed by an extensive search through a pre-calculated grid of theoretical spectra. As a fitting merit indicator we used the sum of squared residuals between the disentangled spectrum and the selected synthetic spectrum using the STARFIT (Kolbas et al. 2014) code. The search procedure is performed by a genetic algorithm based on the PIKAIA subroutine of Charbonneau (1995). The uncertainties are calculated using a Markov Chain Monte Carlo (MCMC) procedure (Ivezic et al. 2014).

Since, in a close binary system, the sum of the fractional light contributions of both components should be unity, we would normally constrain the optimal fitting of the disentangled spectra with that condition. However, as discussed earlier, there are indications of a faint companion to the close pair in VV Ori. We therefore performed the optimal fitting in an unconstrained mode to allow for a light contribution of such a third star. The result of the modelling, with all

**Table 8.** The main atmospheric parameters for the components of VV Ori determined by optimal fitting of their disentangled spectra. Also, the fractional light contributions of the components (light dilution factor, LDF) can be optimized in an unconstrained mode (see subsection 3.2).


| Parameter                        | Run 1             | Run 2             |
|----------------------------------|-------------------|-------------------|
|                                  | Primary           |                   |
| $T_e$ [K]                        | $26\,010 \pm 320$ | $26\,660 \pm 300$ |
| $\log g$ [cm s <sup>-2</sup> ]   | $3.96 \pm 0.02$   | $4.083^*$         |
| $v \sin i$ [km s <sup>-1</sup> ] | $138.4 \pm 2.3$   | $151.4 \pm 2.5$   |
| Light dilution factor            | $0.908 \pm 0.007$ | $0.901 \pm 0.007$ |
|                                  | Secondary         |                   |
| $T_e$ [K]                        | $15\,780 \pm 420$ | $16\,250 \pm 410$ |
| $\log g$ [cm s <sup>-2</sup> ]   | $4.29 \pm 0.03$   | $4.320^*$         |
| $v \sin i$ [km s <sup>-1</sup> ] | $89.3 \pm 3.1$    | $98.1 \pm 3.5$    |
| Light dilution factor            | $0.096 \pm 0.008$ | $0.095 \pm 0.008$ |

atmospheric parameters and fractional light contributions left free, are given in Table 8. Model fits can be seen in Fig. 13.

The sum of the LDF, that is, the fractional light contributions of the components to the total light of the system, determined by our optimal fitting of disentangled components' spectra, then became slightly larger than unity, thus denying any significant light contribution from a third component. This finding does not support the results of the LC analysis in the present work. In the LC analysis of BVR photometry of VV Ori (Table 3), and the *TESS* observations (Table 5) a small third light appeared, in agreement with the astrometric findings of Horch et al. (2017). The light ratio of the main components we derived from the foregoing spectroscopic analysis, however,  $l_{r,sp} = l_2/l_1 = 0.106 \pm 0.009$  is in reasonably good agreement with Terrell et al.'s photometric light ratio in the B and b passbands, that is,  $l_{r,ph} = 0.093 \pm 0.001$  and  $0.098 \pm 0.001$ , respectively.

The eclipsing, double-lined spectroscopic binary nature of VV Ori's close pair allows determination of their masses and radii with high precision. Since one of the main obstacles in determination of the atmospheric parameters from hydrogen line profiles is degeneracy between the  $T_e$  and surface gravity, a determination of the latter from the system's dynamics could be used to lift this degeneracy. In the second run of the optimal fitting of the disentangled components' spectra, the surface gravity values for both components were fixed to those determined from combined spectroscopic and LC analyses. The results of Run 2 with  $\log g$ 's thus fixed, as indicated by an asterisk, are given in Table 8. Breaking the degeneracy in the  $T_e$  and  $\log g$  parameters has significant effects on the  $T_e$  values. These turn out to be about 650 and 250 K higher than the first run estimates for the primary and secondary components, respectively.

The primary's  $T_e$  determined by Terrell, Munari & Siviero (2007) was  $T_{e,1} = 26\,200$  K. This is in fair agreement with our determinations in both runs. This finding is encouraging, since Terrell, Munari & Siviero (2007) based the value of the primary's  $T_e$  on their dereddened B–V, using an estimate of the interstellar reddening from Na I lines. Older estimates of the  $T_e$  for the primary component (Table 1) were based on Eaton (1975), who found  $T_{e,1} = 25\,400 \pm 1\,500$  K from modelling the spectral energy distribution in the UV.



**Figure 13.** Optimal fitting of disentangled spectra for the primary component (lower) and secondary component (upper). Disentangled spectra are in red, optimal fits in black. The region of the spectra centred on the H  $\gamma$  line (left panel) and various metal lines in the spectral range from  $\lambda$  4500 to 4750  $\text{\AA}$  (right panel) are shown.

**Table 9.** Line-modelling parameters for VV Ori averaged from out-of-eclipse observations (see subsection 3.3 for explanation of the parameters).

| Parameter                     | Primary | $\pm$ | Secondary | $\pm$ |
|-------------------------------|---------|-------|-----------|-------|
| $U$                           | 0.954   | 0.001 | 0.955     | 0.001 |
| $I_0$                         | -0.101  | 0.004 | -0.018    | 0.002 |
| $v_{\text{rot}}$ km s $^{-1}$ | 147.1   | 5.7   | 86.1      | 10.3  |
| $s$ km s $^{-1}$              | 5.6     | 2.3   | 34.3      | 1.6   |
| $\Delta f$                    | 0.007   |       | 0.007     |       |
| $\chi^2/\nu$                  | 1.0     |       | 0.93      |       |


### 3.3 Rotational velocities

If the resolution is sufficiently high, spectral-line profiles can be modelled with a parameter set that determines the source’s rotation rate and scale of turbulence in the surrounding plasma, as well as the wavelength of the centre of light. Such modelling has been carried out in numerous previous studies (Shajin & Struve 1929; Huang & Struve 1954; Slettebak 1985; Butland et al. 2019). The profiles of the He I lines in our high-dispersion spectral images, particularly the  $\lambda 6678$  feature, are well suited to this purpose.

The entries in Table 9 follow a similar arrangement to the other tables of model fitting in this paper.  $U$  refers to the continuum reference level at the line’s central wavelength corresponding to the example shown in Fig. 14. The local continuum is assumed to be a straight line, but it may have a definite slope in the raw data that is empirically dealt with in the fitting procedure.

The quantities  $I_0$  correspond to the central depths relative to the local continuum (average of the six profiles examined).

The equatorial rotational velocity of the primary  $v_{\text{rot},1}$  is calculated with an inclination  $i = 79.0^\circ$  (Table 5) to correct for the projection. The parameter  $s$  relates to the scale of Gaussian broadening that is convolved with that from the rotation. It may be interpreted as a measure of the turbulence of the source plasma, or perhaps some other near-symmetric broadening effect. The conspicuous and consistently obtained high value of this parameter for the secondary (compared



**Figure 14.** Profile fitting to the He I  $\lambda 6678$  lines: The lines have been fitted separately and do not quite return to the same sloping continuum level.

to the primary) is noteworthy.  $\Delta f$  indicates the standard deviation of raw measures in the data sample.

For synchronized rotation with aligned axes we would expect the projected equatorial speed of the primary, using Table 10, to be  $\sim 0.368 \times 466/\sin i$  km s $^{-1}$ , that is,  $\sim 175$  km s $^{-1}$ , but from Table 9 it is seen that the observed speed of 147 km s $^{-1}$  is significantly less than that. The ratio of the two rotational speed estimates ( $\sim 0.59$ ) is also significantly different to that of the ratio of radii following from Table 5 (0.492). In fact, if we divided the secondary’s rotational

**Table 10.** Rotational velocities: observed and calculated.

|                   | Primary         | Secondary      |
|-------------------|-----------------|----------------|
| $v \sin i$        | $138.4 \pm 2.3$ | $89.3 \pm 3.1$ |
| $v_{\text{sync}}$ | $176.8 \pm 0.8$ | $86.4 \pm 1.7$ |

**Table 11.** Photospheric elemental abundances for the primary component in VV Ori. For comparison, abundances determined for B-type stars in the Ori I association in Nieva & Simón-Díaz (2011) (abbreviated NSD) are also presented. The elemental abundances for species X is given relative to the hydrogen abundance,  $\epsilon(X) = \log(X/H) + 12$ . The N/O and N/C abundance ratios are also given, as a sensitive probe to mixing processes in the stellar interiors.

| Element | This work<br>[dex] | NSD<br>[dex]     |
|---------|--------------------|------------------|
| C       | $8.28 \pm 0.06$    | $8.35 \pm 0.03$  |
| N       | $7.77 \pm 0.07$    | $7.82 \pm 0.07$  |
| O       | $8.76 \pm 0.07$    | $8.77 \pm 0.03$  |
| Mg      | $7.64 \pm 0.06$    | $7.57 \pm 0.06$  |
| Si      | $7.51 \pm 0.08$    | $7.50 \pm 0.06$  |
| Al      | $6.41 \pm 0.08$    | -                |
| N/O     | $-0.99 \pm 0.10$   | $-0.95 \pm 0.08$ |
| N/C     | $-0.51 \pm 0.09$   | $-0.53 \pm 0.08$ |

velocity by that ratio we would obtain the synchronous rotation value  $175 \text{ km s}^{-1}$  for the primary, that is, the mean rotational speed of the secondary is close to synchronism, unlike the primary.

Following the discussion of Southworth, Bowman & Pavlovski (2021), this can be interpreted as a non-alignment of the primary's rotation and orbit axes, that would correspond to an obliquity angle ( $\epsilon$ ) at zero precession angle ( $\psi$ ) of around  $33^\circ$ . Precession of this spin axis could cause the apparent variation of inclination reported by Southworth, Bowman & Pavlovski (2021).

### 3.4 Elemental abundances for the primary component

As the primary component contributes most of the light from this binary, it has more weight in the spectral disentangling. The S/N of the primary's disentangled spectrum is thus relatively high. This is an important point for detailed abundance analysis. Our adopted procedure is fully described in Pavlovski, Southworth & Tamajo (2018) and Pavlovski et al. (2023). A grid of non-local thermodynamic equilibrium (NLTE) synthetic spectra is calculated using the programmes DETAIL and SURFACE with model atmospheres first produced by ATLAS9 using the local thermodynamic equilibrium (LTE) prescription. Theoretical spectra are broadened according to a given instrumental profile and projected rotational velocity. Abundances are then determined from line-profile fittings of the normalized disentangled spectrum of the primary to the adjustable theoretical one.

The following species were considered: C, N, O, Mg, Si, and Al. The microturbulent velocity,  $\xi = 2 \pm 1 \text{ km s}^{-1}$ , is determined from the oxygen lines, which appear to be the most numerous in the primary's spectrum. The result of the abundance analysis is given in Table 11. Determined abundances are in excellent agreement with previous analyses of high-mass stars in detached binary systems (see Pavlovski, Southworth & Tamajo 2018; Pavlovski et al. 2023). There has been no previous abundance analysis specifically for VV Ori, but stars in the Ori I association have been extensively studied through recent decades (Simón-Díaz 2010; NSD 2011, and

**Table 12.** Adopted absolute parameters of VV Ori, derived from the combined photometric and spectroscopic analyses discussed in Sections 2 and 3. The units labelled with an 'N' are given in terms of the nominal solar quantities defined in IAU 2015 Resolution B3 (Prša et al. 2016).

| Parameters  |                   | Value | Uncertainty |
|---|-------------------|-------|-------------|
| Masses ( $\mathcal{M}_{\odot}^{\text{N}}$ )         | $M_1$             | 11.56 | 0.14        |
|   | $M_2$             | 4.81  | 0.06        |
|   | $M_3$             | 2.0   | 0.3         |
| Radii ( $\mathcal{R}_{\odot}^{\text{N}}$ )          | $R_1$             | 5.11  | 0.03        |
|   | $R_2$             | 2.51  | 0.02        |
|   | $R_3$             | 1.8   | 0.1         |
| Semimajor axis ( $\mathcal{R}_{\odot}^{\text{N}}$ ) | $a$               | 13.91 | 0.05        |
| Temperatures (K)                                    | $T_{e1}$          | 26660 | 300         |
|   | $T_{e2}$          | 16250 | 420         |
|   | $T_{e3}$          | 10000 | 1000        |
| Luminosities $\log(L/L_{\odot}^{\text{N}})$         | $\log L_1$        | 4.07  | 0.02        |
|   | $\log L_2$        | 2.60  | 0.06        |
|   | $\log L_3$        | 1.5   | 0.2         |
| Absolute magnitudes                                 | $M_{\text{bol}1}$ | -5.44 | 0.05        |
|   | $M_{\text{bol}2}$ | -1.75 | 0.14        |
|   | $M_{\text{bol}3}$ | 1.1   | 0.40        |
| Reddening (mags)                                    | $E(B - V)$        | 0.08  | 0.03        |
| Gravities ( $\log [cgs]$ )                          | $\log g_1$        | 4.08  | 0.05        |
|   | $\log g_2$        | 4.32  | 0.06        |
|   | $\log g_3$        | 4.25  | 0.10        |
| Distance (pc)                                       | $\rho$            | 396   | 7           |

references therein). Clearly, the elemental abundances, and nitrogen-to-oxygen (N/O) and nitrogen-to-carbon (N/C) abundance ratios for the primary component in VV Ori is the same as for stars in the Ori I association, within the given uncertainties.

## 4 ABSOLUTE PARAMETERS

It is well known that the actual sizes of the component stars in an eclipsing binary system (radii  $R_1$ ,  $R_2$ ) can be determined by combining the results of LC and RV curve parametrization – the 'eclipse' or 'Russell's' method. The absolute size of the orbit comes from dividing the RV parameter  $a \sin i$  by  $\sin i$ , using the inclination  $i$  from the LC modelling. The sought radii are simply the product of  $a$  and the fractional radii  $r_{1,2}$ . Surface gravities ( $g_{1,2}$ ), are scaled from the solar value ( $\log g_{\odot} = 4.438$ ) using the radii and masses  $M_{1,2}$  similarly derived from the combined RV and LC parameters, or using Kepler's third law, given the period  $P$  and separation  $a$  of the two stars.

We thus determined the physical properties of VV Ori from the photometric results of Section 2 and the spectrometry of Section 3. For this we have used the JK TABSDIM code (Southworth, Maxted & Smalley 2005), modified to apply the IAU system of nominal solar values (Prša et al. 2016), together with the NIST 2018 values for the Newtonian gravitational and the Stefan–Boltzmann constants. Error bars were derived from the perturbation analysis referred to in subsection 2.1. The results are given in Table 12.

We calculated a distance to the system using optical  $UBVR$  magnitudes from Ducati et al. (2001), near-IR  $JHK$ , magnitudes from 2MASS (Cutri et al. 2003) converted to the Johnson system using the transformations from Carpenter (2001), and bolometric corrections from Girardi et al. (2002). The interstellar reddening was determined by requiring the optical and near-IR distances to match. We found a distance of  $396 \pm 7 \text{ pc}$ . This is significantly shorter than the distance of  $441 \pm 22 \text{ pc}$  from the *Gaia* DR3 parallax (Gaia Collaboration et al. 2016, 2018, 2021, 2023). Possible explanations for this are that the

*Gaia* parallax was affected by the brightness of the system and/or the presence of the nearby third body. Evidence in support of this comes from the renormalized unit weight error (RUWE) of 1.347 being close to the upper limit of 1.4, beyond which the *Gaia* parallax is considered unreliable.<sup>3</sup>

The photometric parallax ( $\pi$ ) can be derived from the formula (Budding & Demircan, 2007, equation 3.42)

$$\log \pi = 7.454 - \log R - 0.2V - 2F'_V, \quad (7)$$


where  $F'_V$  is directly proportional to the logarithm of a star's mean surface flux (Barnes & Evans 1976), and is specified by  $F'_V = \log T_e + 0.1BC$ , where  $BC$  is the bolometric correction. Applying equation (7) directly to the stars in VV Ori, with the V magnitudes from Section 2, the  $R$  and  $T_e$  values from Table 12, and the  $BC$  values from Budding & Demircan (2007) Table 3.1, we obtain  $\log \pi_1 = -2.64$ , and  $\log \pi_2 = -2.66$ . The third star would also produce a comparable value  $\log \pi_3 \approx -2.7$ , if we put its  $T_e$  at 10 000 K, but data on that star are still very approximate. These parallaxes are in close agreement with the value cited above from the *Gaia* DR3, but the measured  $B - V$  colour excess argues against adopting the measured V as unaffected by interstellar absorption. Using the relation of Cardelli, Clayton & Mathis (1989) with the reddening  $E = 0.08$ , that is,  $A_V = 0.26$ , the mean distance turns out to be  $\rho = 396 \pm 7$  pc, in good agreement with the foregoing estimate from the  $V - I$  colour.

Evaluation of the absolute luminosities ( $L_{1,2}$ ) and bolometric magnitudes ( $M_{\text{bol},2}$ ) of the component stars requires the  $T_e$  values to be known. These were enumerated from the spectral disentangling results given in Table 8, and checked with the colours derived from Table 2. In these calculations, we adopted the solar calibration values as: effective temperature  $T_e = 5780$  K,  $M_{\text{bol}} = 4.75$  from the IAU-adopted solar constants. The adopted absolute parameters for VV Ori are listed, with their uncertainties, in Table 12.

The parameters of the third star are included in Table 12 for completeness, assuming that it is coeval with the close binary and about 4.5 V mag fainter (Section 1). The third star's properties should still be regarded as quite imprecise compared with those of the main components. Better knowledge of this wide system (VV Ori AB) can be expected in future high-accuracy survey work.

## 5 DISCUSSION

Part of the rationale for this study was the ongoing programme of precise quantification of stellar properties, using modern data and analysis techniques. We have shown that independent use of two different LC analysis procedures on recent data sets from the *TESS* programme resulted in very similar values for the main geometric parameters. Combining these parameters with the RV analysis on high-dispersion FIES and HERCULES spectrograms, then recovers absolute parameter sets that are within reasonable error estimates of each other. This bolsters confidence in the employed analytical methods, and provides reliable evidence to check against theory. Fig. 15 shows such a comparison. The MESA Isochrones and Stellar Tracks MIST facility (Dotter 2016; Choi et al. 2016) based on the MESA models (Paxton et al. 2011) was used to plot evolutionary tracks and isochrones for comparison with our derived



**Figure 15.** Derived  $\log T_e$  and  $\log g$  parameters (Table 12) with their uncertainties for the close pair in VV Ori (filled black circles) are compared with the MIST evolutionary tracks and isochrones (Dotter 2016; Choi et al. 2016). The near-vertical evolutionary tracks are labelled according to masses in  $M_\odot$ . Blue curves are close to the adopted values. Isochrones (dotted) are given in logarithm of time in year.

parameters. The masses and luminosities are in close agreement with corresponding models at a log age (in yr) of  $\sim 6.9$ .

Our results confirm that VV Ori is a young binary system and shares properties with other members of the Orion Ib OB star association, with an age between about 6 and 10 Myr. We confirm a photometric distance of around 400 pc: closer to the value ( $\sim 360$  pc) of Brown, de Geus & de Zeeuw (1994) for the Ib subgroup, but lower than the distance ( $\sim 500$  pc) of Warren & Hesser (1978). The *Gaia* DR3 value ( $\sim 440$  pc) was considered relatively imprecise, perhaps due to calibration difficulties for this bright star in a crowded field containing nebulosity. Post-*Gaia* population studies (Zari, Brown & de Zeeuw 2019) have identified substructures within the Orion Ib Association (see also Warren & Hesser, 1977, 1978). Interestingly, the galactic coordinates of VV Ori would place it in the B7, or, marginally, the E subgroup, which has an estimated mean age of close to 11 Myr.

In Section 4 we have given the log age value as  $\approx 6.9$ , supporting the idea that VV Ori is slightly older than the bulk of the  $\epsilon$  Ori association (see Fig. 1). The age of 8 Myr is, however, young enough to fall within Zahn's (1977) synchronization time-scale for massive stars with radiative envelopes and initial separations of around  $20 R_\odot$ .

In this context, it may be possible to form an independent assessment of the age by referring to the estimated synchronization time-scales. The observed  $v \sin i$ , and synchronized velocities are given in Table 10. The observed rotational velocity of the primary appears low by  $\sim 40 \text{ km s}^{-1}$ . The secondary's rotational velocity is, alternatively, in good agreement with a synchronized state. The low width of the primary's line core implies that any additional broadening effect, such as macro-turbulence, would not improve agreement between the observed primary rotation and the value corresponding to synchronization.

This low value of the apparent rotation is, however, in keeping with the possibility of a displaced spin axis raised in subsection 2.2.

<sup>3</sup>[https://gea.esac.esa.int/archive/documentation/GDR2/Gaia\\_archive/chap\\_datamodel/sec\\_dm\\_main\\_tables/ssec\\_dm\\_ruwe.html](https://gea.esac.esa.int/archive/documentation/GDR2/Gaia_archive/chap_datamodel/sec_dm_main_tables/ssec_dm_ruwe.html)

Given the report of changes in the apparent inclination (Southworth, Bowman & Pavolvski 2021) and keeping in mind the interaction between the heat-transfer driven  $\beta$  Cep pulsations and tides (Townsend, Goldstein & Zweibel 2018; Pedersen 2022), the phenomenon of precession can resolve the various observed oddities of the system. The dynamics of this situation then invites critical attention from relevant theory.

Here, we may note that a major source of uncertainty in modelling stellar structure and pulsation properties of upper main sequence stars is the distribution and transport of internal angular momentum. Asteroseismology allows for the determination of the interior rotation of stars, provided non-radial oscillations are present at a suitable level. Normal modes of oscillation, calculated for non-rotating, single stars, become split by rotation and tidal distortions.

Four main processes have been taken to contribute to angular momentum transport within stellar radiative regions: meridional circulation, turbulence driven by instabilities, magnetism, and internal waves (Goldreich & Nicholson 1989; Zahn 2013; Mathis 2013). During core hydrogen and helium-burning phases, single stars are expected to rotate nearly uniformly. However, in a binary system, even during early phases of evolution, tidal interactions may give rise to non-uniform rotation. In particular, differential rotation in radiative envelopes can induce a diversity of hydrodynamical and magnetohydrodynamical instabilities, that will, in turn, transport and redistribute angular momentum. The secondary component of VV Ori appears to be in near-synchronous rotation, so we expect differential rotation in that star to be relatively unimportant in comparison with that of the asynchronous primary.

Analysis of the low-order oscillation modes observed in  $\beta$  Cep stars should be able to inform us about the star's internal structure. Thus, convective overshoot at the convective core/radiative interface is a major source of turbulence. Mixing length theory contains a free parameter which determines the extent of overshoot in terms of the local pressure scale height  $H_p$ . This overshoot region is probed by comparing observed oscillation frequencies with those of models for a range of prescriptions of the overshoot parameter. So, as well as determining the size and location of the convective core, the frequency spectrum reflects conditions around it that bear on the stellar modelling and angular momentum regime.

A potentially important source of angular momentum transport is internal gravity waves (IGWs) propagating in the stably stratified envelope. The generation and damping of these waves depend sensitively on their frequency and length-scales. They may be present either at the core-radiative interface, or in thin convection zones near the surface, where local opacity increases in accordance with the  $\kappa$ -mechanism, or else by tidal forcing (Zahn 1975). For the latter, tidal forces generate a disturbance at the interface that propagates away from the core. Such waves are large-scale  $l = 2$ ,  $m = 1$  or  $2$  and have frequency equal to the forcing frequency  $\sigma$  (in the corotating frame). In a rotating star,  $\sigma = k\Omega_{\text{orb}} - m\Omega$ , where  $\Omega_{\text{orb}}$  is the orbital frequency and  $\Omega$  that of the rotation; while  $k > 0$  and  $m$  ( $|m| \leq l$ ) are integers. Dissipation of these waves occurs by non-adiabatic thermal damping effects near the surface of the star. This is different for prograde ( $m > 0$ ) and retrograde ( $m < 0$ ) waves, thus giving rise to a net transport of angular momentum. In this way, IGWs couple convective core and radiative envelope regions, and from asteroseismic data it appears that this coupling is strong.

However, the exact quantification of such conditions from seismic analysis appears still unsettled regarding modelling or observational data selections, with implications on the confidence of interpretation (Salmon et al. 2022). In particular, models that include rotation and/or tidal distortions are not yet directly available.

One source of uncertainty is the extent of chemical mixing which, in young intermediate to high-mass stars arises from instabilities in the radiative layers and, based on interpretation of asteroseismic data, is orders of magnitude lower than expected. Chemical gradients at the core-envelope interface should develop as the star evolves and affect the amplitudes of low-frequency  $g$ -mode waves.

With regard to magnetic fields transporting angular momentum, these may be thought to be largely confined to the convective core (MacGregor & Cassinelli 2003), though other convective zones near the surface would generate fields that could transmit angular momentum along connecting field lines.

Townsend, Goldstein & Zweibel (2018) have investigated the transport of angular momentum by heat-driven non-radial  $g$ -modes, producing a table of models covering most of the mass range of young B-type stars. From their Fig. 7, it can be seen that both VV Ori's primary and secondary (Table 12) are on the edge of the region in the HR diagram, in which their torque instability operates. While this may suggest that the proposed mechanism is not viable, the model does not include the effects of tides or rotation. Moreover, Townsend, Goldstein & Zweibel assumed aligned spin and orbital axes, whereas with the putative precession in VV Ori this would not be the case. One of the effects of non-alignment would be that the  $m = \pm 1$  modes would also contribute to angular momentum transport in the radiative region, in addition to the  $m = \pm 2$  modes (for the dominant  $l = 2$  tide).

Clearly, the effects of tides and rotation on the instabilities that give rise to these waves need to be investigated before any definitive predictions can be made. Meanwhile, the juxtaposition of the properties of the stars in VV Ori, as determined from the classical methods reported in this paper, against asteroseismological inferences should have very interesting consequences.

## ACKNOWLEDGEMENTS

Generous allocations of time on the 1-m McLennan Telescope and HERCULES spectrograph at the Mt John University Observatory in support of the Southern Binaries Programme have been made available through its TAC and supported by its Director, Dr. K. Pollard and previous Director, Prof. J. B. Hearnshaw. Useful help at the telescope was provided by the MJUO management (N. Frost and previously A. Gilmore & P. Kilmartin). Considerable assistance with the use and development of the HRSP software was given by its author Dr. J. Skuljan, and very helpful work with initial data reduction was carried out by R. J. Butland.

This work was based on observations made with the NOT, owned in collaboration by the University of Turku and Aarhus University, and operated jointly by Aarhus University, the University of Turku and the University of Oslo, representing Denmark, Finland and Norway, the University of Iceland and Stockholm University at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias.

General support for this programme has been shown by the School of Chemical and Physical Sciences of the Victoria University of Wellington; as well as the Çanakkale Onsekiz Mart University, Turkey, notably Prof. O. Demircan. The Royal Astronomical Society of New Zealand, particularly its Variable Stars South section (<http://www.variablestarssouth.org>), was also supportive.

It is a pleasure to express our appreciation of the high-quality and ready availability, via the Mikulski Archive for Space Telescopes (MAST), of data collected by the *TESS* mission. Funding for the *TESS* mission is provided by the NASA Explorer Programme. This research has made use of the SIMBAD data base, operated at CDS, Strasbourg,

France, and of NASA's Astrophysics Data System Bibliographic Services. We thank the University of Queensland for the use of collaboration software. We are grateful for the helpful comments and guidance by the anonymous referee, which led to an improved paper.

## DATA AVAILABILITY

The *TESS* photometric data used in this study are publicly available from the Barbara A. Mikulski Archive for Space Telescopes (MAST) portal maintained by the Space Telescope Science Institute. The BVR photometric data underlying this article will be shared on reasonable request to M. Blackford and also available as supplementary material to the electronic version of the paper. HERCULES spectroscopy can be sourced from E. Budding. FIES spectra can be obtained from K. Pavlovski, J. Southworth, or the NOT archive at <https://www.not.iac.es/archive/>.

## REFERENCES

- Aerts C., Christensen-Dalsgaard J., Kurtz D. W., 2010, *Asteroseismology*. Springer, Dordrecht ISBN: 978-1-4020-5178-4, <https://doi.org/10.1007/978-1-4020-5803-5>
- Barnes T. G., Evans D. S., 1976, *MNRAS*, 174, 489
- Barr J. M., 1905, *Selected Papers and Proc. Roy. Astron. Soc. Canada*, 1904, 42
- Beltrami G., Galeotti P., 1969, *IAU Inform. Bull. Var. Stars*, 393, 1
- Blaauw A., 1964, *ARA&A*, 2, 213
- Bowman D. M., 2020, *Front. Astron. Space Sci.*, 7, 70
- Bowman D. M., Johnson C., Tkachenko A., Mkrtichian D. E., Gunsriwati K., Aerts C., 2019, *ApJ*, 883, L26
- Brown A. G. A., de Geus E. J., de Zeeuw P. T., 1994, *A&A*, 289, 101
- Budding E., Demircan O., 2007, *Introduction to Astronomical Photometry*. Cambridge Univ. Press, Cambridge ISBN: 9780511536175
- Budding E., Demircan O., 2022, *A Guide to Close Binary Systems*, Taylor & Francis Limited, Boca Raton ISBN: 9781138064386
- Budding E., Najim N. N., 1980, *Astrophys. Space Sci.*, 72, 369
- Budding E., Love T., Blackford M. G., Banks T., Rhodes M. D., 2021, *MNRAS*, 502, 6032
- Bursens S., et al., 2020, *A&A*, 639, 81
- Butland R. J., Erdem A., Özkardeş B., Blackford M. G., Sürgit D., Budding E., Demircan O., 2019, *MNRAS*, 482, 2644
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, L245
- Carpenter J. M., 2001, *AJ*, 121, 2851
- Chambliss C. R., 1983, *Astrophys. Space Sci.*, 89, 15
- Chambliss C. R., 1984, *Astrophys. Space Sci.*, 99, 163
- Charbonneau P., 1995, *ApJS*, 101, 309
- Choi J., Dotter A., Conroy C., Cantiello M., Paxton B., Johnson B. D., 2016, *ApJ*, 823, L102
- Clausen J. V., 1996, *A&A*, 308, 151
- Cutri R. M., et al., 2003, *Explanatory Supplement to the 2MASS All Sky Data Release*. NASA, Washington <http://www.ipac.caltech.edu/2mass/releases/allsky/doc/explsup.html>
- Daniel Z., 1915, *Publ. Allegheny*, 3, 179
- Desnoux V., Buil C., 2005, *Soc. Astron. Sci. Ann. Symp.*, 24, 129
- Dotter A., 2016, *ApJS*, 222, 8
- Ducati J. R., 2002, *CDS/ADC Collection of Electronic Catalogues*, 2237.
- Ducati J. R., Bevilacqua C. M., Rembold S. B., Ribeiro D., 2001, *ApJ*, 558, L309
- Duerbeck H. W., 1975, *A&AS*, 22, 19
- Eaton J. A., 1975, *ApJ*, 197, L379
- Eker Z., et al., 2018, *MNRAS*, 479, 5491
- Erdem A., et al., 2022, *MNRAS*, 515, 6151
- Feiden G. A., 2015, in Rucinski S. M., Torres G., Zejda M., eds, *Living Together: Planets, Host Stars and Binaries*, Proceedings of a conference held 8-12 September 2014 in Litomyšl, Czech Republic; ASP Conference Series, Vol. 496. Astronomical Society of the Pacific, San Francisco, p. 137
- Fuller J., Kurtz D. W., Handler G., Rappaport S., 2020, *MNRAS*, 498, 5730
- Gaia Collaboration, Brown A. G. A., Vallenari A., et al., 2016, *A&A*, 595, 1
- Gaia Collaboration, Brown A. G. A., Vallenari A., Prusti T., et al., 2018, *A&A*, 616, 22
- Gaia Collaboration, Brown A. G. A., Vallenari A., Prusti T., et al., 2021, *A&A*, 694, 1
- Gaia Collaboration, Vallenari A., Brown A. G. A., Prusti T., et al., 2023, *A&A*, 674, 1
- Girardi L., Bertelli G., Bressan A., Chiosi C., Groenewegen M. A. T., Marigo P., Salasnich B., Weiss A., 2002, *A&A*, 391, 195
- Golay M., 1974, *Introduction to Astronomical Photometry*. D. Reidel Publishing Company, Dordrecht ISBN: 978-90-277-0428-3
- Goldreich P., Nicholson P. D., 1989, *ApJ*, 342, L1079
- Hadrava P., 1995, *A&AS*, 498, 5730
- Hambleton K. M., et al., 2013, *MNRAS*, 434, 925
- Handler G., Kurtz D. W., Rappaport S. A., et al., 2020, *Nat Astron*, 4, 684
- Hearnshaw J. B., Barnes S. I., Kershaw G. M., Frost N., Graham G., Ritchie R., Nankivell G. R., 2002, *Exp. Astron.*, 13, 59
- Hensberge H., Pavlovski K., 2007, in Hartkopf W. I., Harmanec P., Guinan E. F., eds, *Binary Stars as Critical Tools & Tests in Contemporary Astrophysics*, Proc. IAU Symp. 240. Cambridge University Press, Cambridge, UK. p. 136
- Horch E. P., et al., 2017, *AJ*, 153, 212
- Huang S. S., Struve O., 1954, *Ann. d'ap.*, 17, 85
- Ilijic S., Hensberge H., Pavlovski K., Freyhammer L. M., 2004, in Hilditch R. W., Hensberge H., Pavlovski K., eds, *Spectroscopically and Spatially Resolving the Components of the Close Binary Stars*, Astronomical Society of the Pacific Conference Series Vol. 318. Astronomical Society of the Pacific, San Francisco. p. 111
- Ivezić Ž., Connolly A. J., VanderPlas J. T., Gray A., 2014, *Statistics, Data Mining, and Machine Learning in Astronomy: A Practical Python Guide for the Analysis of Survey Data*. Princeton Univ. Press, Princeton
- Kolbas V., Dervişoğlu A., Pavlovski K., Southworth J., 2014, *MNRAS*, 444, 3118
- Kopal Z., 1959, *Close Binary Systems*. Chapman & Hall, London
- Kurtz D. W., et al., 2020, *MNRAS*, 494, 5118
- Lampens P., 2006, in Sterken C., Aerts C., eds, *Astrophysics of Variable Stars*, Pecs, Hungary, 5-10 September 2005, ASP Conf. Ser. Vol. 349. Astronomical Society of the Pacific, San Francisco, p. 153
- Langer D., 2012, *ARA&A*, 50, 107
- Len P., Breger M., 2004, *Proc. IAU 2004, AUS224*, 786
- Lesh J. R., Aizenman M. L., 1978, *ARA&A*, 16, 215
- MacGregor K. B., Cassinelli J. P., 2003, *ApJ*, 586, L480
- Mathis S., 2013, in Goupil M., Belkacem K., Neiner C., Lignières F., Green J. J., eds, *Lecture Notes in Physics*, Vol. 865. Springer-Verlag, Berlin
- Murphy S., 2018, <https://doi.org/10.48550/arXiv.1811.12659>
- Nieva M. F., Simón-Díaz S., 2011, *A&A*, 532, 2
- Pavlovski K., Hensberge H., 2010, in Prša A., Zejda M., eds, *Binaries - Key to Comprehension of the Universe*, Astronomical Society of the Pacific Conference Series, Vol. 435, Astronomical Society of the Pacific, San Francisco. p. 207
- Pavlovski K., Tamajo E., Koubsky P., Southworth J., Yang S., Kolbas V., 2009, *MNRAS*, 400, 791
- Pavlovski K., Southworth J., Tamajo E., 2018, *MNRAS*, 481, 3129
- Pavlovski K., et al., 2022, *A&A*, 658, 92
- Pavlovski K., Southworth J., Tkachenko A., Van Reeth T., Tamajo E., 2023, *A&A*, 671, 139
- Paxton B., Bildsten L., Dotter A., Herwig F., Lesaffre P., Timmes F., 2011, *ApJS*, 192, 3
- Pedersen M. G., et al., 2019, *ApJ*, 872, L9
- Pedersen M. G., 2022, *ApJ*, 940, L49
- Popper D. M., 1993, *PASP*, 105, 721
- Prša A., et al., 2016, *AJ*, 152, 41
- Ratajczak M., Pigulski A., Pavlovski K., 2017, in Zwintz K., Poretti E., eds, *Proc. Polish Astronomical Society, Second BRITe-Constellation*



- Science Conf.: Small Satellites–Small satellites – Big Science. Polish Astronomical Society, Warsaw, p. 128
- Rhodes M. D., 2021, winfitter manual. <https://michaelrhodesbyu.weebly.com>
- Ricker G. R., et al., 2014, Proc. SPIE Vol. 9143. e-ID: 914320.
- Russell H. M., 1912a, *ApJ*, 35, L315
- Russell H. M., 1912b, *ApJ*, 36, L54
- Russell H. M., 1942, *ApJ*, 95, L345
- Russell H. M., Shapley H., 1912, *ApJ*, 36, L239
- Salmon S. J. A. J., Eggenberger P., Montalban J., Miglio A., Noels A., Buldgen G., Moyanol F., Meyneti G., 2022, *A&A*, 659, 142
- Sana H., et al., 2012, *Science*, 337, 444
- Sarma M. B. K., Vivekananda R. P., 1995, *J. Astrophys. Astron.*, 16, 407
- Shajin G., Struve O., 1929, *MNRAS*, 89, 222
- Simon K. P., Sturm E., 1994, *A&A*, 281, 286
- Simón-Díaz S., 2010, *A&A*, 510, 22
- Skuljan J., 2004, in Kurtz D. W., Pollard K. R., eds, Variable Stars in the Local Group, IAU Colloquium 193, Proceedings of the conference held 6–11 July, 2003 at Christchurch, New Zealand, ASP Conference Proceedings. Astronomical Society of the Pacific, San Francisco, p. 575
- Slettebak A., 1985, *ApJS*, 59, 769
- Southworth J., 2020, *The Observatory*, 140, 247
- Southworth J., 2021, *Universe*, 7, 369
- Southworth J., Bowman D. M., 2022, *MNRAS*, 513, 3191.
- Southworth J., Maxted P. F. L., Smalley B., 2005, *A&A*, 429, 645
- Southworth J., et al., 2011, *MNRAS*, 414, 2413
- Southworth J., Bowman D. M., Tkachenko A., Pavlovski K., 2020, *MNRAS*, 497, 19
- Southworth J., Bowman D. M., Pavolvski K., 2021, *MNRAS*, 501, 65
- Stankov A., Handler G., 2005, *ApJS*, 158, 193
- Sterne T. E., 1939, *MNRAS*, 99, 451
- Struve O., Luyten W. J., 1949, *ApJ*, 110, L160
- Tamajo E., Pavlovski K., Southworth J., 2011, *A&A*, 526, 76
- Teltin J. H., et al., 2014, *Astron. Nach.*, 335, 41
- Terrell D., Munari U., Siviero A., 2007, *MNRAS*, 374, 530
- Townsend R. H. D., Goldstein J., Zweibel E. G., 2018, *MNRAS*, 475, 879
- Van Hamme W., Wilson R. E., 2007, *ApJ*, 661, L1129
- Waelkens C., 1991, *A&A*, 246, 453
- Warren W. H., Jr, Hesser J. E., 1977, *ApJS*, 34, 115
- Warren W. H., Jr, Hesser J. E., 1978, *ApJS*, 36, 497
- Wells D. C., Greisen E. W., Harten R. H., 1991, *A&A Supp.*, 44, 363
- Welsh W. F., Orosz J. A., Aerts C., Brown T. M., et al., 2011, *ApJS*, 197, 4
- Wilson R. E., 1979, *ApJ*, 234, L1054
- Wilson R. E., Devinney E. J., 1972, *ApJ*, 166, L605
- Wilson R. E., Van Hamme W., 2004, Computing Binary Star Observables (Wilson–Devinney program user guide). Available at <ftp://ftp.astro.ufl.edu/pub/wilson>
- Wood F. B., 1946, Contr. Princeton Univ. Obs., 21, 25
- Wright N. J., 2020, *NewAR*, 90, 101549
- Zahn J.-P., 1975, *A&A*, 41, 329
- Zahn J.-P., 1977, *A&A*, 57, 383
- Zahn J.-P., 2013, in Alecian G., Lebreton Y., Richard O., Vauclair G., eds, EAS Publications Series, EDP Sciences, 91944 Les Ulis Cedex A, France, vol. 63
- Zari E., Brown A. G. A., de Zeeuw P. T., 2019, *A&A*, 628, 123
- Zucker C., Alves J., Goodman A., et al., 2023, in Inutsuka S.-i., Aikawa Y., Muto T., Tomida K., Tamura M., eds, Protostars and Planets VII, ASP Conference Series, Vol. 534, Proceedings of a conference held 10–15 April 2023 at Kyoto, Japan. Astronomical Society of the Pacific, San Francisco, p. 43
- İnlek G., Budding E., Demircan O., 2017, *ApSS*, 362, 167

## SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article

## APPENDIX A:

**Table A1.** Significant peaks in the frequency spectrum of VV Ori.

| Frequency $\text{d}^{-1}$ | Amplitude (mmag) | Phase (rad)     |
|---------------------------|------------------|-----------------|
| 0.0042 ± 0.0000           | 11.4053 ± 0.0047 | 0.7062 ± 0.0001 |
| 0.1166 ± 0.0001           | 1.1453 ± 0.0047  | 0.7511 ± 0.0007 |
| 0.2353 ± 0.0010           | 0.1014 ± 0.0047  | 0.7168 ± 0.0074 |
| 0.3087 ± 0.0012           | 0.0816 ± 0.0047  | 0.6794 ± 0.0092 |
| 0.4459 ± 0.0015           | 0.0674 ± 0.0047  | 0.0007 ± 0.0112 |
| 0.5902 ± 0.0012           | 0.0862 ± 0.0047  | 0.3041 ± 0.0087 |
| 0.6696 ± 0.0006           | 0.1589 ± 0.0047  | 0.6249 ± 0.0047 |
| 0.7776 ± 0.0013           | 0.0797 ± 0.0047  | 0.4689 ± 0.0095 |
| 0.8196 ± 0.0011           | 0.0948 ± 0.0047  | 0.7560 ± 0.0079 |
| 1.1574 ± 0.0016           | 0.0615 ± 0.0047  | 0.1898 ± 0.0122 |
| 1.3075 ± 0.0003           | 0.2899 ± 0.0047  | 0.6053 ± 0.0026 |
| 1.4047 ± 0.0007           | 0.1513 ± 0.0047  | 0.4425 ± 0.0050 |
| 1.6139 ± 0.0016           | 0.0631 ± 0.0047  | 0.1496 ± 0.0119 |
| 1.9836 ± 0.0015           | 0.0683 ± 0.0047  | 0.3762 ± 0.0110 |
| 2.5417 ± 0.0009           | 0.1099 ± 0.0047  | 0.9549 ± 0.0069 |
| 2.7875 ± 0.0011           | 0.0897 ± 0.0047  | 0.4184 ± 0.0084 |
| 2.8376 ± 0.0004           | 0.2318 ± 0.0047  | 0.8943 ± 0.0033 |
| 4.0018 ± 0.0009           | 0.1075 ± 0.0047  | 0.2149 ± 0.0070 |
| 4.7099 ± 0.0010           | 0.1028 ± 0.0047  | 0.1232 ± 0.0073 |
| 5.3827 ± 0.0006           | 0.1798 ± 0.0047  | 0.0331 ± 0.0042 |
| 5.4875 ± 0.0014           | 0.0731 ± 0.0047  | 0.9579 ± 0.0103 |
| 5.6319 ± 0.0008           | 0.1216 ± 0.0047  | 0.4694 ± 0.0062 |
| 7.1503 ± 0.0008           | 0.1283 ± 0.0047  | 0.9558 ± 0.0059 |
| 7.6831 ± 0.0008           | 0.1243 ± 0.0047  | 0.9236 ± 0.0061 |
| 7.8308 ± 0.0003           | 0.3680 ± 0.0047  | 0.8973 ± 0.0020 |
| 8.0721 ± 0.0017           | 0.0600 ± 0.0047  | 0.1402 ± 0.0126 |
| 8.3589 ± 0.0005           | 0.1973 ± 0.0047  | 0.1094 ± 0.0038 |
| 8.4197 ± 0.0014           | 0.0718 ± 0.0047  | 0.7448 ± 0.0105 |
| 8.4772 ± 0.0011           | 0.0930 ± 0.0047  | 0.5730 ± 0.0081 |
| 8.5029 ± 0.0005           | 0.1847 ± 0.0047  | 0.6060 ± 0.0041 |
| 8.7186 ± 0.0015           | 0.0690 ± 0.0047  | 0.0109 ± 0.0109 |
| 9.0322 ± 0.0002           | 0.5902 ± 0.0047  | 0.2577 ± 0.0013 |
| 9.0958 ± 0.0013           | 0.0753 ± 0.0047  | 0.3013 ± 0.0100 |
| 9.1773 ± 0.0001           | 1.4365 ± 0.0047  | 0.2618 ± 0.0005 |
| 10.0176 ± 0.0013          | 0.0754 ± 0.0047  | 0.4459 ± 0.0100 |
| 10.3193 ± 0.0009          | 0.1093 ± 0.0047  | 0.8415 ± 0.0069 |
| 10.3798 ± 0.0001          | 1.2726 ± 0.0047  | 0.0622 ± 0.0006 |
| 10.4817 ± 0.0015          | 0.0656 ± 0.0047  | 0.7674 ± 0.0115 |
| 11.0542 ± 0.0008          | 0.1322 ± 0.0047  | 0.1982 ± 0.0057 |
| 11.1685 ± 0.0004          | 0.2789 ± 0.0047  | 0.1763 ± 0.0027 |
| 11.2012 ± 0.0003          | 0.3549 ± 0.0047  | 0.7060 ± 0.0021 |
| 11.7989 ± 0.0006          | 0.1617 ± 0.0047  | 0.7024 ± 0.0047 |
| 11.8698 ± 0.0001          | 0.9092 ± 0.0047  | 0.4085 ± 0.0008 |
| 13.0721 ± 0.0002          | 0.5211 ± 0.0047  | 0.6850 ± 0.0014 |
| 13.2177 ± 0.0002          | 0.5919 ± 0.0047  | 0.1865 ± 0.0013 |
| 13.7346 ± 0.0014          | 0.0715 ± 0.0047  | 0.9988 ± 0.0105 |
| 13.8899 ± 0.0005          | 0.1992 ± 0.0047  | 0.3467 ± 0.0038 |
| 14.3084 ± 0.0013          | 0.0772 ± 0.0047  | 0.0037 ± 0.0098 |
| 14.4155 ± 0.0009          | 0.1104 ± 0.0047  | 0.9842 ± 0.0068 |
| 14.5005 ± 0.0016          | 0.0612 ± 0.0047  | 0.2587 ± 0.0123 |
| 14.5623 ± 0.0004          | 0.2594 ± 0.0047  | 0.0344 ± 0.0029 |
| 15.2349 ± 0.0012          | 0.0842 ± 0.0047  | 0.6404 ± 0.0089 |
| 20.9609 ± 0.0009          | 0.1082 ± 0.0047  | 0.5285 ± 0.0078 |
| 22.3940 ± 0.0010          | 0.0975 ± 0.0047  | 0.7536 ± 0.0077 |
| 23.5953 ± 0.0007          | 0.1417 ± 0.0047  | 0.0482 ± 0.0053 |
| 25.0870 ± 0.0006          | 0.1600 ± 0.0047  | 0.3657 ± 0.0047 |
| 25.7319 ± 0.0020          | 0.0511 ± 0.0047  | 0.9643 ± 0.0148 |
| 26.2917 ± 0.0012          | 0.0807 ± 0.0047  | 0.6437 ± 0.0093 |
| 27.7816 ± 0.0014          | 0.0697 ± 0.0047  | 0.9514 ± 0.0108 |

**Table A2.** Identified spectral lines for VV Ori (p ≡ primary; s ≡ secondary).

| Species       | Order number | Adopted $\lambda$ | Comment                            |
|---------------|--------------|-------------------|------------------------------------|
| He I          | 85           | 6678.149          | well-defined p; s weak, noisy      |
| H $_{\alpha}$ | 87           | 6562.817          | s noticeable in p profile          |
| He I          | 97           | 5875.340          | well-defined p; s weak             |
| Na II         | 97           | 5895.923          | deep, narrow structured<br>D-lines |
| Na II         | 97           | 5889.953          | –                                  |
| Si III        | 99           | 5739.620          | p and s visible                    |
| C III         | 100          | 5696.00           | p only                             |
| N II          | 100          | 5679.56           |                                    |
| Si II         | 101          | 5639.49           |                                    |
| C III         | 109          | 5217.93           |                                    |
| O II          | 109          | 5206.73           |                                    |
| He I          | 112, 113     | 5047.736          | p strong, s visible                |
| He I          | 113          | 5015.675          | p only                             |
| N II          | 114          | 5001.30           |                                    |
| O II          | 115          | 4942.1            | blend                              |
| C III         | 115          | 4922.14           |                                    |
| H $_{\beta}$  | 117          | 4861.332          | s visible but blended              |
| C III         | 118          | 4819.63           | p blend                            |
| N II          | 121          | 4718.4            | weak                               |
| He I          | 120, 121     | 4713.258          | weak p and s                       |
| O II          | 121          | 4699.21           |                                    |
| He II         | 122          | 4685.682          | p only                             |
| O II          | 122          | 4674.213          |                                    |
| C III         | 122          | 4663.53           |                                    |
| C III         | 122          | 4650.160          |                                    |
| C III + O II  | 123          | 4649.86           | strong blend                       |
| N II          | 123          | 4630.567          |                                    |
| N II          | 123          | 4621.39           |                                    |
| O II          | 123          | 4610.14           |                                    |
| O II          | 124          | 4602.11           |                                    |
| O II          | 124          | 4596.174          |                                    |
| C III         | 124          | 4593.47           |                                    |
| Si III        | 125          | 4574.78           | edge of order                      |
| Si III        | 125          | 4569.67           | blend                              |
| C III         | 125          | 4567.872          |                                    |
| Si III        | 125          | 4552.65           | blend                              |

This paper has been typeset from a  $\text{T}_{\text{E}}\text{X}/\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$  file prepared by the author.