

Odderon mechanism for transverse single spin asymmetry in the Wandzura-Wilczek approximation

Benić, Sanjin; Horvatić, Davor; Kaushik, Abhiram; Vivoda, Eric Andreas

Source / Izvornik: **Physical Review D**, 2022, 106

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1103/PhysRevD.106.114025>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:217:335415>

Rights / Prava: [Attribution 4.0 International](#)/[Imenovanje 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-03-24**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



Odderon mechanism for transverse single spin asymmetry in the Wandzura-Wilczek approximation

Sanjin Benić, Davor Horvatić[✉], Abhiram Kaushik, and Eric Andreas Vivoda[✉]

*Department of Physics, Faculty of Science, University of Zagreb,
Bijenička cesta 32, 10000 Zagreb, Croatia*

 (Received 24 October 2022; accepted 5 December 2022; published 26 December 2022)

We compute the transverse single spin asymmetry in forward $p^\uparrow p \rightarrow hX$ and $p^\uparrow A \rightarrow hX$ collisions from the odderon mechanism originally suggested by Kovchegov and Sievert [Phys. Rev. D **86**, 034028 (2012)]. Working in the hybrid approach of the color glass condensate effective theory, we first identify the relevant collinear parton distribution function (PDF) of the transversely polarized proton p^\uparrow as the intrinsic twist-3 $g_T(x)$ distribution. We further argue that the complete polarized cross section also contains contributions from the kinematical and the dynamical twist-3 PDFs, in addition to the intrinsic twist-3 PDF. By restricting to the Wandzura-Wilczek approximation, where the dynamical twist-3 PDFs are dropped, we find that the odderon contribution to the polarized cross section for inclusive hadron production is exactly zero at the next-to-leading order in the strong coupling.

DOI: 10.1103/PhysRevD.106.114025

I. INTRODUCTION AND MOTIVATION

Transverse single spin asymmetry (SSA) [1–4] is a phenomenon associated with azimuthally asymmetric particle production in collisions involving a transversely polarized proton p^\uparrow . SSA is characterized by a sine modulation $\mathbf{P}_{h\perp} \times \mathbf{S}_\perp = P_{h\perp} S_\perp \sin(\phi_h - \phi_S)$. Here, $\mathbf{P}_{h\perp}$ is the transverse momentum of the produced hadron, and \mathbf{S}_\perp is the spin of the transversely polarized proton. Decades of dedicated measurements have demonstrated its persistence even at the highest collision energies, with the SSA being largest in the forward region of the produced particle, typically a hadron. This is so across different collision systems such as ep^\uparrow and $p^\uparrow p$ but also most recently for $p^\uparrow A$ collisions [5,6].

On the theory front, it is known that the presence of the phase in the cross section is crucial to generate SSA. In the forward region, where the momentum fraction x in the target is small, one naturally expects the phenomenon of gluon saturation [7–10] to play an important role in determining SSA [11–19]. In this work, we are revisiting the computation by Kovchegov and Sievert [11], where they used the color glass condensate (CGC) effective theory [7–10] for gluon saturation, to suggest a new mechanism

for SSA. The special property of this mechanism is in supplying the phase by the odderon distribution [20–22]

$$\mathcal{O}(\mathbf{x}_\perp, \mathbf{y}_\perp) \equiv \frac{1}{2iN_c} \text{tr}(V(\mathbf{x}_\perp)V^\dagger(\mathbf{y}_\perp) - V(\mathbf{y}_\perp)V^\dagger(\mathbf{x}_\perp)), \quad (1)$$

that is, the imaginary part of the dipole distribution $\text{tr}(V(\mathbf{x}_\perp)V^\dagger(\mathbf{y}_\perp))/N_c$. Here, $V(\mathbf{x}_\perp)$ is a fundamental Wilson line with $\langle \dots \rangle$ denoting the color average. This “odderon mechanism,” as we will refer to it in this work, leads to a substantial A suppression of SSA, $\sim A^{-7/6}$ parametrically [11].

In the following Sec. II, we take as a starting point the polarized cross section in the hybrid approach [14–17] with a transversely polarized proton described by the collinear twist-3 PDFs and the dense target (a nuclei or a proton in the forward collision) by Wilson line correlators arising from the CGC. In the context of the twist-3 parton distribution functions (PDFs), the computation of Ref. [11] is clarified in terms of the $g_T(x)$ distribution. We carefully emphasize, however, that the complete twist-3 hadronic cross section contains additional terms associated with the kinematic twist-3 function $g_{1T}^{(1)}(x)$, that is, the first moment of the worm-gear transverse-momentum dependent (TMD) distributions [23], as well as the dynamical Efremov-Teryaev-Qiu-Sterman (ETQS) functions [24,25]; see, e.g., Eq. (3) below. Working in the Wandzura-Wilczek (WW) approximation [26], that neglects the dynamical twist-3 part of the cross section, our computation first confirms that the cross section is proportional to the odderon distribution (1). However, as we explicitly show

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

in Secs. III–V, due to the specific form of the resulting hard factor, the odderon contribution to SSA for inclusive hadron production turns out to be exactly zero at the next-to-leading order (NLO) in the strong coupling α_S for all possible partonic channels. In the concluding Sec. VI, we also briefly outline several new ways the odderon could appear in SSA after all.

II. GENERAL REMARKS

In the hybrid approach, a dilute projectile proton is described using collinear PDFs, while the distributions of the dense target (nuclei, or a proton in forward collisions) are given in terms of Wilson line correlators. To set up our notations, we first write down the unpolarized $pA \rightarrow hX$ cross section in terms of the familiar twist-2 PDFs and fragmentation functions (FFs). For convenience, this is given in the following way:

$$E_h \frac{d\sigma}{d^3P_h} \equiv \frac{1}{2(2\pi)^3} \int \frac{dz_h}{z_h^2} D(z_h) \times \int dx_p \left\{ \frac{1}{2} f(x_p) \text{Tr}[\not{P}_p S^{(0)}(p_1)] + \frac{1}{2} G(x_p) (-g_{\perp}^{\alpha\beta}) S_{\alpha\beta}^{(0)}(p_1) \right\}, \quad (2)$$

where we have separated out the twist-2 hadron FF $D(z_h)$ and the twist-2 PDF in the quark (gluon) $f(x_p)$ [$G(x_p)$] initiated channel. The proton and the nucleus move along

the light cone with momenta in the center-of-mass frame given as $P_p^+ = P_A^- = \sqrt{s/2}$, where s is the collision energy squared per nucleon,¹ while P_h is the momenta of the produced hadron h . Here and in the following, we use the light-cone variables $p^\pm = (p^0 \pm p^3)/\sqrt{2}$. Furthermore, p_1 is the momentum of the parton moving collinearly with the proton $p_1 = x_p P_p$, and $g_{\perp}^{\alpha\beta} = g^{\alpha\beta} - n^\alpha \bar{n}^\beta - \bar{n}^\alpha n^\beta$ with $n^\alpha = \delta^{\alpha-}$, $\bar{n}^\alpha = \delta^{\alpha+}$ is a transverse projector to the physical gluon polarizations. $S^{(0)}(p_1)$ [$S_{\alpha\beta}^{(0)}(p_1)$] is an all-order two-parton scattering kernel in the quark (gluon) initiated channel containing the hard factor and also the target distribution that we will be computing within the CGC hybrid approach. We have absorbed the CGC flux factor $1/2P_p^+$ into the definition of $S^{(0)}(p_1)$.

A. Polarized cross section

In order to generalize to collisions with a transversely polarized proton, the polarized cross section $d\Delta\sigma$ is computed up to twist 3 in the polarized proton PDF. We will be restricting here to the usual twist-2 FF, $D(z_h)$ —the particular contribution arising from twist-3 FFs [27,28] has been computed in $p^\uparrow A$ [17]. We are also not considering various pole contributions to $d\Delta\sigma$; see [16]. Our starting point is a separate (nonpole) contribution that has already been discussed in semi-inclusive deep inelastic scattering (SIDIS) [29,30]. Adapting to the $p^\uparrow A$ computation, we have the following gauge-invariant all-order expression²:

$$E_h \frac{d\Delta\sigma}{d^3P_h} = \frac{1}{2(2\pi)^3} \int \frac{dz_h}{z_h^2} D(z_h) \left\{ \frac{M_N}{2} \int dx_p g_T(x_p) \text{Tr}[\gamma_5 \not{S}_\perp S^{(0)}(p_1)] + \frac{M_N}{2} \int dx_p g_{1T}^{(1)}(x_p) \text{Tr} \left[\gamma_5 \not{P}_p S_\perp^\lambda \left(\frac{\partial S^{(0)}(k_1)}{\partial k_{1\perp}^2} \right)_{k_1=p_1} \right] + \frac{iM_N}{4} \int dx_p dx'_p \text{Tr} \left[\left(\not{P}_p e^{\bar{n}n\lambda S_\perp} \frac{G_F(x_p, x'_p)}{x_p - x'_p} + i\gamma_5 \not{P}_p S_\perp^\lambda \frac{\tilde{G}_F(x_p, x'_p)}{x_p - x'_p} \right) S_\lambda^{(1)}(x_p P_p, x'_p P_p) \right] \right\}. \quad (3)$$

The first part of Eq. (3) is arising from the $g_T(x_p)$ distribution function. This is also referred to as an *intrinsic* (i.e., $\sim \langle P_p S_\perp | \bar{\psi} \psi | P_p S_\perp \rangle$) contribution. The second part is a *kinematical* ($\langle P_p S_\perp | \bar{\psi} \partial \psi | P_p S_\perp \rangle$) contribution that is proportional to $g_{1T}^{(1)}(x_p)$, namely, the first moment of the worm-gear TMD.³ In the third (*dynamical* $\sim \langle P_p S_\perp | \bar{\psi} F \psi | P_p S_\perp \rangle$) part, we have the ETQS distributions $G_F(x_p, x'_p)$ and $\tilde{G}_F(x_p, x'_p)$. Note the appearance of the same two-parton scattering kernel $S^{(0)}(p_1)$ as in Eq. (2).

To compute the cross section, we also need its finite- $k_{1\perp}$ variant $S^{(0)}(k_1)$, as well as $S_\lambda^{(1)}(x_p P_p, x'_p P_p)$, which is a three-parton scattering kernel containing an additional gluon from the polarized proton. We should appreciate the appearance of the $k_{1\perp}$ derivative as a consequence of performing the computation up to twist 3 but also due to the connection of $\partial S^{(0)}(k_1)/\partial k_{1\perp}$ with $S_\lambda^{(1)}(x_p P_p, x'_p P_p)$ through the Ward identity for the gluon from the proton [29,30].

We point out here that the computation in Ref. [11] is on the parton level, taking transversely polarized spinors $u(p, S_\perp)$ for the initial quark. Thanks to the decomposition $u(p, S_\perp) \bar{u}(p, S_\perp) = (\not{p} + m)(1 + \gamma_5 \not{S}_\perp)/2$ [33], only the $\frac{m}{2} \gamma_5 \not{S}_\perp$ piece is relevant for the polarized quark in the current context (the remaining S_\perp -dependent term eventually gets interpreted as the transversity PDF, but this does

¹Here, P_A^- is the center-of-mass momentum per nucleon.

²We are using the convention $\epsilon_{0123} = +1 = -\epsilon^{0123}$ and $\gamma_5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$.

³Instead of $g_{1T}^{(1)}(x)$, sometimes a function $\tilde{g}(x)$ is used [31], with the relation $\tilde{g}(x) = -2g_{1T}^{(1)}(x)$ [32].

not contribute in what follows). Thus, the computation in Ref. [11] clearly corresponds to the term in Eq. (3) that is proportional to $g_T(x_p)$. Namely, when changing from a polarized free quark projectile to a polarized nucleon projectile, one naturally replaces the quark mass m by the nucleon mass M_N .

The above introduced distributions satisfy the QCD equation of motion identity [31,32]

$$xg_T(x) = g_{1T}^{(1)}(x) - \frac{1}{2} \int dx' \frac{G_F(x, x') + \tilde{G}_F(x, x')}{x - x'}. \quad (4)$$

The $g_T(x)$ distribution satisfies another important relation connecting it to the twist-2 helicity PDF $\Delta q(x)$ [31,32], thus revealing that $g_T(x)$ itself has a twist-2 piece

$$E_h \frac{d\Delta\sigma}{d^3P_h} \simeq \frac{1}{2(2\pi)^3} \frac{M_N}{2} \int \frac{dz_h}{z_h^2} D(z_h) \int dx_p g_T(x_p) \times \left(S_{\perp}^{\lambda} \frac{\partial}{\partial k_{1\perp}^{\lambda}} \text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] \right)_{k_1=p_1}, \quad (6)$$

that we will take as the starting point of our explicit computations below.

The attractive feature of the WW approximation is that we can compute $g_T(x)$ [via Eq. (5)] and, thus, also Eq. (6) from, say, global fits of the helicity PDFs. The distribution $g_T(x)$ has been recently studied on the lattice [34], and a separate global analysis was performed to constrain the $g_{1T}^{(1)}(x)$ distributions in Ref. [35]. The general conclusion from these works is that, given the current uncertainties, both $g_T(x)$ and $g_{1T}^{(1)}(x)$ are roughly consistent with the WW approximation. While this provides a justification for the present computation, we acknowledge that the current uncertainties found in the aforementioned works do

$$g_T(x) = \int_x^1 \frac{dx'}{x'} \Delta q(x') + (\text{genuine twist} - 3). \quad (5)$$

The remainder is given in terms of the ETQS functions; see [31,32] for the explicit expression. Our computation will be based on the WW approximation, that is, ignoring genuine (dynamical) twist-3 contributions *everywhere*. This, in particular, means that we are taking into account the $g_T(x_p)$ and the $g_{1T}^{(1)}(x_p)$ contributions to $d\Delta\sigma$ in Eq. (3), while we are ignoring the genuine twist-3 contributions in the definition of $g_T(x)$ in Eq. (5) and in the equation of motion identity, Eq. (4). This allows us to fix $g_{1T}^{(1)}(x)$ through Eq. (4) as $g_{1T}^{(1)}(x) \simeq xg_T(x)$, rendering Eq. (3) into the following compact form:

accommodate sizeable deviations (even up to 40% in certain regions of x) from the WW approximation. A complete computation that goes beyond the WW approximation does not mean only considering $g_T(x)$ or $g_{1T}^{(1)}(x)$ with genuine twist-3 pieces included, but would also require dealing with the full complexity of the dynamical twist-3 contributions at NLO contained in the second line of Eq. (3). While we reflect on this in the concluding section, a full NLO computation with genuine twist-3 effects is beyond the scope of this work.

The analogous expression for the gluon initiated channel is adapted from Eqs. (17) and (25) in Ref. [36] [see also Eq. (32) in Ref. [37]] to read

$$\begin{aligned} E_h \frac{d\Delta\sigma}{d^3P_h} = & \frac{1}{2(2\pi)^3} \int \frac{dz_h}{z_h^2} D(z_h) \left[iM_N \int dx_p \mathcal{G}_{3T}(x_p) \frac{1}{P_1^+} e^{n\alpha\beta S_{\perp}} S^{(0)\alpha\beta'}(p_1) \omega_{\alpha\alpha'} \omega_{\beta'\beta} \right. \\ & - iM_N \int \frac{dx_p}{x_p^2} \tilde{g}(x_p) (g_{\perp}^{\beta\lambda} e^{a\bar{n}nS_{\perp}} - g_{\perp}^{\alpha\lambda} e^{\beta\bar{n}nS_{\perp}}) \left(\frac{\partial S_{\alpha\beta}^{(0)}(k_1)}{\partial k_1^{\lambda}} \right)_{k_1=p_1} \\ & \left. - \frac{1}{2} \int \frac{dx_p dx'_p}{x_p x'_p} M_F^{\alpha\beta\gamma}(x_p, x'_p) \frac{S^{(1)\alpha\beta'\gamma'}(x_p P_p, x'_p P_p)}{x'_p - x_p} \omega_{\alpha\alpha'} \omega_{\beta'\beta} \omega_{\gamma'\gamma} \right], \quad (7) \end{aligned}$$

where $\omega_{\alpha\beta} = g_{\alpha\beta} - \bar{n}_{\alpha} n_{\beta}$. In the first line, we have the intrinsic contribution with $\mathcal{G}_{3T}(x_p)$ being the gluonic counterpart of $g_T(x_p)$. In the second line, $\tilde{g}(x_p)$ is the gluonic kinematical function (see [38,39] for the definition), and $M_F^{\alpha\beta\gamma}(x_p, x'_p)$ is the three-gluon correlator. In the WW approximation, $\mathcal{G}_{3T}(x)$ becomes related to the gluon helicity PDF $\Delta G(x)$ as [38]

$$\mathcal{G}_{3T}(x) \simeq \frac{1}{2} \int_x^1 \frac{dx'}{x'} \Delta G(x'), \quad (8)$$

while $\tilde{g}(x) \simeq x^2 \mathcal{G}_{3T}(x)$ [38]. The WW truncation then amounts to the first two lines of Eq. (7).

B. A recap of the leading-order inclusive hadron production

The leading-order (LO) amplitude for inclusive hadron production from the $q(k_1) \rightarrow q(q)$ channel in the $n \cdot A = A^+ = 0$ gauge is simply given as [40]

$$\mathcal{M} = \gamma^+ \int_{\mathbf{x}_\perp} e^{i(\mathbf{q}_\perp - \mathbf{k}_{1\perp}) \cdot \mathbf{x}_\perp} [V(\mathbf{x}_\perp) - 1]. \quad (9)$$

Here, $V(\mathbf{x}_\perp) = \mathcal{P} \exp [ig \int_{-\infty}^{\infty} dx^+ A_a^-(x) t^a]$ is the fundamental Wilson line with $A_a^-(x)$ being the classical field of the target, and we use $\int_{\mathbf{x}_\perp} \equiv \int d^2\mathbf{x}_\perp$. In Eq. (9), we have omitted the overall light-cone delta function $(2\pi)\delta(k_1^+ - q^+)$, as well as the initial and final state spinors, thus leaving a matrix in spinor (and color) space. From \mathcal{M} , we obtain the leading-order result for $S^{(0)}(k_1)$ as

$$\begin{aligned} S^{(0)}(k_1) &\equiv \frac{1}{2P_p^+} \frac{1}{N_c} \langle \bar{\mathcal{M}} \not{q} \mathcal{M} \rangle (2\pi) \delta(k_1^+ - q^+) \\ &= (2\pi) \delta(k_1^+ - q^+) \frac{x_p}{2k_1^+} \gamma^+ \not{q} \gamma^+ \\ &\quad \times \int_{\mathbf{x}_\perp, \mathbf{x}'_\perp} \mathcal{S}(\mathbf{x}_\perp, \mathbf{x}'_\perp) e^{i(\mathbf{q}_\perp - \mathbf{k}_{1\perp}) \cdot (\mathbf{x}_\perp - \mathbf{x}'_\perp)}, \end{aligned} \quad (10)$$

where $1/2P_p^+$ is the flux factor, $1/N_c$ is coming from averaging over the color of the initial state quark, and

$$\mathcal{S}(\mathbf{x}_\perp, \mathbf{x}'_\perp) \equiv \frac{1}{N_c} \text{tr} \langle V(\mathbf{x}_\perp) V^\dagger(\mathbf{x}'_\perp) \rangle \quad (11)$$

is the color averaged dipole distribution. Here and in the following, we are suppressing the dependence of the nuclear distributions on the momentum fraction $x_A = k_2^-/P_A^-$, where k_2 is the partonic momenta from the nuclei. At the LO we have the momentum conservation $k_1 + k_2 = q$. We readily conclude that $\text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] \sim \text{tr}[\gamma_5 \not{k}_1 \gamma^+ \not{q} \gamma^+] \sim e^{++qk_1} = 0$, and so Eq. (6) vanishes at the LO. As we also have $\text{tr}[\gamma_5 \not{g}_\perp S^{(0)}(p_1)] = 0$, the $g_T(x)$ and the $g_{1T}^{(1)}(x)$ terms in Eq. (3) vanish separately at the LO.

The analogous expressions in the $g(k_1) \rightarrow g(k_g)$ channel are [41]

$$\mathcal{M} = (-2k_1^+) \int_{\mathbf{x}_\perp} e^{i(\mathbf{k}_{g\perp} - \mathbf{k}_{1\perp}) \cdot \mathbf{x}_\perp} [U(\mathbf{x}_\perp) - 1] \quad (12)$$

and

$$\begin{aligned} S_{\alpha\beta}^{(0)}(k_1) &= \frac{1}{2P_p^+} \frac{1}{N_c^2 - 1} \langle \mathcal{M}^\dagger \mathcal{M} \rangle d_{\alpha\beta}(k_g) \\ &= (2\pi) \delta(k_1^+ - k_g^+) (2k_1^+)^2 d_{\alpha\beta}(k_g) \\ &\quad \times \int_{\mathbf{x}_\perp, \mathbf{x}'_\perp} \mathcal{S}_A(\mathbf{x}_\perp, \mathbf{x}'_\perp) e^{i(\mathbf{k}_{g\perp} - \mathbf{k}_{1\perp}) \cdot (\mathbf{x}_\perp - \mathbf{x}'_\perp)}, \end{aligned} \quad (13)$$

where

$$d^{\alpha\beta}(k) = -g^{\alpha\beta} + \frac{n^\alpha k^\beta + n^\beta k^\alpha}{k^+} \quad (14)$$

is the gluon polarization tensor. Note that $d^{\alpha\beta}(p_1) = -g_\perp^{\alpha\beta}$. $U(\mathbf{x}_\perp)$ is the adjoint Wilson line, and

$$\mathcal{S}_A(\mathbf{x}_\perp, \mathbf{x}'_\perp) \equiv \frac{1}{N_c^2 - 1} \text{tr} \langle U(\mathbf{x}_\perp) U^\dagger(\mathbf{x}'_\perp) \rangle \quad (15)$$

is the adjoint dipole distribution. The contribution from this channel also vanishes at the LO simply due to the realness of the adjoint Wilson line.

III. NLO INCLUSIVE HADRON PRODUCTION IN $p^\uparrow A \rightarrow hX$: THE $q \rightarrow qg$ CHANNEL

At the NLO, we have, in general, the $q \rightarrow qg$, $g \rightarrow q\bar{q}$, and $g \rightarrow gg$ channels. In this section, we compute the $q \rightarrow qg$ channel (together with the accompanying virtual contribution), while the $g \rightarrow q\bar{q}$ and the $g \rightarrow gg$ channels are discussed separately in Secs. IV and V, respectively.

A. $q \rightarrow q$: Real contribution

We consider the $q(k_1) \rightarrow q(q)g(k_g)$ partonic channel, where in the final state a real gluon with momentum k_g gets radiated in addition to the quark with momentum q . Here, k_2 is set by momentum conservation at NLO: $k_1 + k_2 = q + k_g$. We will focus on the case where the quark fragments into a final state hadron and we integrate over the (untagged) gluon phase space to compute the inclusive hadron cross section according to Eq. (6). The main quantity to compute is $S^{(0)}(k_1)$, which takes the following form:

$$\begin{aligned} S^{(0)}(k_1) &= \frac{1}{2P_p^+} \int \frac{d^3k_g}{(2\pi)^3 2E_g} \frac{1}{N_c} \langle \bar{\mathcal{M}}^{\mu'} \not{q} \mathcal{M}^\mu \rangle d_{\mu\mu'}(k_g) \\ &\quad \times (2\pi) \delta(k_1^+ - q^+ - k_g^+) \\ &= \frac{q^+}{P_p^+} \int_{k_{g\perp}} \frac{1}{N_c} \frac{1}{4q^+ k_g^+} \langle \bar{\mathcal{M}}^{\mu'} \not{q} \mathcal{M}^\mu \rangle d_{\mu\mu'}(k_g), \end{aligned} \quad (16)$$

where $\int_{k_{g\perp}} \equiv \int \frac{d^2k_{g\perp}}{(2\pi)^2}$ and similar for other transverse momenta integrations. Using the quark and gluon propagators in the CGC background [42–44], we can compute the following amplitudes:

$$\begin{aligned}
\mathcal{M}_1^\mu &= -ig\gamma^\mu \frac{\not{q} + \not{k}_g}{(q + k_g)^2 + i\epsilon} \gamma^+ \int_{\mathbf{x}_\perp} e^{i\mathbf{k}_{2\perp} \cdot \mathbf{x}_\perp} t^a [V(\mathbf{x}_\perp) - 1], \\
\mathcal{M}_2^\mu &= -ig\gamma^+ \frac{\not{k}_1 - \not{k}_g}{(k_1 - k_g)^2 + i\epsilon} \gamma^\mu \int_{\mathbf{x}_\perp} e^{i\mathbf{k}_{2\perp} \cdot \mathbf{x}_\perp} [V(\mathbf{x}_\perp) - 1] t^a, \\
\mathcal{M}_3^\mu &= ig(2k_g^+) \gamma_\nu \frac{d^{\nu\mu}(k_1 - q)}{(k_1 - q)^2 + i\epsilon} \int_{\mathbf{x}_\perp} e^{i\mathbf{k}_{2\perp} \cdot \mathbf{x}_\perp} t^b [U^{ab}(\mathbf{x}_\perp) - \delta^{ab}], \\
\mathcal{M}_4^\mu &= -g(2k_g^+) \int_{\mathbf{k}_\perp} \int_{-\infty}^{\infty} \frac{dk^-}{(2\pi)} \gamma^+ \frac{\not{q} - \not{k}}{(q - k)^2 + i\epsilon} \gamma_\nu \frac{d^{\nu\mu}(k_1 + k - q)}{(k_1 + k - q)^2 + i\epsilon} \\
&\quad \times \int_{\mathbf{x}_\perp \mathbf{y}_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} [V(\mathbf{x}_\perp) - 1] t^b [U^{ab}(\mathbf{y}_\perp) - \delta^{ab}], \tag{17}
\end{aligned}$$

with the total amplitude \mathcal{M} given as $\mathcal{M}^\mu = \sum_{k=1}^4 \mathcal{M}_k^\mu$. We find that the result (17) agrees with Ref. [45] except for the overall sign and the adjoint indices in the eikonal gluon vertex that enters \mathcal{M}_3^μ and \mathcal{M}_4^μ ; see, for example, Eq. (3.2) in Ref. [46]. In the special case when the final quark and gluon are on shell,⁴ \mathcal{M}^μ takes the following simple form:

$$\begin{aligned}
\mathcal{M}^\mu &= -ig \int_{\mathbf{k}_\perp} \int_{\mathbf{x}_\perp \mathbf{y}_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} [T_q^\mu t^a V(\mathbf{x}_\perp) \\
&\quad + T_{qg}^\mu(\mathbf{k}_\perp) V(\mathbf{x}_\perp) t^b U^{ab}(\mathbf{y}_\perp)], \tag{18}
\end{aligned}$$

where

$$T_q^\mu = \gamma^\mu \frac{\not{q} + \not{k}_g}{(q + k_g)^2} \gamma^+ \tag{19}$$

and

$$T_{qg}^\mu(\mathbf{k}_\perp) = -\gamma^+ \frac{\not{q} - \not{k}}{(q - k)^2} \gamma_\nu d^{\nu\mu}(k_1 + k - q). \tag{20}$$

Here, we evaluated the k^- integral in favor of $(k_1 + k - q)^2 + i\epsilon = 0$ so that

$$(q - k)^2 = -\frac{1}{k_1^+ k_g^+} [q^+ \mathbf{k}_{1\perp} + k_1^+ (\mathbf{k}_\perp - \mathbf{q}_\perp)]^2. \tag{21}$$

Inserting Eq. (17) into Eq. (16), we find

$$\begin{aligned}
S^{(0)}(k_1) &= \frac{q^+}{P_p^+} \frac{g^2 C_F}{4q^+ k_g^+} \int_{\mathbf{k}_{g\perp} \mathbf{k}_{1\perp} \mathbf{k}'_\perp} \int_{\mathbf{x}_\perp \mathbf{x}'_\perp \mathbf{y}_\perp \mathbf{y}'_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} \\
&\quad \times e^{-i\mathbf{k}'_\perp \cdot \mathbf{x}'_\perp} e^{-i(\mathbf{k}_{2\perp} - \mathbf{k}'_\perp) \cdot \mathbf{y}'_\perp} d_{\mu\mu'}(k_g) [\mathcal{S}(\mathbf{x}_\perp, \mathbf{x}'_\perp) \bar{T}_q^\mu \not{q} T_q^\mu + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}'_\perp) \bar{T}_{qg}^{\mu'}(\mathbf{k}'_\perp) \not{q} T_q^\mu \\
&\quad + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp) \bar{T}_q^{\mu'} \not{q} T_{qg}^\mu(\mathbf{k}_\perp) + \mathcal{S}_{qqgq}(\mathbf{x}'_\perp, \mathbf{y}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp) \bar{T}_{qg}^{\mu'}(\mathbf{k}'_\perp) \not{q} T_{qg}^\mu(\mathbf{k}_\perp)], \tag{22}
\end{aligned}$$

where $\mathcal{S}(\mathbf{x}_\perp, \mathbf{x}'_\perp)$ is the dipole defined in Eq. (11) and the additional distributions are given as

$$\begin{aligned}
\mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}'_\perp) &\equiv \frac{1}{C_F N_c} \langle \text{tr}(V^\dagger(\mathbf{x}'_\perp) t^b V(\mathbf{x}_\perp) t^a) U^{ba}(\mathbf{y}'_\perp) \rangle, \\
\mathcal{S}_{qqgq}(\mathbf{x}'_\perp, \mathbf{y}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp) &\equiv \frac{1}{C_F N_c} \langle \text{tr}(V^\dagger(\mathbf{x}'_\perp) V(\mathbf{x}_\perp) t^a t^b) [U^\dagger(\mathbf{y}'_\perp) U(\mathbf{y}_\perp)]^{ba} \rangle. \tag{23}
\end{aligned}$$

We now show that the first and the fourth terms in the square brackets in Eq. (22) do not contribute to the polarized cross section when integrated over the gluon

⁴Without loss of generality, the initial quark can be taken as on shell even at finite $\mathbf{k}_{1\perp}$, that is, $k_1^2 = 0$. Equation (6) is not affected due to the $\partial/\partial k_{1\perp}^2$ derivative.

momenta. This is intuitively clear, as the SSA must come from interferences of different amplitudes, that are given by the second and the third terms (cf. Fig. 1), while the first and the fourth term are squares of amplitudes. The analogous structure can also be identified in the computation in Ref. [11]. Consider the first term in Eq. (22), where in the context of Eq. (6) we have

$$\mathcal{S}_{qqg}(\mathbf{x}'_{\perp}, \mathbf{x}_{\perp}, \mathbf{y}'_{\perp}) \simeq \frac{1}{2C_F N_c} (N_c^2 \mathcal{S}(\mathbf{y}'_{\perp}, \mathbf{x}'_{\perp}) \mathcal{S}(\mathbf{x}_{\perp}, \mathbf{y}'_{\perp}) - \mathcal{S}(\mathbf{x}_{\perp}, \mathbf{x}'_{\perp})). \quad (32)$$

The second (dipole) term in Eq. (32) drops out when combined with the hard factors in Eq. (30). To see this, note that the \mathbf{y}_{\perp} and the \mathbf{y}'_{\perp} integrations in Eq. (30) result in δ functions that yield $\mathbf{k}'_{\perp} = \mathbf{k}_{\perp} = \mathbf{k}_{2\perp}$. This gives $-\mathcal{H}(\mathbf{k}_{2\perp}, \mathbf{k}_{1\perp}) + \mathcal{H}(\mathbf{k}_{2\perp}, \mathbf{k}_{1\perp}) = 0$.

Now we split the dipole into its real and imaginary parts [21]:

$$\mathcal{S}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) = \mathcal{P}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) + i\mathcal{O}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}), \quad (33)$$

where

$$\begin{aligned} \mathcal{P}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) &\equiv \frac{1}{2} (\mathcal{S}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) + \mathcal{S}(\mathbf{y}_{\perp}, \mathbf{x}_{\perp})), \\ \mathcal{O}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) &\equiv \frac{1}{2i} (\mathcal{S}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) - \mathcal{S}(\mathbf{y}_{\perp}, \mathbf{x}_{\perp})) \end{aligned} \quad (34)$$

are the pomeron and the odderon distributions, respectively. We also replace the primed and unprimed transverse coordinate and momenta labels in the first term in Eq. (30), namely, $\mathbf{k}'_{\perp} \leftrightarrow \mathbf{k}_{\perp}$, $\mathbf{x}'_{\perp} \leftrightarrow \mathbf{x}_{\perp}$, and $\mathbf{y}'_{\perp} \leftrightarrow \mathbf{y}_{\perp}$. By compensating for the reversed sign in the exponentials with $\mathbf{x}_{\perp} \rightarrow -\mathbf{x}_{\perp}$, and using overall invariance under reflections for the distributions in the unpolarized target, we obtain

$$\begin{aligned} &\text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] \\ &= ig^2 N_c \frac{q^+}{P_p^+} \int_{\mathbf{k}_{2\perp}, \mathbf{k}_{\perp}} \int_{\mathbf{x}_{\perp}, \mathbf{x}'_{\perp}, \mathbf{y}_{\perp}} e^{i\mathbf{k}_{\perp} \cdot (\mathbf{x}_{\perp} - \mathbf{y}_{\perp})} e^{-i\mathbf{k}_{2\perp} \cdot (\mathbf{x}'_{\perp} - \mathbf{y}_{\perp})} \\ &\quad \times [\mathcal{P}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) \mathcal{O}(\mathbf{x}'_{\perp}, \mathbf{y}_{\perp}) - \mathcal{O}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp}) \mathcal{P}(\mathbf{x}'_{\perp}, \mathbf{y}_{\perp})] \mathcal{H}(\mathbf{k}_{\perp}, \mathbf{k}_{1\perp}). \end{aligned} \quad (35)$$

Here, we have used the following symmetry properties: $\mathcal{P}(\mathbf{y}_{\perp}, \mathbf{x}_{\perp}) = \mathcal{P}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp})$ and $\mathcal{O}(\mathbf{y}_{\perp}, \mathbf{x}_{\perp}) = -\mathcal{O}(\mathbf{x}_{\perp}, \mathbf{y}_{\perp})$, which follow from Eq. (34). We have also passed from $\mathbf{k}_{g\perp}$ to $\mathbf{k}_{2\perp}$ integration. This result clearly demonstrates that the polarized cross section is proportional to the odderon operator.

The Dirac trace is easy to calculate, and we find

$$\mathcal{H}(\mathbf{k}_{\perp}, \mathbf{k}_{1\perp}) = 4i(\bar{z} + 1) \frac{\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp}}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2}, \quad (36)$$

where $\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp} \equiv \epsilon^{-+v_{1\perp}v_{2\perp}} = v_{1\perp} v_{2\perp} \sin(\phi_1 - \phi_2)$ and

$$\begin{aligned} \mathbf{v}_{1\perp} &\equiv z\mathbf{q}_{\perp} - \bar{z}\mathbf{k}_{g\perp} = \mathbf{q}_{\perp} - \bar{z}\mathbf{k}_{1\perp} - \bar{z}\mathbf{k}_{2\perp}, \\ \mathbf{v}_{2\perp} &\equiv \mathbf{q}_{\perp} - \bar{z}\mathbf{k}_{1\perp} - \mathbf{k}_{\perp}. \end{aligned} \quad (37)$$

Equations (35) and (36) represent the main results of this section. The vectors $\mathbf{v}_{1\perp}$ and $\mathbf{v}_{2\perp}$ reflect the collinear gluon

radiations so that when $\mathbf{v}_{1\perp} \rightarrow 0$ ($\mathbf{v}_{2\perp} \rightarrow 0$) the radiated gluon would be collinear to the final (initial) state quark. Note, however, that in Eq. (36) both of these limits are completely finite, meaning that the usual collinear divergences one encounters in the NLO computations of an unpolarized cross section for inclusive hadron production (see, for example, [47]) are absent in this particular computation of the polarized cross section. In addition, when $z \rightarrow 0$ ($\bar{z} \rightarrow 1$), i.e., when the radiated gluon is collinear to the nucleus ($k_g^+ \rightarrow 0$ and so $k_g^- \rightarrow \infty$ effectively), the hard factor is also finite. In fact, a close inspection reveals that the resulting cross section is zero in this limit. Namely, when $z \rightarrow 1$ there is a symmetry in the hard factor such that, by interchanging $\mathbf{k}_{\perp} \leftrightarrow \mathbf{k}_{2\perp}$ so that $\mathbf{v}_{1\perp} \leftrightarrow \mathbf{v}_{2\perp}$, the hard factor picks up a sign $\mathcal{H} \rightarrow -\mathcal{H}$. On the other hand, the soft part in Eq. (35) is even under such a transformation, and so the overall cross section is zero in this limit. In the case of the NLO unpolarized cross section, the $z \rightarrow 0$ divergence recovers a part of the small- x evolution of the nuclear wave function [47].

We reflect here also on the computation in Ref. [11] that takes into account only the $g_T(x_p)$ contribution (on the parton level) in Eq. (6). The resulting hard factor associated with $g_T(x)$ is found to be

$$\begin{aligned} \mathcal{H}^{(gr)}(\mathbf{k}_{\perp}) &= \frac{1}{4q^+ k_g^+} d_{\mu\mu'}(k_g) \text{tr}[\gamma_5 \not{s}_{\perp} \bar{T}_q^{\mu} \not{q} T_{qg}^{\mu'}(\mathbf{k}_{\perp})]_{\mathbf{k}_{1\perp}=0} \\ &= 4i\bar{z}^2 \frac{\hat{\mathbf{v}}_{1\perp} \times \mathbf{S}_{\perp}}{\hat{\mathbf{v}}_{1\perp}^2 \hat{\mathbf{v}}_{2\perp}^2}, \end{aligned} \quad (38)$$

with $\hat{\mathbf{v}}_{1\perp}$ and $\hat{\mathbf{v}}_{2\perp}$ obtained from $\mathbf{v}_{1\perp}$ and $\mathbf{v}_{2\perp}$ by setting $\mathbf{k}_{1\perp} = 0$; see Eq. (37). It is important to observe that, while the final state collinear divergence ($\hat{\mathbf{v}}_{1\perp} \rightarrow 0$) is absent, the hard factor has a divergence when the radiated gluon is collinear to the initial state proton ($\hat{\mathbf{v}}_{2\perp} \rightarrow 0$). This divergence is also present in Ref. [11], as can be seen from their Eq. (15) by setting the quark mass $m \rightarrow 0$.⁵ In hindsight, this means that the result in Ref. [11] must be incomplete in the sense that the lowest-order computation should be free from any divergences. That is, by taking into account also the $g_{1T}^{(1)}(x)$ part of the full cross section (3), as per the WW approximation (6), we indeed find that the initial state collinear divergence is canceled between the $g_T(x)$ and the $g_{1T}^{(1)}(x)$ parts, resulting in a finite hard factor (36). A similar conclusion was also reached in a collinear framework in SIDIS; see [30,37] where the $g_T(x)$ contribution to the cross section contained an initial state collinear divergence that gets exactly canceled with the collinear divergence in the $g_{1T}^{(1)}(x)$ part.

⁵One should be careful here in first factoring out one power of m in Eq. (15) in Ref. [11], as per the definition of $g_T(x)$.

B. Proof that the real contribution in the $q \rightarrow q$ channel vanishes

We now argue that, in fact, Eq. (35) is exactly zero. Before performing an explicit computation, we can appreciate it in an intuitive way as follows. In general, for the polarized cross section to be nonzero, we need two vectors: the transverse momentum of the final state and the spin so that we can form the familiar cross product $\mathbf{q}_\perp \times \mathbf{S}_\perp$. In the case of Eq. (35), we have \mathbf{q}_\perp , while we can think of $\mathbf{k}_{1\perp}$ as a proxy for the spin, thanks to the derivative $S_\perp^\lambda \partial / \partial k_{1\perp}^\lambda$. However, owing to the particular form of the hard factor (36), the two vectors \mathbf{q}_\perp and $\mathbf{k}_{1\perp}$ enter the cross section only through the linear combination $\mathbf{q}_{1\perp} \equiv \mathbf{q}_\perp - \bar{z}\mathbf{k}_{1\perp}$ (the soft part of the cross section is independent of $\mathbf{k}_{1\perp}$). Thus, the final result depends only on a single vector, $\mathbf{q}_{1\perp}$, and therefore must be zero.

To see the above statement explicitly, we start by switching to the coordinates

$$\begin{aligned} \mathbf{r}_\perp &= \mathbf{x}_\perp - \mathbf{y}_\perp, & \mathbf{b}_\perp &= \frac{\mathbf{x}_\perp + \mathbf{y}_\perp}{2}, \\ \mathbf{r}'_\perp &= \mathbf{y}_\perp - \mathbf{x}'_\perp, & \mathbf{b}'_\perp &= \frac{\mathbf{y}_\perp + \mathbf{x}'_\perp}{2}, \end{aligned} \quad (39)$$

to obtain

$$\begin{aligned} & \text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] \\ &= ig^2 N_c \frac{q^+}{P^+} \int_{\mathbf{k}_{2\perp}, \mathbf{k}_{1\perp}} \int_{\mathbf{r}_\perp, \mathbf{b}_\perp, \mathbf{r}'_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{r}_\perp} e^{i\mathbf{k}'_\perp \cdot \mathbf{r}'_\perp} \\ & \quad \times [\mathcal{P}(\mathbf{r}_\perp, \mathbf{b}_\perp) \mathcal{O}(\mathbf{r}'_\perp, \mathbf{b}'_\perp) - \mathcal{O}(\mathbf{r}_\perp, \mathbf{b}_\perp) \mathcal{P}(\mathbf{r}'_\perp, \mathbf{b}'_\perp)] \mathcal{H}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}). \end{aligned} \quad (40)$$

Note that not all transverse coordinates in Eq. (40) are independent—we have the following relation for \mathbf{b}'_\perp :

$$\mathbf{b}'_\perp = \mathbf{b}_\perp - \frac{1}{2}(\mathbf{r}_\perp + \mathbf{r}'_\perp). \quad (41)$$

This is an important point, because $\mathcal{O}(\mathbf{r}_\perp, -\mathbf{b}_\perp) = -\mathcal{O}(\mathbf{r}_\perp, \mathbf{b}_\perp)$ and so an integral over \mathbf{b}_\perp would superficially vanish simply via $\mathbf{b}_\perp \rightarrow -\mathbf{b}_\perp$. Next, in order to deconvolve the transverse integrals in Eq. (40), we Fourier transform the distributions as

$$\mathcal{P}(\mathbf{r}_\perp, \mathbf{b}_\perp) = \int_{\mathbf{k}_\perp, \mathbf{\Delta}_\perp} e^{-i\mathbf{k}_\perp \cdot \mathbf{r}_\perp} e^{-i\mathbf{\Delta}_\perp \cdot \mathbf{b}_\perp} \mathcal{P}(\mathbf{k}_\perp, \mathbf{\Delta}_\perp), \quad (42)$$

and similarly for $\mathcal{O}(\mathbf{r}_\perp, \mathbf{b}_\perp)$. In terms of the Fourier-transformed distributions, Eq. (40) becomes

$$\begin{aligned} \text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] &= ig^2 N_c \frac{q^+}{P^+} \int_{\mathbf{k}_\perp, \mathbf{\Delta}_\perp} [\mathcal{P}(\mathbf{k}_\perp, \mathbf{\Delta}_\perp) \mathcal{O}(\mathbf{k}'_\perp, \mathbf{\Delta}'_\perp) \\ & \quad - \mathcal{O}(\mathbf{k}_\perp, \mathbf{\Delta}_\perp) \mathcal{P}(\mathbf{k}'_\perp, \mathbf{\Delta}'_\perp)] \mathcal{H}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}), \end{aligned} \quad (43)$$

where $\mathbf{k}_\perp = \mathbf{k}_\perp + \frac{1}{2}\mathbf{\Delta}_\perp$, $\mathbf{k}_{2\perp} = \mathbf{k}'_\perp + \frac{1}{2}\mathbf{\Delta}_\perp$, and $\mathbf{\Delta}'_\perp = -\mathbf{\Delta}_\perp$.

The key quantity to consider in Eq. (43) is the integral over the angular variables. While the pomeron carries no angular dependence, the odderon has the following modulation $\mathcal{O}(\mathbf{k}_\perp, \mathbf{\Delta}_\perp) \propto (\mathbf{k}_\perp \cdot \mathbf{\Delta}_\perp)$ [this is simply the momentum space counterpart of the more familiar $\mathcal{O}(\mathbf{r}_\perp, \mathbf{b}_\perp) \sim (\mathbf{r}_\perp \cdot \mathbf{b}_\perp)$ modulation]; see, e.g., [52–54]. Focusing on the first part in Eq. (43), we start from the following expression:

$$\int_0^{2\pi} \frac{d\phi_\Delta}{2\pi} \int_0^{2\pi} \frac{d\phi_\kappa}{2\pi} \int_0^{2\pi} \frac{d\phi_{\kappa'}}{2\pi} (\mathbf{k}'_\perp \cdot \mathbf{\Delta}'_\perp) \frac{(\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp})}{v_{1\perp}^2 v_{2\perp}^2}. \quad (44)$$

Introducing $\delta_{1\perp} \equiv \mathbf{q}_{1\perp} - \bar{z}\mathbf{\Delta}_\perp/2$ and $\delta_{2\perp} \equiv \mathbf{q}_{1\perp} - \mathbf{\Delta}_\perp/2$, we have

$$\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp} = \delta_{1\perp} \times \delta_{2\perp} - \delta_{1\perp} \times \mathbf{k}_\perp - \bar{z}\mathbf{k}'_\perp \times \delta_{2\perp} + \bar{z}\mathbf{k}_\perp \times \mathbf{k}'_\perp \quad (45)$$

and

$$\begin{aligned} v_{1\perp}^2 &= \delta_{1\perp}^2 + \bar{z}^2 \mathbf{k}'_\perp{}^2 - 2\bar{z}\mathbf{k}'_\perp \cdot \delta_{1\perp} \cos(\phi_{\kappa'} - \phi_{\delta_1}), \\ v_{2\perp}^2 &= \delta_{2\perp}^2 + \mathbf{k}_\perp{}^2 - 2\mathbf{k}_\perp \cdot \delta_{2\perp} \cos(\phi_\kappa - \phi_{\delta_2}). \end{aligned} \quad (46)$$

Now we compute the integrals over ϕ_κ and $\phi_{\kappa'}$. From the first term in Eq. (45), we obtain

$$\begin{aligned} & \int_0^{2\pi} \frac{d\phi_\kappa}{2\pi} \int_0^{2\pi} \frac{d\phi_{\kappa'}}{2\pi} (\mathbf{k}'_\perp \cdot \mathbf{\Delta}'_\perp) \frac{(\delta_{1\perp} \times \delta_{2\perp})}{v_{1\perp}^2 v_{2\perp}^2} \\ &= -\frac{1}{4\bar{z}} (\delta_{1\perp} \cdot \mathbf{\Delta}_\perp) \frac{(\mathbf{q}_{1\perp} \times \mathbf{\Delta}_\perp)}{|\delta_{2\perp}^2 - \mathbf{k}_\perp{}^2|} \left(1 - \frac{\delta_{1\perp}^2 + \bar{z}^2 \mathbf{k}'_\perp{}^2}{|\delta_{1\perp}^2 - \bar{z}^2 \mathbf{k}'_\perp{}^2|} \right), \end{aligned} \quad (47)$$

where we used $\delta_{1\perp} \times \delta_{2\perp} = -z(\mathbf{q}_{1\perp} \times \mathbf{\Delta}_\perp)/2$. Inserting now the definition of $\delta_{1\perp}$, we will, in general, have an expression of the type

$$\begin{aligned} & \int_0^{2\pi} \frac{d\phi_\Delta}{2\pi} \sin(\phi_{q_1} - \phi_\Delta) f(\cos(\phi_{q_1} - \phi_\Delta)) \\ &= - \int_{\phi_{q_1}}^{\phi_{q_1} - 2\pi} \frac{d\phi}{2\pi} \sin \phi f(\cos \phi), \end{aligned} \quad (48)$$

but this is simply zero, as

$$\begin{aligned} \int_{\phi_1}^{\phi_1 - 2\pi} d\phi \sin \phi f(\cos \phi) &= - \int_{\phi_1}^{\phi_1 - 2\pi} d(\cos \phi) f(\cos \phi) \\ &= F(\cos \phi) \Big|_{\phi_1}^{\phi_1 - 2\pi} = 0, \end{aligned} \quad (49)$$

where $F(\cos \phi)$ is a primitive function of $f(\cos \phi)$. By a completely analogous computation, we can show that each of the remaining three pieces in Eq. (45) is also zero and, thus, conclude that the complete real contribution in the $q \rightarrow q$ channel vanishes.

C. $q \rightarrow q$: Virtual contribution

For the virtual correction to the $q \rightarrow q$ channel, we have the following amplitude:

$$\begin{aligned} \mathcal{M} = & g^2 \int_{-\infty}^{\infty} \frac{dk_g^+}{(2\pi)} \int_{k_{g\perp}, k_{1\perp}} \int_{x_{\perp}, y_{\perp}} e^{ik_{1\perp} \cdot x_{\perp}} e^{i(q_{\perp} - k_{1\perp} - k_{\perp}) \cdot y_{\perp}} \\ & \times [t^a t^a V(\mathbf{x}_{\perp}) \mathcal{T}_{q,1} + V(\mathbf{x}_{\perp}) t^a t^a \mathcal{T}_{q,2} \\ & + t^a V(\mathbf{x}_{\perp}) t^b U^{ab}(\mathbf{y}_{\perp}) \mathcal{T}_{qg}], \end{aligned} \quad (50)$$

where

$$\begin{aligned} \mathcal{T}_{q,1} = & i \int_{-\infty}^{\infty} \frac{dk_g^-}{(2\pi)} \gamma^{\mu} \frac{\not{q} - \not{k}_g}{(q - k_g)^2 + i\epsilon} \gamma^{\nu} \frac{\not{q}}{q^2 + i\epsilon} \gamma^{\rho} \frac{d_{\mu\nu}(k_g)}{k_g^2 + i\epsilon}, \\ \mathcal{T}_{q,2} = & i \int_{-\infty}^{\infty} \frac{dk_g^-}{(2\pi)} \gamma^{\rho} \frac{\not{k}_1}{k_1^2 + i\epsilon} \gamma^{\mu} \frac{\not{k}_1 - \not{k}_g}{(k_1 - k_g)^2 + i\epsilon} \gamma^{\nu} \frac{d_{\mu\nu}(k_g)}{k_g^2 + i\epsilon}, \\ \mathcal{T}_{qg} = & i \int_{-\infty}^{\infty} \frac{dk_g^-}{(2\pi)} \gamma^{\mu} \frac{\not{q} - \not{k}_g}{(q - k_g)^2 + i\epsilon} \gamma^{\rho} \frac{\not{q} - \not{k} - \not{k}_g}{(q - k - k_g)^2 + i\epsilon} \\ & \times \gamma_{\rho} d^{\nu\mu}(k + k_1 + k_g - q) \frac{d_{\mu\nu}(k_g)}{k_g^2 + i\epsilon}. \end{aligned} \quad (51)$$

Above, in the first line of \mathcal{T}_{qg} we have evaluated the k^- integration in favor of the singularity at $(k_1 + k + k_g - q)^2 + i\epsilon = 0$ so that

$$\begin{aligned} S^{(0)}(k_1) = & (2\pi) \delta(k_1^+ - q^+) C_F g^2 \frac{1}{2P_p^+} \int_{-\infty}^{\infty} \frac{dk_g^+}{(2\pi)} \int_{k_{g\perp}, k_{1\perp}, k'_{1\perp}} \int_{x_{\perp}, x'_{\perp}, y_{\perp}, y'_{\perp}} e^{ik_{1\perp} \cdot x_{\perp}} e^{i(k_{2\perp} - k_{1\perp}) \cdot y_{\perp}} e^{-ik'_{1\perp} \cdot x'_{\perp}} e^{-i(k_{2\perp} - k'_{1\perp}) \cdot y'_{\perp}} \\ & \times [\mathcal{S}_q(x'_{\perp}, \mathbf{x}_{\perp}) \gamma^+ \not{q} \mathcal{T}_q + \mathcal{S}_q(x'_{\perp}, \mathbf{x}_{\perp}) \bar{\mathcal{T}}_q \not{q} \gamma^+ + \mathcal{S}_{qg}(x'_{\perp}, \mathbf{x}_{\perp}, \mathbf{y}_{\perp}) \gamma^+ \not{q} \mathcal{T}_{qg}(k_{\perp}) + \mathcal{S}_{qg}(x'_{\perp}, \mathbf{x}_{\perp}, \mathbf{y}'_{\perp}) \bar{\mathcal{T}}_{qg}(k'_{\perp}) \not{q} \gamma^+], \end{aligned} \quad (53)$$

where now $\mathbf{k}_{2\perp} \equiv \mathbf{q}_{\perp} - \mathbf{k}_{1\perp}$ and $\mathcal{T}_q \equiv \mathcal{T}_{q,1} + \mathcal{T}_{q,2}$. Analogous to the case of real production, the terms in Eq. (53) that are proportional to the dipole operator will not contribute as a consequence of C parity. This includes the first two terms of the second line and the dipole pieces in the last two terms of the second line according to Eq. (32). Repeating further the steps of the calculation used for real production, we find

$$\begin{aligned} \text{tr}[\gamma_5 \not{k}_1 S^{(0)}(k_1)] = & i(2\pi) \delta(k_1^+ - q^+) N_c g^2 \frac{q^+}{P_p^+} \int_{-\infty}^{\infty} \frac{dk_g^+}{(2\pi)} \\ & \times \int_{k_{g\perp}, k_{1\perp}} \int_{x_{\perp}, x'_{\perp}, y_{\perp}} e^{ik_{1\perp} \cdot (x_{\perp} - y_{\perp})} e^{-ik_{2\perp} \cdot (x'_{\perp} - y_{\perp})} \\ & \times [\mathcal{P}(x_{\perp}, y_{\perp}) \mathcal{O}(x'_{\perp}, y_{\perp}) - \mathcal{O}(x_{\perp}, y_{\perp}) \\ & \times \mathcal{P}(x'_{\perp}, y_{\perp})] \mathcal{H}(k_{\perp}, k_{1\perp}). \end{aligned} \quad (54)$$

See Fig. 2 for the diagram corresponding to this remaining contribution. Here, now $\mathcal{H}(k_{\perp}, k_{1\perp})$ is

$$\begin{aligned} \mathcal{H}(k_{\perp}, k_{1\perp}) = & \frac{1}{2q^+} \text{tr}[\gamma_5 \not{k}_1 \gamma^+ \not{q} \mathcal{T}_{qg}(k_{\perp})] \\ = & -4i(\bar{y} + 1) \frac{\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp}}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2}, \end{aligned} \quad (55)$$




FIG. 2. An interference diagram that determines $S^{(0)}(k_1)$ in the virtual correction to the $q \rightarrow q$ channel. The vertical gluons denote Wilson lines arising from multiple scattering on the dense nucleus.

$$\begin{aligned} (q - k - k_g)^2 = & -\frac{1}{k_1^+ k_g^+} [k_1^+ (\mathbf{k}_{\perp} + \mathbf{k}_{1\perp} + \mathbf{k}_{g\perp} - \mathbf{q}_{\perp}) \\ & - k_g^+ \mathbf{k}_{1\perp}]^2. \end{aligned} \quad (52)$$

To get the NLO virtual contribution to $S^{(0)}(k_1)$, we combine the virtual amplitude (50) with the LO amplitude (9) and find

where we have introduced $y \equiv k_g^+ / k_1^+ = k_g^+ / q^+ = z / \bar{z}$ and $\mathbf{v}_{1\perp}$ ($\mathbf{v}_{2\perp}$) are associated with final (initial) state collinear configurations explicitly given as $\mathbf{v}_{1\perp} \equiv y \mathbf{q}_{\perp} - \mathbf{k}_{g\perp}$ and $\mathbf{v}_{2\perp} \equiv \mathbf{k}_{\perp} + \bar{y} \mathbf{k}_{1\perp} + \mathbf{k}_{g\perp} - \mathbf{q}_{\perp}$. To compute Eq. (55), we have evaluated the k_g^- integral in $\mathcal{T}_{qg}(k_{\perp})$ in favor of the singularity $k_g^2 + i\epsilon = 0$. Proceeding with the $\mathbf{k}_{g\perp}$ loop integral, we pass from the variable $\mathbf{k}_{g\perp}$ to $\mathbf{v}_{2\perp}$ and write

$$\int_{\mathbf{k}_{g\perp}} \frac{\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp}}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2} = - \int_{\mathbf{v}_{2\perp}} \frac{\mathbf{v}_{\perp} \times \mathbf{v}_{2\perp}}{(\mathbf{v}_{\perp} + \mathbf{v}_{2\perp})^2 \mathbf{v}_{2\perp}^2}, \quad (56)$$

where $\mathbf{v}_{\perp} \equiv \bar{y} \mathbf{q}_{\perp} - y \mathbf{k}_{1\perp} - \mathbf{k}_{\perp}$. But this contains an angular integral that is precisely of the form (49) and, therefore, vanishes.

D. $q \rightarrow g$

In this case, we have only the real diagram with gluon fragmenting into a final state hadron. The expression for $S^{(0)}(k_1)$ takes the same form as Eq. (22), with the only difference being that now we are integrating over \mathbf{q}_{\perp} (the momenta of the untagged quark) instead of over $\mathbf{k}_{g\perp}$:

$$\begin{aligned}
S^{(0)}(k_1) &= \frac{k_g^+}{P_p^+} \frac{g^2 C_F}{(2q^+)(2k_g^+)} \int_{\mathbf{q}_\perp \mathbf{k}_\perp \mathbf{k}'_\perp} \int_{\mathbf{x}_\perp \mathbf{x}'_\perp \mathbf{y}_\perp \mathbf{y}'_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} e^{-i\mathbf{k}'_\perp \cdot \mathbf{x}'_\perp} e^{-i(\mathbf{k}_{2\perp} - \mathbf{k}'_\perp) \cdot \mathbf{y}'_\perp} d_{\mu\mu'}(k_g) [\mathcal{S}(\mathbf{x}_\perp, \mathbf{x}'_\perp) \bar{T}'_{q'} \not{q} T_q^\mu \\
&\quad + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}'_\perp) \bar{T}'_{qg}(\mathbf{k}'_\perp) \not{q} T_q^\mu + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp) \bar{T}'_{q'} \not{q} T_{qg}^\mu(\mathbf{k}_\perp) + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{y}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp) \bar{T}'_{qg}(\mathbf{k}'_\perp) \not{q} T_{qg}^\mu(\mathbf{k}_\perp)].
\end{aligned} \tag{57}$$

The first term again vanishes due to C parity. To show that the last term vanishes, we first need to make the following replacements: $\mathbf{k}_\perp - \mathbf{q}_\perp \rightarrow \mathbf{k}_\perp$ and $\mathbf{k}'_\perp - \mathbf{q}_\perp \rightarrow \mathbf{k}'_\perp$ for the \mathbf{k}_\perp and \mathbf{k}'_\perp integrals, which makes $T_{gq}(\mathbf{k}_\perp)$ independent of \mathbf{q}_\perp . Additionally, since \not{q} is sandwiched between two γ^+ matrices, it does not give a \mathbf{q}_\perp contribution. Therefore, the respective hard factor does not depend on \mathbf{q}_\perp , and the only \mathbf{q}_\perp dependence in the last term appears in the exponential. From here, we take the analogous steps as in the $q \rightarrow q$ channel. First performing the \mathbf{q}_\perp integration

$$\int_{\mathbf{q}_\perp} e^{i\mathbf{q}_\perp \cdot (\mathbf{x}_\perp - \mathbf{x}'_\perp)} = \delta^{(2)}(\mathbf{x}_\perp - \mathbf{x}'_\perp), \tag{58}$$

$S_{qqg}(\mathbf{x}'_\perp, \mathbf{y}'_\perp, \mathbf{x}_\perp, \mathbf{y}_\perp)$ collapses to an adjoint dipole [see Eq. (23)] when $\mathbf{x}'_\perp = \mathbf{x}_\perp$, which is, in addition, independent of \mathbf{x}_\perp . This allows us to perform the \mathbf{x}_\perp integral to conclude that $\mathbf{k}'_\perp = \mathbf{k}_\perp$. With this, we can utilize C parity to find the hard factor from the last term vanishes. For the remaining interference terms, the trace is given in Eq. (36). With the vectors $\mathbf{v}_{1\perp}$ and $\mathbf{v}_{2\perp}$ suitably rewritten in the form $\mathbf{v}_{1\perp} = (z\mathbf{k}_{1\perp} - \mathbf{k}_{g\perp}) + \mathbf{k}_{2\perp}$ and $\mathbf{v}_{2\perp} = (z\mathbf{k}_{1\perp} - \mathbf{k}_{g\perp}) + \mathbf{k}_{2\perp} - \mathbf{k}_\perp$, we see that a unique combination $z\mathbf{k}_{1\perp} - \mathbf{k}_{g\perp}$ appears. Thus, the steps to show that the cross section also vanishes in the $q \rightarrow g$ channel are from this point completely analogous to those for the $q \rightarrow q$ channel; see Sec. III B. Together with the result from Secs. III B and III C, this completes the statement that in the $q \rightarrow qg$ channel the

odderon mechanism in the WW approximation does not contribute to SSA at NLO.

IV. THE $g \rightarrow q\bar{q}$ CHANNEL

In this channel, we label the momenta as $g(k_1) \rightarrow q(q)\bar{q}(p)$. The NLO amplitude can be written as

$$\begin{aligned}
\mathcal{M}^\alpha &= (+g) \int_{\mathbf{k}_\perp} \int_{\mathbf{x}_\perp \mathbf{y}_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} [T_g^\alpha t^b U^{ba}(\mathbf{x}_\perp) \\
&\quad + T_{q\bar{q}}^\alpha(\mathbf{k}_\perp) V(\mathbf{x}_\perp) t^a V^\dagger(\mathbf{y}_\perp)],
\end{aligned} \tag{59}$$

where

$$T_g^\alpha \equiv 2k_1^+ \gamma_p \frac{d^{p\alpha}(q+p)}{(q+p)^2} \tag{60}$$

and

$$T_{q\bar{q}}^\alpha(\mathbf{k}_\perp) \equiv \frac{1}{2p^+} \gamma^+ \frac{\not{q} - \not{k}}{(q-k)^2} \gamma^\alpha (\not{q} - \not{k} - \not{k}_1) \gamma^+. \tag{61}$$

In the above expressions, similar to the discussion in Sec. III A, k^- is obtained by picking up the pole from the condition $(q-k-k_1)^2 + i\epsilon = 0$. The above results can be shown to agree with the corresponding amplitude in Eq. (38) in Ref. [55] used for unpolarized pA collisions after taking the collinear limit for the gluon from the proton. Using Eq. (59), we calculate $S_{\alpha\beta}^{(0)}(k_1)$ as

$$\begin{aligned}
S_{\alpha\beta}^{(0)}(k_1) &= \frac{1}{2P_p^+} \int \frac{d^3p}{(2\pi)^3 2E_p} \frac{1}{N_c^2 - 1} \text{Tr} \langle \bar{\mathcal{M}}_\alpha \not{q} \mathcal{M}_\beta \not{p} \rangle \\
&= \frac{q^+}{P_p^+} \frac{g^2 T_R}{(2q^+)(2p^+)} \int_{\mathbf{p}_\perp \mathbf{k}_\perp \mathbf{k}'_\perp} \int_{\mathbf{x}_\perp \mathbf{x}'_\perp \mathbf{y}_\perp \mathbf{y}'_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} e^{-i\mathbf{k}'_\perp \cdot \mathbf{x}'_\perp} e^{-i(\mathbf{k}_{2\perp} - \mathbf{k}'_\perp) \cdot \mathbf{y}'_\perp} \\
&\quad \times \{ \mathcal{S}_A(\mathbf{x}_\perp, \mathbf{x}'_\perp) \text{tr}[\bar{T}'_{g,\alpha} \not{q} T_{g,\beta} \not{p}] + \mathcal{S}_{qqg}(\mathbf{x}'_\perp, \mathbf{y}'_\perp, \mathbf{x}_\perp) \text{tr}[\bar{T}'_{q\bar{q},\alpha}(\mathbf{k}'_\perp) \not{q} T_{g,\beta} \not{p}] \\
&\quad + \mathcal{S}_{qqg}(\mathbf{y}_\perp, \mathbf{x}_\perp, \mathbf{x}'_\perp) \text{tr}[\bar{T}'_{g,\alpha} \not{q} T_{q\bar{q},\beta}(\mathbf{k}_\perp) \not{p}] + \mathcal{S}_{qqq}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}'_\perp, \mathbf{y}_\perp) \text{tr}[\bar{T}'_{q\bar{q},\alpha}(\mathbf{k}'_\perp) \not{q} T_{q\bar{q},\beta}(\mathbf{k}_\perp) \not{p}] \},
\end{aligned} \tag{62}$$

where $T_R = 1/2$. Here, \mathcal{S}_A and \mathcal{S}_{qqg} are defined in Eq. (15) and the first line of Eq. (23), respectively, while

$$\begin{aligned}
\mathcal{S}_{qqq}(\mathbf{x}'_\perp, \mathbf{x}_\perp, \mathbf{y}'_\perp, \mathbf{y}_\perp) \\
\equiv \frac{1}{C_F N_c} \text{tr} \langle V^\dagger(\mathbf{x}_\perp) V(\mathbf{x}_\perp) t^a V^\dagger(\mathbf{y}_\perp) V(\mathbf{y}_\perp) t^a \rangle
\end{aligned} \tag{63}$$

is an additional gluon distribution of the target. Similar to the findings in Sec. III A, the first and the fourth terms in Eq. (62) vanish. For the first term, this can be argued from C parity on the Dirac trace [another way is simply from the fact that the adjoint dipole $\mathcal{S}_A(\mathbf{x}_\perp, \mathbf{x}'_\perp)$ is real]. For the fourth term, the key point is that the hard factor does not depend on \mathbf{p}_\perp (\not{p} is sandwiched between γ^+ , and so the \mathbf{p}_\perp




FIG. 3. An interference diagram that determines $S_{\alpha\beta}^{(0)}(k_1)$ in the $g \rightarrow q\bar{q}$ channel. The vertical gluons denote Wilson lines arising from multiple scattering on the dense nucleus.

dependence drops out). Then, the fourth term does not contribute by the same steps used in Sec. III A. This leaves the interference term in Eq. (62) that is represented graphically in Fig. 3. According to the WW truncation in Eq. (7), these require the evaluation of the following hard factors:

$$\begin{aligned} \mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) &\equiv \frac{1}{4q^+p^+} \frac{1}{p^+} \epsilon^{+\alpha\beta S_\perp} \omega_{\alpha'\alpha} \omega_{\beta'\beta} \\ &\quad \times \text{tr}[\bar{T}_g^{\alpha'} \not{q} T_{q\bar{q}}^{\beta'}(\mathbf{k}_\perp) \not{p}], \\ \mathcal{H}^{(\bar{g})}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) &\equiv \frac{1}{4q^+p^+} (g_{\beta\lambda}^{\beta\lambda} \epsilon^{\alpha-+S_\perp} - g_{\beta\lambda}^{\alpha\lambda} \epsilon^{\beta-+S_\perp}) \\ &\quad \times \text{tr}[\bar{T}_{g,\alpha} \not{q} T_{q\bar{q},\beta}(\mathbf{k}_\perp) \not{p}]. \end{aligned} \quad (64)$$

We find

$$\mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) = 4z\bar{z}(z - \bar{z}) \frac{\hat{\mathbf{v}}_{1\perp} \times \mathbf{S}_\perp}{\hat{\mathbf{v}}_{1\perp}^2 \hat{\mathbf{v}}_{2\perp}^2}, \quad (65)$$

$$\begin{aligned} \mathcal{H}^{(\bar{g})}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) &= \frac{4}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2} \{ -[(z^2 + \bar{z}^2)(\mathbf{v}_{2\perp} \times \mathbf{S}_\perp) \\ &\quad + z\bar{z}(z - \bar{z})(\mathbf{k}_{1\perp} \times \mathbf{S}_\perp)] v_{1\perp}^\lambda \\ &\quad + (z^2 + \bar{z}^2)(\mathbf{v}_{1\perp} \times \mathbf{S}_\perp) v_{2\perp}^\lambda \\ &\quad + z\bar{z}(z - \bar{z})(\mathbf{v}_{1\perp} \times \mathbf{S}_\perp) k_{1\perp}^\lambda \}, \end{aligned} \quad (66)$$

where now $\mathbf{v}_{1\perp} \equiv z\mathbf{q}_\perp - \bar{z}\mathbf{p}_\perp = \mathbf{q}_\perp - \bar{z}\mathbf{k}_{1\perp} - \bar{z}\mathbf{k}_{2\perp}$ and $\mathbf{v}_{2\perp} \equiv \mathbf{q}_\perp - \bar{z}\mathbf{k}_{1\perp} - \mathbf{k}_\perp$, with $z \equiv p^+/k_1^+$ the momentum fraction of the recoiling antiquark. By $\hat{\mathbf{v}}_{1\perp}$ ($\hat{\mathbf{v}}_{2\perp}$) in Eq. (65), we again denote $\mathbf{v}_{1\perp}$ ($\mathbf{v}_{2\perp}$) at $\mathbf{k}_{1\perp} = 0$. According to the WW truncation of the polarized cross section, we also need to take a derivative of $\mathcal{H}^{(\bar{g})}(\mathbf{k}_\perp, \mathbf{k}_{1\perp})$ with respect to $k_{1\lambda}$ (and sum over λ); cf. second line in Eq. (7). After this, we could proceed with the angular integrals as in Sec. III B, but this time the expressions would involve \mathbf{S}_\perp . A simpler way to proceed is to first combine the first and the second line in Eq. (7), leading to

$$\begin{aligned} &\mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) - \left[\frac{\partial}{\partial k_1^\lambda} \mathcal{H}^{(\bar{g})}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) \right]_{k_1=p_1} \\ &= -\frac{4\bar{z}(z^2 + \bar{z}^2)}{\hat{\mathbf{v}}_{1\perp}^4 \hat{\mathbf{v}}_{2\perp}^4} [(\hat{\mathbf{v}}_{1\perp}^2 \hat{\mathbf{v}}_{2\perp}^2 + 2(\hat{\mathbf{v}}_{1\perp} \cdot \hat{\mathbf{v}}_{2\perp}) \mathbf{v}_{2\perp}^2)(\hat{\mathbf{v}}_{1\perp} \times \mathbf{S}_\perp) \\ &\quad - (\hat{\mathbf{v}}_{1\perp}^2 \hat{\mathbf{v}}_{2\perp}^2 + 2(\hat{\mathbf{v}}_{1\perp} \cdot \hat{\mathbf{v}}_{2\perp}) \hat{\mathbf{v}}_{1\perp}^2)(\hat{\mathbf{v}}_{2\perp} \times \mathbf{S}_\perp)]. \end{aligned} \quad (67)$$

After some inspection, this can be also rewritten in a more convenient form as

$$\begin{aligned} &\mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) - \left[\frac{\partial}{\partial k_1^\lambda} \mathcal{H}^{(\bar{g})}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) \right]_{k_1=p_1} \\ &= \left[S_\perp^\lambda \frac{\partial}{\partial k_{1\perp}^\lambda} \left(4(z^2 + \bar{z}^2) \frac{\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp}}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2} \right) \right]_{k_1=p_1}, \end{aligned} \quad (68)$$

where S_\perp^λ is now factored out and the effective hard factor inside the brackets in Eq. (68) now has finite $\mathbf{k}_{1\perp}$ through $\mathbf{v}_{1\perp}$ and $\mathbf{v}_{2\perp}$. To prove the equivalence of Eqs. (67) and (68), we have used the Schouten identity $\hat{\mathbf{v}}_{1\perp}(\hat{\mathbf{v}}_{2\perp} \times \mathbf{S}_\perp) + \hat{\mathbf{v}}_{2\perp}(\mathbf{S}_\perp \times \hat{\mathbf{v}}_{1\perp}) + \mathbf{S}_\perp(\hat{\mathbf{v}}_{1\perp} \times \hat{\mathbf{v}}_{2\perp}) = 0$. Equation (68) reveals that the general structure of the $g \rightarrow q$ hard factor is the same as in the $q \rightarrow q$ channel; see Eq. (36). Therefore, by following the same logic as in Sec. III B, we conclude that the corresponding polarized cross section in the $g \rightarrow q\bar{q}$ channel also vanishes.

V. THE $g \rightarrow gg$ CHANNEL

In the case of the $g \rightarrow gg$ channel, there is a great simplification due to the fact that the purely gluonic contributions involve only adjoint Wilson lines, which are real, and, therefore, the odderon mechanism is absent. The only exception is the quark loop correction to the tree-level $g(k_1) \rightarrow g(k_g)$ amplitude (see Fig. 4), which we compute below. Discarding immediately the dipole pieces, $S_{\alpha\beta}^{(0)}(k_1)$ takes the following form:




FIG. 4. An interference diagram that determines $S_{\alpha\beta}^{(0)}(k_1)$ in the virtual correction to the $g \rightarrow g$ channel. The vertical gluons denote Wilson lines arising from multiple scattering on the dense nucleus.

$$\begin{aligned}
& S_{\alpha\beta}^{(0)}(k_1) \\
&= \frac{1}{2P_p^+} (2\pi) \delta(k_1^+ - k_g^+) (-ig^2) N_f N_c T_R \int_{-\infty}^{\infty} \frac{dq^+}{(2\pi)} \int_{\mathbf{q}_\perp, \mathbf{k}_\perp, \mathbf{k}'_\perp} \\
&\times \int_{\mathbf{x}_\perp, \mathbf{x}'_\perp, \mathbf{y}_\perp, \mathbf{y}'_\perp} e^{i\mathbf{k}_\perp \cdot \mathbf{x}_\perp} e^{i(\mathbf{k}_{2\perp} - \mathbf{k}_\perp) \cdot \mathbf{y}_\perp} e^{-i\mathbf{k}_\perp \cdot \mathbf{x}'_\perp} e^{-i(\mathbf{k}_{2\perp} - \mathbf{k}'_\perp) \cdot \mathbf{y}'_\perp} \\
&\times [S_{qqg}(\mathbf{x}_\perp, \mathbf{y}_\perp, \mathbf{x}'_\perp) (-2k_1^+) d_{\alpha\mu}(k_g) T_{q\bar{q},\beta}^\mu(\mathbf{k}_\perp) \\
&+ S_{qqg}(\mathbf{y}'_\perp, \mathbf{x}'_\perp, \mathbf{x}_\perp) (-2k_1^+) d_{\alpha\mu}(k_g) T_{q\bar{q},\beta}^{\mu\dagger}(\mathbf{k}'_\perp)], \quad (69)
\end{aligned}$$

where q is the quark loop momentum and

$$\begin{aligned}
T_{q\bar{q}}^{\mu\beta}(\mathbf{k}_\perp) &\equiv \frac{1}{2q^+} \int_{-\infty}^{\infty} \frac{dq^-}{(2\pi)} \text{tr} \left[\frac{\not{q} - \not{k}_g}{(q - k_g)^2 + i\epsilon} \gamma^\mu \frac{\not{q}}{q^2 + i\epsilon} \right. \\
&\times \left. \gamma^+ (\not{q} - \not{k}_g + \not{k}_1 + \not{k}) \gamma^\beta \frac{\not{q} - \not{k}_g + \not{k}}{(q - k_g + k)^2} \gamma^+ \right]. \quad (70)
\end{aligned}$$

Similar to the computation in Sec. IV, and according to Eq. (7), we are to evaluate the following combination:

$$\mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) - \left[\frac{\partial}{\partial k_1^\lambda} \mathcal{H}^{(\bar{g})\lambda}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) \right]_{k_1=p_1}, \quad (71)$$

where we define

$$\begin{aligned}
\mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) &\equiv \frac{1}{(2q^+)^2} \frac{1}{p_1^+} e^{n\alpha\beta S_\perp} \omega_{\alpha'\alpha} \omega_{\beta'\beta} \\
&\times (-2k_1^+) d_{\alpha\mu}(k_g) T_{q\bar{q},\beta}^\mu(\mathbf{k}_\perp), \\
\mathcal{H}^{(\bar{g})\lambda}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) &\equiv \frac{1}{(2q^+)^2} (g_\perp^{\beta\lambda} e^{a\bar{n}n S_\perp} - g_\perp^{\alpha\lambda} e^{\beta\bar{n}n S_\perp}) \\
&\times (-2k_1^+) d_{\alpha\mu}(k_g) T_{q\bar{q},\beta}^\mu(\mathbf{k}_\perp). \quad (72)
\end{aligned}$$

A direct computation leads to

$$\begin{aligned}
& \mathcal{H}^{(G_{3T})}(\mathbf{k}_\perp) - \left[\frac{\partial}{\partial k_1^\lambda} \mathcal{H}^{(\bar{g})\lambda}(\mathbf{k}_\perp, \mathbf{k}_{1\perp}) \right]_{k_1=p_1} \\
&= \left[S_\perp^\lambda \frac{\partial}{\partial k_{1\perp}^\lambda} \left(4(y^2 + \bar{y}^2) \frac{\mathbf{v}_{1\perp} \times \mathbf{v}_{2\perp}}{\mathbf{v}_{1\perp}^2 \mathbf{v}_{2\perp}^2} \right) \right]_{k_1=p_1}, \quad (73)
\end{aligned}$$

where now $\mathbf{v}_{1\perp} \equiv -\mathbf{q}_\perp + \mathbf{k}_{g\perp} - \bar{y}\mathbf{k}_{1\perp} - \mathbf{k}_\perp$ and $\mathbf{v}_{2\perp} \equiv \mathbf{q}_\perp - y\mathbf{k}_{g\perp}$. But this is completely analogous to the result for the loop correction in Sec. III C, and so $S_{\alpha\beta}^{(0)}(k_1)$ vanishes after the \mathbf{q}_\perp integral.

VI. DISCUSSION AND CONCLUSIONS

We have revisited the odderon mechanism for SSA in $p^\uparrow A$ originally suggested in Ref. [11] at the quark level.

At the hadron level, this mechanism would involve the $g_T(x)$ distribution. We have considered the WW truncation of the full twist-3 polarized cross section and argued that, in addition to $g_T(x)$, we also need to take into account the $g_{1T}^{(1)}(x)$ for a consistent computation. Our main finding is that under this truncation the polarized cross section vanishes exactly up to NLO for all possible partonic channels.

It is natural to consider whether any of the above assumptions can be relaxed so that a nonzero contribution to SSA from the odderon mechanism may be found after all. One option is to go beyond the WW approximation, namely, including the ETQS pieces in Eq. (6). Already at the LO, this could potentially yield a new contribution where the phase is obtained from the odderon. Note the difference from the more conventional pole calculus—here, one needs to pick up the principal value of internal propagators so that the general functional forms of the ETQS functions would be required. Going beyond the WW approximation at the NLO is more challenging. In practice, restricting merely to $S^{(0)}(k_1)$ and its $k_{1\perp}$ derivative is not enough, as one needs to take into account also the full three-body kernel $S_\lambda^{(1)}(x_p P_p, x'_p P_p)$ that includes pole and nonpole pieces. Already the collinear divergence found in the hard factor associated with $g_T(x)$ [see Eq. (38)] remains beyond the WW approximation, underlying the need to compute $S_\lambda^{(1)}(x_p P_p, x'_p P_p)$ in order to get a finite cross section.⁶ One can also consider the twist-3 FF mechanism, where we would pick up the real part of the twist-3 FFs with the phase provided by the odderon. Once again, this in contrast to the conventional computations where the phase is supplied by the imaginary part of twist-3 FFs. Given that the current global fits constrain only the imaginary part of the twist-3 FFs [57,58], the phenomenological implications of this alternative would be worth exploring.

Another possibility would be to retain the WW approximation but compute the hard factor up to NNLO. While, of course, only an explicit computation can reveal whether the odderon appears at NNLO, we mention here a competing mechanism that is already known to appear at NNLO. The basic premise is very simple: At higher orders, it is an imaginary part of the loop amplitude that can supply the phase. A specific NNLO contribution illustrating this is given in Fig. 5, where the crosses denote cut propagators. Physically, the initial $q \rightarrow qg$ splitting occurs inside the target nucleus in the amplitude. The qg system subsequently rescatters with a t -channel quark into the final state, providing a phase with respect to the amplitude on the opposite side of the final state cut. Such final state rescattering is sometimes referred to as the lensing

⁶It is worth mentioning here the computation of target SSA in deep inelastic scattering, where it was found that the divergent pieces cancel among the two-body and three-body kernels [56].




FIG. 5. A sample interference diagram appearing at NNLO. The crosses denote cut propagators that determine the imaginary part of the loop.

mechanism and was considered in Ref. [19]. In fact, this idea [59] is closely related to the very first estimate of SSA in perturbative QCD [60]. The computation in Ref. [19] was in the quark-diquark model. As a future work, it would be important to consider this in the hybrid approach.

ACKNOWLEDGMENTS

S. B. thanks Yoshitaka Hatta for suggesting to work on the odderon mechanism for SSA. We thank Yoshitaka Hatta and Yuri Kovchegov for useful comments on the manuscript. S. B., A. K., and E. A. V. are supported by the Croatian Science Foundation (HRZZ) No. 5332 (UIP-2019-04).

- [1] V. Barone, A. Drago, and P. G. Ratcliffe, *Phys. Rep.* **359**, 1 (2002).
- [2] U. D’Alesio and F. Murgia, *Prog. Part. Nucl. Phys.* **61**, 394 (2008).
- [3] D. Pitonyak, *Int. J. Mod. Phys. A* **31**, 1630049 (2016).
- [4] M. Grosse Perdekamp and F. Yuan, *Annu. Rev. Nucl. Part. Sci.* **65**, 429 (2015).
- [5] C. Aidala *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **123**, 122001 (2019).
- [6] J. Adam *et al.* (STAR Collaboration), *Phys. Rev. D* **103**, 072005 (2021).
- [7] E. Iancu and R. Venugopalan, The color glass condensate and high-energy scattering in QCD, in *Quark-Gluon Plasma 3*, edited by R. C. Hwa and X.-N. Wang (World Scientific, Singapore, 2003), pp. 249–3363.
- [8] F. Gelis, E. Iancu, J. Jalilian-Marian, and R. Venugopalan, *Annu. Rev. Nucl. Part. Sci.* **60**, 463 (2010).
- [9] Y. V. Kovchegov and E. Levin, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **33**, 1 (2012).
- [10] J.-P. Blaizot, *Rep. Prog. Phys.* **80**, 032301 (2017).
- [11] Y. V. Kovchegov and M. D. Sievert, *Phys. Rev. D* **86**, 034028 (2012); **86**, 079906(E) (2012).
- [12] D. Boer, A. Dumitru, and A. Hayashigaki, *Phys. Rev. D* **74**, 074018 (2006).
- [13] Z.-B. Kang and F. Yuan, *Phys. Rev. D* **84**, 034019 (2011).
- [14] A. Schäfer and J. Zhou, *Phys. Rev. D* **90**, 034016 (2014).
- [15] J. Zhou, *Phys. Rev. D* **92**, 014034 (2015).
- [16] Y. Hatta, B.-W. Xiao, S. Yoshida, and F. Yuan, *Phys. Rev. D* **94**, 054013 (2016).
- [17] Y. Hatta, B.-W. Xiao, S. Yoshida, and F. Yuan, *Phys. Rev. D* **95**, 014008 (2017).
- [18] S. Benić and Y. Hatta, *Phys. Rev. D* **99**, 094012 (2019).
- [19] Y. V. Kovchegov and M. G. Santiago, *Phys. Rev. D* **102**, 014022 (2020).
- [20] Y. V. Kovchegov, L. Szymanowski, and S. Wallon, *Phys. Lett. B* **586**, 267 (2004).
- [21] Y. Hatta, E. Iancu, K. Itakura, and L. McLerran, *Nucl. Phys. A* **760**, 172 (2005).
- [22] S. Jeon and R. Venugopalan, *Phys. Rev. D* **71**, 125003 (2005).
- [23] A. Bacchetta, D. Boer, M. Diehl, and P. J. Mulders, *J. High Energy Phys.* **08** (2008) 023.
- [24] A. V. Efremov and O. V. Teryaev, *Sov. J. Nucl. Phys.* **36**, 140 (1982).
- [25] J.-w. Qiu and G. F. Sterman, *Phys. Rev. D* **59**, 014004 (1998).
- [26] S. Wandzura and F. Wilczek, *Phys. Lett.* **72B**, 195 (1977).
- [27] X.-D. Ji, *Phys. Rev. D* **49**, 114 (1994).
- [28] F. Yuan and J. Zhou, *Phys. Rev. Lett.* **103**, 052001 (2009).
- [29] P. G. Ratcliffe, *Nucl. Phys.* **B264**, 493 (1986).
- [30] S. Benic, Y. Hatta, H.-n. Li, and D.-J. Yang, *Phys. Rev. D* **100**, 094027 (2019).
- [31] H. Eguchi, Y. Koike, and K. Tanaka, *Nucl. Phys.* **B752**, 1 (2006).
- [32] K. Kanazawa, Y. Koike, A. Metz, D. Pitonyak, and M. Schlegel, *Phys. Rev. D* **93**, 054024 (2016).
- [33] E. Leader, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **15**, 1 (2011).
- [34] S. Bhattacharya, K. Cichy, M. Constantinou, A. Metz, A. Scapellato, and F. Steffens, *Phys. Rev. D* **102**, 111501 (2020).
- [35] S. Bhattacharya, Z.-B. Kang, A. Metz, G. Penn, and D. Pitonyak, *Phys. Rev. D* **105**, 034007 (2022).
- [36] Y. Hatta, K. Kanazawa, and S. Yoshida, *Phys. Rev. D* **88**, 014037 (2013).
- [37] S. Benić, Y. Hatta, A. Kaushik, and H.-n. Li, *Phys. Rev. D* **104**, 094027 (2021).
- [38] Y. Hatta, K. Tanaka, and S. Yoshida, *J. High Energy Phys.* **02** (2013) 003.
- [39] Y. Koike, K. Yabe, and S. Yoshida, *Phys. Rev. D* **101**, 054017 (2020).
- [40] A. Dumitru and J. Jalilian-Marian, *Phys. Rev. Lett.* **89**, 022301 (2002).
- [41] A. Dumitru, A. Hayashigaki, and J. Jalilian-Marian, *Nucl. Phys. A* **765**, 464 (2006).

- [42] A. Ayala, J. Jalilian-Marian, L. D. McLerran, and R. Venugopalan, *Phys. Rev. D* **52**, 2935 (1995).
- [43] L. D. McLerran and R. Venugopalan, *Phys. Rev. D* **59**, 094002 (1999).
- [44] I. I. Balitsky and A. V. Belitsky, *Nucl. Phys.* **B629**, 290 (2002).
- [45] J. Jalilian-Marian and Y. V. Kovchegov, *Phys. Rev. D* **70**, 114017 (2004); **71**, 079901(E) (2005).
- [46] P. Caucal, F. Salazar, and R. Venugopalan, *J. High Energy Phys.* **11** (2021) 222.
- [47] G. A. Chirilli, B.-W. Xiao, and F. Yuan, *Phys. Rev. D* **86**, 054005 (2012).
- [48] J. Jalilian-Marian, A. Kovner, A. Leonidov, and H. Weigert, *Nucl. Phys.* **B504**, 415 (1997).
- [49] J. Jalilian-Marian, A. Kovner, and H. Weigert, *Phys. Rev. D* **59**, 014015 (1998).
- [50] E. Iancu, A. Leonidov, and L. D. McLerran, *Nucl. Phys.* **A692**, 583 (2001).
- [51] E. Iancu, A. Leonidov, and L. D. McLerran, *Phys. Lett. B* **510**, 133 (2001).
- [52] T. Lappi, A. Ramnath, K. Rummukainen, and H. Weigert, *Phys. Rev. D* **94**, 054014 (2016).
- [53] H. Dong, D.-X. Zheng, and J. Zhou, *Phys. Lett. B* **788**, 401 (2019).
- [54] R. Boussarie, Y. Hatta, L. Szymanowski, and S. Wallon, *Phys. Rev. Lett.* **124**, 172501 (2020).
- [55] J. P. Blaizot, F. Gelis, and R. Venugopalan, *Nucl. Phys.* **A743**, 57 (2004).
- [56] M. Schlegel, *Phys. Rev. D* **87**, 034006 (2013).
- [57] J. Cammarota *et al.* (Jefferson Lab Angular Momentum (JAM) Collaboration), *Phys. Rev. D* **102**, 054002 (2020).
- [58] L. Gamberg *et al.* (Jefferson Lab Angular Momentum (JAM) Collaboration), *Phys. Rev. D* **106**, 034014 (2022).
- [59] S. J. Brodsky, D. S. Hwang, and I. Schmidt, *Phys. Lett. B* **530**, 99 (2002).
- [60] G. L. Kane, J. Pumplin, and W. Repko, *Phys. Rev. Lett.* **41**, 1689 (1978).