Addressing shape and extent of Weyl cones in TaAs by Landau level spectroscopy

Santos-Cottin, D.; Wyzula, J.; Le Mardelé, F.; Crassee, I.; Martino, E.; Novák, J.; Eguchi, G.; Rukelj, Zoran; Novak, M.; Orlita, M; ...

Source / Izvornik: Physical Review B (condensed matter and materials physics), 2022, 105

Journal article, Published version Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

https://doi.org/10.1103/PhysRevB.105.L081114

Permanent link / Trajna poveznica: https://urn.nsk.hr/um:nbn:hr:217:115574

Rights / Prava: In copyright/Zaštićeno autorskim pravom.

Download date / Datum preuzimanja: 2025-03-23



Repository / Repozitorij:

Repository of the Faculty of Science - University of Zagreb





Addressing shape and extent of Weyl cones in TaAs by Landau level spectroscopy

D. Santos-Cottin,¹ J. Wyzula,² F. Le Mardelé,¹ I. Crassee,² E. Martino,^{1,3} J. Novák,⁴ G. Eguchi⁰,⁵

¹Department of Physics, University of Fribourg, CH-1700 Fribourg, Switzerland

²LNCMI, CNRS-UGA-UPS-INSA, 25, avenue des Martyrs, F-38042 Grenoble, France

³IPHYS, EPFL, CH-1015 Lausanne, Switzerland

⁴Masaryk University, Faculty of Science, CZ-61137 Brno, Czech Republic

⁵Institute of Solid State Physics, Vienna University of Technology, Wiedner Hauptstrasse 8-10, AT-1040 Vienna, Austria

⁶Department of Physics, Faculty of Science, University of Zagreb, Bijenička 32, HR-10000 Zagreb, Croatia

⁷Institute of Physics, Charles University in Prague, CZ-12116 Prague, Czech Republic

(Received 20 November 2021; accepted 10 February 2022; published 25 February 2022)

TaAs is a prime example of a topological semimetal with two types of Weyl nodes, W_1 and W_2 , whose bulk signatures have proven elusive. We apply Landau level spectroscopy to crystals with multiple facets and identify—among other low-energy excitations between parabolic bands—the response of a cone extending over a wide energy range. Comparison with density functional theory studies allows us to associate this conical band with nearly isotropic W_2 nodes. In contrast, W_1 cones, which are more anisotropic and less extended in energy, appear to be buried too deep beneath the Fermi level. They cannot be accessed directly. Instead, the excitations in their vicinity give rise to an optical response typical of a narrow-gap semiconductor rather than a Weyl semimetal.

DOI: 10.1103/PhysRevB.105.L081114

Topological systems offer glimpses of relativisticlike physics within a crystal. The two most famous manifestations are Klein tunneling in graphene [1], and the chiral anomaly [2] in Weyl semimetals. These are materials whose broken inversion symmetry leads to pairs of Weyl nodes in the reciprocal space. The elementary excitations near those nodes are chiral Weyl quasiparticles. In TaAs-the first discovered and the most prominent Weyl semimetal-24 such nodes are sprinkled around the Brillouin zone [3–5]. Among them, eight are strongly anisotropic W1 nodes, and 16 are fairly isotropic W₂ nodes [6]. The large number and two different kinds of Weyl nodes have complicated our understanding of the underlying Weyl physics. TaAs was proposed as a platform to benefit from the special properties of Weyl fermions [7–9]. Yet, most basic questions remain unanswered: How can one directly access the linearly dispersing Weyl cones? What is their energy range? When do their excitations dominate over other collective excitations?

We address the Weyl cones of TaAs through Landau level and infrared spectroscopy, measuring several samples with multiple crystalline facets. Our key finding is that only one kind of Weyl cones in TaAs significantly contributes to its optical response. These cones are isotropic and large, extending over more than 250 meV. Comparison with density functional theory (DFT) studies allows us to recognize them as W_2 cones. In contrast, the W_1 cones are buried too deep below the Fermi energy to be optically accessible. Regardless, the electronic bands in their vicinity strongly contribute to the optical and magneto-optical response, because of their profound anisotropy, mirrored in a large joint density of states. This contribution dominates when the magnetic field is oriented along the [001] axis, effectively masking excitations in the W_2 cones.

We grew single crystals of TaAs exposing large mirrorlike surfaces in different crystalline planes: (001), (101), (112). The samples were characterized through magnetotransport and quantum oscillations [10]. Infrared spectra were then taken using a Fourier-transform infrared (FTIR) Vertex 70v spectrometer and employing an *in situ* coating technique [11]. Magneto-optical spectra were measured at 2 K, in fields up to 34 T. A single sample with large (001), (101), and (112) planes allowed us to confirm that all the measurements on different samples are reproducible and consistent [10].

The tetragonal symmetry of TaAs leads to a high number of Weyl points, shown in Fig. 1. Eight W_1 nodes with $k_c = 0$ sit near the time-reversal invariant momentum point Σ of the first Brillouin zone. Sixteen W2 nodes lurk further away from high symmetry points, with $k_c \neq 0$. The connecting lines between all pairs of Weyl nodes are parallel with the (001) plane. The crystal unit cell is four times longer along the c axis than along the *a* axis. Such an elongated shape creates a steep angle of $\sim 74^{\circ}$ between the (001) and (101) planes, and an angle of $\sim 67^{\circ}$ between the planes (001) and (112). The (001) plane of TaAs is most commonly addressed in the literature. In contrast, other directions are much less explored. Figure 1(a)schematically shows the shape of W_1 and W_2 cones in the (001) plane, after the DFT calculations [12]. There is no consensus in the literature on the extent and k-space separation of these Weyl cones. This is why we only give a schematic drawing of the cone shapes. A cut through the two Weyl cones is shown for the (101) plane in Fig. 1(b). Evidently, the W_2 cone is isotropic, but the W_1 cone becomes rather flat in the (101) plane. This flat energy dispersion of the W_1 cone

Z. Rukelj,^{1,6} M. Novak⁰,⁶ M. Orlita,^{2,7} and Ana Akrap^{1,*}

^{*}ana.akrap@unifr.ch



FIG. 1. Top: Brillouin zone of TaAs, with (001) and (101) planes shown in red and blue, respectively. Following the band structure calculations [12], we sketch the W_1 and W_2 Weyl cones (a) in the (001) plane, and (b) in the (101) plane.

along the c axis leads to an increased density of states (DOS). One may expect the same for the electronic states surrounding these cones.

Topological semimetals can be understood in fine detail using Landau level (LL) spectroscopy [13-19]. Their small Fermi pockets and high carrier mobility enable us to reach the quantum regime and observe the inter-LL transitions, letting us map the band structure. Figures 2(a) and 2(b) and 2(c) and 2(f) show the color plots of relative magnetoreflectance and its energy derivative, for (101) and (001) facets, respectively. The observed inter-LL resonances are narrow, so we associate the positions of maxima in R_B/R_0 spectra directly with the transition energies. The spectra were confirmed on different samples (see Supplemental Material [10] and Refs. [20,21] therein). The carrier concentrations were similar to the literature [6,22]. Magneto-optical spectra are exceptionally rich. Far below 1 T, TaAs reaches its quantum regime and the inter-LL transitions become apparent [23]. Figures 2(a)-2(c) show a striking series of inter-LL transitions along the (101) facet. The spectra are dominated by a series of sublinear lines-a nearly \sqrt{B} dependence of interband excitations on a magnetic field. This indicates a strongly nonparabolic dispersion, as expected for a conical dispersion stemming from Weyl nodes [17,24].

TaAs is a multiband system with a complex magnetooptical response, therefore we cannot analyze the experimental data using simple effective Hamiltonians [14–18]. Instead, to understand our spectra, we adopt a semiclassical approach [25]. The Lifshitz-Onsager formula says that in a magnetic field B, the area in momentum space encircled during cyclotron motion of an electron with crystal momentum k_n equals $S_n = \pi k_n^2 = (eB/h)(n + \gamma)$. Here, *n* is an integer enumerating individual cyclotron orbits, in fact, Landau levels. The parameter γ describes the Berry phase of the explored band. In the semiclassical approach, we may assume that the optical excitations conserve the momentum of electrons which are promoted between the electronic bands. Each interband inter-LL transition, observed at a given B and energy $\hbar\omega$, then measures the distance between bands at a given momentum $k_n = \sqrt{(n+\gamma)2eB/\hbar}$. In our case, rather than a Berry phase, γ represents an additional tuning parameter, compensating for the fact that momentum is no longer a good quantum number in quantizing magnetic fields. In reality, the excited electrons change their LL index, or the cyclotron orbit. For a simple system with full rotational symmetry around the direction of B, the selection rules $n \rightarrow n \pm 1$ apply in the Faraday configuration.

The described procedure allows us to deduce the joint profile, $E_c - E_v$, of the involved conduction and valence bands. Assigning each excitation a particular orbital integer n, we obtain a fairly smooth profile [Fig. 2(d)], leading us to conclude that a well-defined conical band extends over more than 250 meV, in agreement with the photoemission work [5]. There is no deviation from linearity up to the highest experimentally probed energies. Assuming a full electronhole symmetry, the slope gives a velocity of $v = (2.0 \pm$ $(0.2) \times 10^5$ m/s. This simple result of our analysis is surprising. We observe only one type of Weyl cones, while the other one does not manifest in our data. The occupation effect-Pauli blocking-is the most likely explanation. We may also conclude that the observed conical band is nearly isotropic. Otherwise, the pairs of Weyl cones oriented along the a and b crystallographic axes would have to contribute differently to the magneto-optical response when Bis applied along the [101] direction (see the Brillouin zone in Fig. 1). A weak anisotropy of the traced conical bands may explain the appearance of additional, fairly weak lines in our data [gray points in Fig. 2(c)]. The analysis of the data collected on the (112)-oriented facet-43° away from the (101) plane—also indicates a single conical band with a nearly identical velocity parameter [10]. DFT studies [4,12] show that W₂ cones should be isotropic. In contrast, W₁ cones have a flat dispersion along the tetragonal c axis, and their character is closer to quasi-two-dimensional rather than threedimensional (3D). Therefore, the conical feature we observe is W₂.

In stark contrast, the (001) data in Figs. 2(e)-2(h) show over 50 discernible lines, where linear-in-*B* transitions completely dominate the response. Clearly, in the (001) plane we are primarily looking at excitations between paraboliclike bands. No trace is seen of \sqrt{B} -like transitions, which would



FIG. 2. Inter-Landau-level transitions in (101) (top) and (001) planes (bottom). Relative magnetoreflectance R_B/R_0 is shown in (a) and (e), and its derivative in (b) and (f). The inter-LL transitions are isolated in (c) and (g). They are plotted in (d) and (h) as a function of momentum $k_n = \sqrt{2eB(n + \gamma)/\hbar}$, with $\gamma = 0$.

be primarily expected in a 3D Weyl semimetal. We may identify at least four separate series of transitions, plotted in different colors in Fig. 2(g). The oscillator strengths of these transitions are considerably stronger than for the excitations observed on other facets [10], effectively masking the \sqrt{B} -like transitions. At the same time, these linear-in-*B* excitations are completely missing from the response of other facets. This apparent anisotropy allows us to establish that these excitations originate in the vicinity of the W₁ cones—around the Σ point—where the band structure becomes significantly flatter in the tetragonal *c* direction [12].

Let us now analyze the response of the (001) facet in greater detail. Repeating the semiclassical approach [26], we first identify which transitions belong together. The energy of each inter-LL transition is then plotted as a function of k_n , after assigning each relevant transition an index n. We recognize up to four families of inter-LL transitions on the (001) facet. The most pronounced series of lines extrapolate to 15 meV in the limit of vanishing magnetic fields. More than 15 transitions collapse onto the same line in the energy versus the k_n plot in Fig. 2(h). From this "collapsed" plot, representing joint bands, we see that the underlying nearly parabolic bands have a gap of $\Delta \sim 15$ meV. A vertical line at any high field in Fig. 2(g) intersects a series of equidistant excitations. Their spacing, amounting to 15–20 meV at 30 T, gives the effective mass of these carriers, $m^* \approx 0.4m_e$. The lowest transition, $0 \rightarrow 1$, emerges at 6–8 T, and is marked in dark red in Fig. 2(g). This value gives us the size of the relevant Fermi pocket. In quantum oscillations, we have a frequency of 7–7.5 T, a size of the W₁ pocket which corresponds well to previously reported values [6,10,22]. Such a small pocket size would imply that W₁ is in the chiral regime. Yet, the response in the (001) plane appears to be dictated by a narrow-gap parabolic band. This striking absence of any observable Weyl cone signatures in the (001) orientation is an important result. It tells us that the W₁ cones are inaccessible for optical excitations, despite a very low carrier concentration.

TaAs is closely related to TaP, whose (001) plane shows peculiar down-dispersing inter-LL transitions [17]. Those transitions are related to the band inversion and the linear dispersion from W_1 cones. In TaAs, nothing similar can be seen: No lines shift to lower energies as the field increases. This is another strong indication that the Fermi level in TaAs is not placed within the W_1 cones, and the system is above the topological separatrix. Finally, on the (001) facet of TaAs we see no evidence of nonparabolic behavior at high energies, contrary to TaP [17]. In TaAs, the W_1 nodes do not seem to play a role, and the conical dispersion from W_2 points becomes remarkably evident once we move away from the (001) plane.

Figure 2(g) contains several unusual low-energy lines shown in black. These lines are only seen in high fields. The lowest line begins at 25 T, crosses 7 meV at 30 T, and extrapolates to zero energy at zero field. Assuming this is a cyclotron resonance (CR), its associated effective mass is $m_H^* = 0.5m_e$. Such heavy carriers in TaAs may be the trivial holes seen in quantum oscillations [6,10]. The CR line onset at 25 T corresponds to the heavy-hole pocket quantum limit. Three roughly equidistant lines appear above the lowest line. One may speculate that these are CR harmonics appearing due to strongly anisotropic band dispersion, similar to bulk graphite with electrons in the vicinity of a Lifshitz transition [27]. The trivial hole pocket in TaAs is indeed anisotropic and warped [6].

One pronounced yet unassigned inter-LL transition in the (001) plane is the line indicated in blue in Fig. 2(g). This line does not naturally belong with the straight lines emanating from 15 meV. Instead, it extrapolates linearly into zero energy at zero field, but flattens significantly above 20 T. Such a response may represent CR absorption of massive electrons in one of the partly occupied, strongly nonparabolic bands.

For both (101) and (001) facets, LL spectroscopy paints a full and complex picture of low-energy excitations in TaAs. This knowledge helps us better understand its zero-field infrared spectra. Optical conductivity primarily reflects the joint density of states, and thereby becomes strongly sensitive to the out-of-plane dispersion. Figure 3 shows the zero-field optical spectra of TaAs for (101) and (001) planes. The lowtemperature reflectivity shows sharp plasma edges at very low energies (15-20 meV), in line with low carrier concentration and small scattering rate $1/\tau$, which are necessary conditions to observe the LL quantization in fields below 1 T [Figs. 2(a) and 2(e)]. Through Kramers-Kronig transformations, we obtain the complex optical conductivity, $\hat{\sigma} = \sigma_1 + i\sigma_2$. Its real part $\sigma_1(\omega)$ is shown in Figs. 3(c) and 3(d). The large carrier mobility is reflected in a narrow Drude response, whose characteristic width is $1/\tau \sim 1 \text{ meV}$. Although TaAs is a multiband system, we only fit a single Drude contribution [28]. The Drude response is so narrow that it cannot be meaningfully separated into two or more contributions. The inset in Fig. 3 shows that the fitted Drude scattering rate $1/\tau$ follows a quadratic temperature dependence.

Most relevant to the Weyl node physics are the interband excitations, which in TaAs set in at remarkably low energies. The 10 K absorption onset is at 15 meV for the (001) facet, in agreement with the gap from Fig. 2(h). Interestingly, $\sigma_1(\omega)$ has sharper features for the (001) facet [Fig. 3(d)] than for the (101) facet [Fig. 3(c)]. This is caused by the flat dispersion of the bands near the Σ point [12], enhancing their optical response. Contrary to initial reports [29,30], we see several sharp kinks in $\sigma_1(\omega)$. Besides the absorption onset kink at 15 meV, the peaks at 70 and 128 meV can be linked to the zero-field extrapolation of several lines in Fig. 2(h). These may be van Hove singularities that could originate anywhere in the Brillouin zone, but because of the DOS effect, their source is most likely in the bands near the Σ point. All those fine features are superimposed upon a broad background. This



FIG. 3. Zero-field optical properties of TaAs, for (101) (top) and (001) plane (bottom panels). Far-infrared reflectance is shown in (a) and (b) as a function of photon energy, at different temperatures. The real part of optical conductivity σ_1 is shown in (c) and (d). Gray arrows indicate the 15 meV absorption onset, and van Hove peaks at 70 and 128 meV. The inset shows the temperature dependence of scattering rate $1/\tau$ for the two planes.

broad background is a sum of interband excitations from W_1 and W_2 pockets. We find that there is no simple meaning of the slopes in $\sigma_1(\omega)$, nor a link to Weyl cone effective velocities. The W_1 pocket is not in its chiral limit, and is therefore not expected to give a linear contribution to $\sigma_1(\omega)$. In the (101) plane [Fig. 3(c)], $\sigma_1(\omega)$ has a smoother profile, with a narrow Drude component and a broad maximum at 120 meV. We expect that W_2 nodes dominate in this orientation, although it is difficult to say what exactly they contribute to $\sigma_1(\omega)$. It seems abundantly clear that the W_2 contribution is not simply a straight line from 0 to the maximum cone extent.

In conclusion, we found a strong difference in the optical and magneto-optical response of TaAs for different orientations, specifically planes (001) and (101). This anisotropic response is caused by the anisotropy of bands giving rise to the W_1 nodes. Crucially, we found that W_1 nodes are not optically accessible even in low-carrier-density crystals of TaAs. The magneto-optical response of the (001) facet, most often addressed experimentally, does not show any direct signatures of Weyl physics. Instead, optical experiments can only probe the neighboring parabolic bands. In contrast, our magneto-optical data give strong evidence that W_2 cones are highly isotropic and extend over more than 250 meV. They are characterized by a relatively

low velocity parameter of $(2.0 \pm 0.2) \times 10^5$ m/s. Therefore, in the (101) plane and other planes closer to the crystal *c* axis, optical experiments can readily detect topologically nontrivial bands.

We thank F. Bechstedt for deeply informative discussions, and N. Miller for helpful comments. A.A. acknowledges funding from the Swiss National Science Foundation through

- [1] A. F. Young and P. Kim, Nat. Phys. 5, 222 (2009).
- [2] N. P. Armitage, E. J. Mele, and A. Vishwanath, Rev. Mod. Phys. 90, 015001 (2018).
- [3] B. Q. Lv, N. Xu, H. M. Weng, J. Z. Ma, P. Richard, X. C. Huang, L. X. Zhao, G. F. Chen, C. E. Matt, F. Bisti, V. N. Strocov, J. Mesot, Z. Fang, X. Dai, T. Qian, M. Shi, and H. Ding, Nat. Phys. 11, 724 (2015).
- [4] H. Weng, C. Fang, Z. Fang, B. A. Bernevig, and X. Dai, Phys. Rev. X 5, 011029 (2015).
- [5] S.-Y. Xu, I. Belopolski, N. Alidoust, M. Neupane, G. Bian, C. Zhang, R. Sankar, G. Chang, Z. Yuan, C.-C. Lee, S.-M. Huang, H. Zheng, J. Ma, D. S. Sanchez, B. Wang, A. Bansil, F. Chou, P. P. Shibayev, H. Lin, S. Jia, and M. Z. Hasan, Science 349, 613 (2015).
- [6] F. Arnold, M. Naumann, S. C. Wu, Y. Sun, M. Schmidt, H. Borrmann, C. Felser, B. Yan, and E. Hassinger, Phys. Rev. Lett. 117, 146401 (2016).
- [7] Q. Ma, S.-Y. Xu, C.-K. Chan, C.-L. Zhang, G. Chang, Y. Lin, W. Xie, T. Palacios, H. Lin, S. Jia, P. A. Lee, P. Jarillo-Herrero, and N. Gedik, Nat. Phys. 13, 842 (2017).
- [8] Y. Gao, S. Kaushik, E. J. Philip, Z. Li, Y. Qin, Y. P. Liu, W. L. Zhang, Y. L. Su, X. Chen, H. Weng, D. E. Kharzeev, M. K. Liu, and J. Qi, Nat. Commun. 11, 720 (2020).
- [9] G. B. Osterhoudt, L. K. Diebel, M. J. Gray, X. Yang, J. Stanco, X. Huang, B. Shen, N. Ni, P. J. W. Moll, Y. Ran, and K. S. Burch, Nat. Mater. 18, 471 (2019).
- [10] See Supplemental Material at http://link.aps.org/supplemental/ 10.1103/PhysRevB.105.L081114 for additional data to support our work.
- [11] C. C. Homes, M. Reedyk, D. A. Cradles, and T. Timusk, Appl. Opt. 32, 2976 (1993).
- [12] D. Grassano, O. Pulci, A. Mosca Conte, and F. Bechstedt, Sci. Rep. 8, 3534 (2018).
- [13] M. Orlita, D. M. Basko, M. S. Zholudev, F. Teppe, W. Knap, V. I. Gavrilenko, N. N. Mikhailov, S. A. Dvoretskii, P. Neugebauer, C. Faugeras, A.-L. Barra, G. Martinez, and M. Potemski, Nat. Phys. **10**, 233 (2014).
- [14] A. Akrap, M. Hakl, S. Tchoumakov, I. Crassee, J. Kuba, M. O. Goerbig, C. C. Homes, O. Caha, J. Novák, F. Teppe, W. Desrat, S. Koohpayeh, L. Wu, N. P. Armitage, A. Nateprov, E. Arushanov, Q. D. Gibson, R. J. Cava, D. van der Marel, B. A. Piot, C. Faugeras, G. Martinez, M. Potemski, and M. Orlita, Phys. Rev. Lett. **117**, 136401 (2016).
- [15] M. Hakl, S. Tchoumakov, I. Crassee, A. Akrap, B. A. Piot, C. Faugeras, G. Martinez, O. Caha, J. Novk, A. Nateprov, E. Arushanov, W. L. Lee, M. O. Goerbig, F. Teppe, M. Potemski, and M. Orlita, in 2017 42nd International Conference on

Project No. PP00P2_170544. M.N. acknowledges support of CeNIKS project cofinanced by the Croatian Government and the EU through the European Regional Development Fund Competitiveness and Cohesion Operational Program (Grant No. KK.01.1.1.02.0013). This work has been supported by ANR projects DIRAC3D and COLECTOR. We acknowledge the support of LNCMI-CNRS, a member of the European Magnetic Field Laboratory (EMFL).

Infrared, Millimeter, and Terahertz Waves (IRMMW-THz) (IEEE, New York, 2017), pp. 1–2.

- [16] E. Martino, I. Crassee, G. Eguchi, D. Santos-Cottin, R. D. Zhong, G. D. Gu, H. Berger, Z. Rukelj, M. Orlita, C. C. Homes, and A. Akrap, Phys. Rev. Lett. **122**, 217402 (2019).
- [17] S. Polatkan, M. O. Goerbig, J. Wyzula, R. Kemmler, L. Z. Maulana, B. A. Piot, I. Crassee, A. Akrap, C. Shekhar, C. Felser, M. Dressel, A. V. Pronin, and M. Orlita, Phys. Rev. Lett. 124, 176402 (2020).
- [18] Y. Shao, Z. Sun, Y. Wang, C. Xu, R. Sankar, A. J. Breindel, C. Cao, M. M. Fogler, A. J. Millis, F. Chou, Z. Li, T. Timusk, M. B. Maple, and D. N. Basov, Proc. Natl. Acad. Sci. USA 116, 1168 (2019).
- [19] Y. Shao, A. N. Rudenko, J. Hu, Z. Sun, Y. Zhu, S. Moon, A. J. Millis, S. Yuan, A. I. Lichtenstein, D. Smirnov, Z. Q. Mao, M. I. Katsnelson, and D. N. Basov, Nat. Phys. 16, 636 (2020).
- [20] X. Huang, L. Zhao, Y. Long, P. Wang, D. Chen, Z. Yang, H. Liang, M. Xue, H. Weng, Z. Fang, X. Dai, and G. Chen, Phys. Rev. X 5, 031023 (2015).
- [21] R. Sankar, G. Peramaiyan, I. P. Muthuselvam, S. Xu, M. Z. Hasan, and F. C. Chou, J. Phys.: Condens. Matter 30, 015803 (2018).
- [22] B. J. Ramshaw, K. A. Modic, A. Shekhter, Y. Zhang, E.-A. Kim, P. J. W. Moll, M. D. Bachmann, M. K. Chan, J. B. Betts, F. Balakirev, A. Migliori, N. J. Ghimire, E. D. Bauer, F. Ronning, and R. D. McDonald, Nat. Commun. 9, 2217 (2018).
- [23] Z. Lu, P. Hollister, M. Ozerov, S. Moon, E. D. Bauer, F. Ronning, D. Smirnov, L. Ju, and B. J. Ramshaw, Sci. Adv. 8, eabj1076 (2022).
- [24] P. E. C. Ashby and J. P. Carbotte, Phys. Rev. B 89, 245121 (2014).
- [25] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Holt-Saunders, Philadelphia, 1976).
- [26] A. Miyata, Z. Yang, A. Surrente, O. Drachenko, D. K. Maude, O. Portugall, L. B. Duffy, T. Hesjedal, P. Plochocka, and R. J. Nicholas, Phys. Rev. B 96, 121111(R) (2017).
- [27] M. Orlita, P. Neugebauer, C. Faugeras, A. L. Barra, M. Potemski, F. M. D. Pellegrino, and D. M. Basko, Phys. Rev. Lett. 108, 017602 (2012).
- [28] A. B. Kuzmenko, Rev. Sci. Instrum. 76, 083108 (2005).
- [29] B. Xu, Y. M. Dai, L. X. Zhao, K. Wang, R. Yang, W. Zhang, J. Y. Liu, H. Xiao, G. F. Chen, A. J. Taylor, D. A. Yarotski, R. P. Prasankumar, and X. G. Qiu, Phys. Rev. B 93, 121110(R) (2016).
- [30] S.-i. Kimura, H. Yokoyama, H. Watanabe, J. Sichelschmidt, V. Süß, M. Schmidt, and C. Felser, Phys. Rev. B 96, 075119 (2017).