

Kemoprevencija tumora polifenolnim sastavnicama

Marošević, Đurđica

Undergraduate thesis / Završni rad

2009

Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj: **University of Zagreb, Faculty of Science / Sveučilište u Zagrebu, Prirodoslovno-matematički fakultet**

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:357211>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-11-22**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



Sveučilište u Zagrebu
Prirodoslovno-matematički fakultet
Biološki odsjek
Rooseveltova trg 6

**KEMOPREVENCIJA TUMORA POLIFENOLNIM
SASTAVNICAMA**

Chemoprevention of carcinogenesis by polyphenolic compounds

Student: Miroslava Marošević

Molekularna biologija, molecular biology

Mentor: prof.dr. Nada Oršolić

Zagreb, 2009.

Sadržaj

KEMOPREVENCIJA TUMORA POLIFENOLNIM SASTAVNICAMA	1
Chemoprevention of carcinogenesis by polyphenolic compounds	1
Sadržaj	2
Uvod	3
Kemijska svojstva	3
Biološka dostupnost	4
Flavonoli	5
Izoflavoni (genistein i daidzein)	5
Stilbeni	5
Katehini [(+)-katehin, (-)-epikatehin, (-)-epigalokehin, (-)-epikatehin galat, (-)-epigallocatehin galat]	6
Inhibicija rasta tumora i mehanizam djelovanja	7
Katehini	7
Stilbeni	9
Propolis i polifenolne sastavnice propolisa	12
Literatura	17
Sažetak	22
Summary	23

Uvod

Polifenoli predstavljaju najbrojniju skupinu biljnih sekundarnih metabolita. Biljkama služe kao zaštita od fotosintetskog stresa, reaktivnih kisikovih radikala, herbivora ili ozljeda. Važan su dio ljudske prehrane, ponajprije fenolna kiselina i flavonoidi, zato što ovjek dnevno unese i do 1 g polifenola, što je mnogo više nego svih drugih poznatih antioksidansa i fitokemikalija. Za usporedbu, to je 10 puta više od vitamina C ili 100 puta više od vitamina E koji ovjek dnevno unese u organizam (Manach i sur., 2004; Scalbert i sur. 2000). Najprije su u centru pažnje bili zbog svog „antinutricijskog u inka“, smanjenja apsorpcije i probave određene hrane, te zbog svoje mogući vezanja proteina. Danas pobu uju interes svojim protuupalnim, protuoksidativnim i protutumorskim osobitostima. U ovom radu pokazana su potencijalna preventivna protutumorska svojstva.


Kemijska svojstva

Prema kemijskoj strukturi polifenoli su fenilbenzo-pironi (fenilhormoni) s brojnim skupinama različitih struktura temeljenih na jezgri od 3 prstena. Najčešće se dijele na osnovu supstituenata na flavanole, antocijane, flavone, flavanone i halkone. Ova osnovna struktura obuhvaća 2 benzenska prstena (A i B) povezani heterocikličkim prstenom pirana ili pirona (C) u sredini. Podjela se temelji na prisutstvu ili odsutstvu dvostruke veze na poziciji 4 C prstena, ili dvostruke veze između 2 i 3 ugljikovog atoma C prstena i prisutnosti hidroksilne grupe na B prstenu. U flavonoidnoj strukturi, fenilna grupa je obično zamijenjena na poziciju 2.pironskog prstena. Kod izoflavonoida zamjena je na poziciji 3. Flavonoidi i tokoferoli (vitamin E) dijele zajedničku strukturu, kromanski prsten (Middleton i sur., 2000).

Tablica 1. Primjer podjele flavonoida (Middleton i sur., 2000)

Class	Flavonoids	Substituents					
		3'	5'	7'	8'	4'	6'
Flavan-3-ols	(-)-Catechin	OH	OH	OH	OH	OH	H
Anthocyanidins	Cyanidin	OH	OH	OH	OH	OH	H
	Pelargonidin	OH	OH	OH	OH	H	H
Flavones	Apigenin	H	OH	OH	H	OH	H
	Diosmin	H	OH	OH	OH	OH	H
	Luteolin	H	OH	OH	OH	OH	H
Flavanones	Naringenin	H	OH	OH	H	OH	H
	Naringin	H	OH	OH	H	OH	H
	Hesperetin	H	OH	OH	OH	OH	H
	Hesperidin	H	OH	OH	OH	OH	H
Chalcones	Floretin	OH (2 ^a)	OH (4)	OH (6)	H	H	OH (6')
	Flavichin	OH (2)	H (4)	OH (6)	H	H	OH (6')
Flavan-3-ols	Quercetin	OH	OH	OH	H	OH	H
	Kaempferol	OH	OH	OH	H	OH	H
	Myricetin	OH	OH	OH	OH	OH	OH
	Fisetin	OH	H	OH	OH	OH	H
	Morin ^b	OH	OH	OH	H	OH	H

OH, hydroxyl.
^a Number in parentheses denotes additional similar substituent at the position indicated by the number.
^b Morin has one more OH group at position 2'.



Slika 1. Kemijske strukture naj eš ih flavonoidnih podskupina (Middleton i sur., 2000)

Biološka dostupnost

Iako je napravljeno ve na tisu e radova o u incima polifenola na biološke sustave *in vitro*, oni ne uzimaju u obzir dostupnost i metabolizam u organizmu *in vivo*, tako da u inci koji su tamo uo eni ne moraju nužno važiti i za *in vivo* istraživanja.

In vitro istraživanja su ra ena s razli itim koncentracijama polifenola od <0,1 μmol/L pa sve do >100 μmol/L, no u fiziološkim uvjetima je zamijeno da ta koncentracija nikad ne

premašuje 10 $\mu\text{mol/L}$ i zato se svi rezultati koji su dobiveni pri višim koncentracijama ne uzimaju u obzir. Iznimka je probavna šupljina, za koju se pretpostavlja, ali još nije dokazano da je u doticaju s višom koncentracijom polifenola.

Još jedan važan detalj je to što su u organizmu polifenoli konjugirani s različitim skupinama, te da nisu istog oblika koji imaju u hrani i koji se primjenjuje u *in vitro* istraživanjima.

Flavonoli

Kvercetin kao jedan od predstavnika flavonola prisutan u luku, jabukama, aju, brokuli, crvenom vinu i biljci ginko biloba, iako pokazuje mnoštvo pozitivnih u inaka *in vitro*, dokazi o njegovom djelovanju *in vivo* u ljudima izostaju. Pretpostavlja se da je najveća prepreka njegovu djelovanju to što je u organizmu konjugiran sa sulfatima, metilom, ili šećerima, zato što u plazmi nikad nije pronađen u slobodnom obliku (Day i sur. 2001). Nema mnogo dokaza o aktivnosti konjugiranih oblika, ali se zaključiti da su mnogo manje biološki aktivni u usporedbi sa aglikonima (Day i sur., 2000; Cano i sur., 2002)

Izoflavoni (genistein i daidzein)

Najviše u inaku pokazuju na biomarkerima kosti (povećanje gustoće, sadržaj minerala, smanjena ekskrecija piridinium križnih veza, povećana koncentracija za kosti specifičnih alkalnih fosfataza i osteokalcina u serumu itd.) Mnoge, ali ne sve promjene mogu biti rezultat vezanja za estrogene receptore (Setchell i sur., 1999; 2002). U plazmi su pronađeni i u slobodnom obliku i konjugirani kao sulfati ili glukuronidi (Setchell i sur., 2001) i ali nije jasno da li je to rezultat aglikona ili su aktivni i konjugirani oblici, ili dolazi do dekonjugacije u tkivu. Postoje dokazi i da su neki od konjugata aktivni *in vitro* (Zhang i sur., 1999).

Stilbeni

Nisu česti u prirodi, biljke ih proizvode samo kao odgovor na infekciju ili ozlijeđenu. Najčešće istraživani je *trans*-resveratrol (3,5,4'-trihidroksstilben) koji je prisutan u grožđu, vinu i kikirikiju. Može postojati u *cis*- i *trans*- stereizomernom obliku, te je ishodišna molekula za obitelj polimera poznatu pod nazivom viniferini. Iz različitih *in vivo* istraživanja se može zaključiti da se apsorbira iz gastrointestinalnog sustava, no uinkovitost apsorpcije, raspodjela, metabolizam i ekskrecije nisu još dobro razjašnjeni. Pronađen je u visokim koncentracijama u crvenom vinu (2-40 μM), no uočeno je da ima kratak poluživot što rezultira niskom dostupnošću (Kopp i sur., 1998). Kao i kod ostalih polifenola ovo predstavlja najveći i problem za njegovu upotrebu *in vivo* i u kliničke svrhe kao protutumorskog spoja.

Koncentracije potrebne za inhibiciju rasta stanica su izme u 5-10 μM (Gescher i sur., 2003), dok nedavna istraživanja na ljudima pokazuju koncentracije slobodnog resveratrola u plazmi oko 21 nM, nakon uzimanja 25 mg, ili 360 $\mu\text{g/kg}$ (Goldberg i sur., 2003). No, pretpostavlja se da bi i glukoronirani i sulfatni konjugati mogli biti aktivni, budu i da dolazi do njihove dekonjugacije na ciljnom mjestu (Walle i sur., 2004; Kuhnle i sur., 2000; Signorelli i sur., 2005).

Katehini [(+)-katehin, (-)-epikatehin, (-)-epigalokehin, (-)-epikatehin galat, (-)-epigalocatehin galat]

(+)-Katechini su prisutni u crnom grož u, kajsijama, jagodama.

(-)-epikatehini su prisutni u jabukama, kupinama, malinama, trešnjama, crnom grož u, kruškama, okoladi, dok (-)-epigalokehin (EGC), (-)-epikatehin galat (ECG), (-)-epigalocatehin galat (EGCG), možemo na i isklju ivo u ajevima, posebno zelenom.

Za sve njih je dokazan širok spektar djelovanja *in vitro*, ali je teže provoditi istraživanja *in vivo*, zato što su široko rasprostranjeni i nikada ne dolaze sami u namirnici, dok je sa EGC, ECG i EGCG jednostavnije raditi. EGCG se u plazmi može otkriti i kao slobodan (Lee i sur., 2002), nekonjugirani oblik, dok su (+)-katehini (Holt i sur., 2002; Donovan i sur., 1999) i (-)-epikatehini prisutni isklju ivo kao konjugati (Holt i sur., 2002; Yang i sur., 1998; Bell i sur., 2000).

Velika ve ina istraživanja temelji se na intaktnim polifenolima nazo nim u hrani, koji nakon unosa u organizam i prije no što dospiju u stanice crijevnog lumena, pro u kroz crijevnu mikrofloru i tamo budu metabolizirani. Daidzein pod utjecajem crijevnih mikroorganizmima metabolizira se do ekvola i tek kao takav ulazi u krvotok; proces se doga a u 30-40% ljudske populacije. Setchell i sur. (2002) u svom istraživanju su pokazali kako „ekvol proizvo a i“ pokazuju bolje rezultate nekih biomarkera nakon uzimanja izoflavona (Setchell i sur., 2002). Otkriveno je i nekoliko metabolita u ljudi nakon konzumacije kokosovih procianidina (oligomerni katehini, kovalentno povezanih), ali nije istražena njihova biološka aktivnost (Rios i sur., 2003). Potrebna su još detaljnija, dugoro nija i iscrpnija istraživanja, kao i ona koja pokazuju ovisnost utjecaja i djelovanja o koncentraciji, te je potrebno raditi sa istim, izoliranim komponentama, a ne s hranom, kako bi se isklju ila mogu nost me udjelovanja razli itih sastojaka. Nužno je istraživati ulogu mikroflora u metabolizmu i biološkoj aktivnosti polifenola, kako bi se mogli izvesti zaklju ci o mogu oj klini koj primjeni.

Inhibicija rasta tumora i mehanizam djelovanja

Katehini

Kao što je već napomenuto polifenoli i njihova protutumorska aktivnost se intenzivno istražuju zadnjih desetljeća, no još uvijek nije napravljen dovoljan pomak i dobivena saznanja koja bi omogućila kliničku primjenu.

U *in vitro* istraživanjima većina komponenti pokazuje bolju ili lošiju protutumorsku aktivnost dok u *in vivo* istraživanjima bolje djeluju na inhibiciju tumora crijeva (Yang i sur., 2002), a nešto slabije na tumore unutarnjih organa poput pluća (Witschi i sur., 1998). Pripravci zelenog ili crnog čaja u inkubito inhibiraju tumore pluća prouzročene 4-(metilnitrosamino)-1-(3-piridil)-1-butanonom (NNK) u A/J miševima. Brojnost tumora bila je smanjena, spontana pojavnost tumora, odnosno proliferacija i progresija iz adenoma u karcinom (Yang i sur., 1997, Landau i sur., 1998). Također je zabilježena inhibicija invazije i metastaziranja tumora u transplantiranih i spontano metastaziraju ih modela prilikom oralnog uzimanja zelenog čaja ili pripravaka sa EGCG (Sazuka i sur., 1995, Liu i sur., 2001). Nedavnim istraživanjima je dokazano kako pripravci čaja djeluju na sve stadije tumora pluća, te da pojačavaju djelovanje protutumorskog lijeka doksorubicina (Yang i sur., 2002, 2005). Slično u inak pokazuje i vitamin E i pirolidineditiokarbamat (PDTC), te predobrada propolisom (Chinery i sur. 1997, Han i sur., 2000). I za polyphenon E i kofein je dokazano da smanjuju pojavu i učestalost plućnih adenokarcinoma, dok nemaju utjecaja na normalne plućne stanice; indeks proliferacije ili stupanj apoptoze nije bio promijenjen u normalnih stanica. Pripravci čaja smanjuju fosforilaciju proteina c-Jun i Erk1/2 što je pokazatelj mogućeg mehanizma djelovanja polifenola (Yang i sur., 2007).

U nekim istraživanjima se pokazalo da životinje obrađene 1% čajem ili kofeinom imaju manju tjelesnu težinu za 14% i manje tjelesnih masti za 35% od kontrolnih grupa (Yang i sur., 1997, Landau i sur., 1998, Xu i sur. 1992). Nije još jasno da li postoji veza između smanjene težine i mehanizma prevencije tumora ili je to slučajnost.

Li i sur. (2002) su u svom istraživanju na hrčcima pokazali da 0,6% pripravak zelenog čaja smanjuje pojavnost tumora prouzročenu oralnom karcinogenezom sa 7,12-dimetilbenz[a]antracenom (DMBA) za 35% i veličinu tumora za 57%. Imunohistokemijskim metodama pokazan je smanjeni proliferacijski indeks a povećani indeks apoptoze (Li i sur., 2002).

Na modelu transgeni nog miša s adenokarcinomom prostate TRAMP (engl. transgenic adenocarcinoma mouse prostate) [transgeni model miša soja C57Bl/6 koji nosi ulomak -

426/+28 štakorskog probasin (PB) gena fuzioniran sa SV40 T antigenom (Tag)] pokazano je da unos 0,1% polifenola zelenog aja, smanjuje pojavnost tumora za 65% i metastaza u odnosu na kontrolnu grupu za 25-95%. Biokemijske i histološke analize pokazale su značajan pad antigena proliferiraju ih stanica, te deseterostruki porast apoptoze tumorskih stanica (Gupta i sur., 2001). I najnovije istraživanje sa TRAMP mišjim modelom napravljenim u svrhu otkrivanja najpogodnijeg stadija raka prostate za obradu polifenolima zelenog aja, potvrdilo je logičnu pretpostavku kako kemopreventivni potencijal polifenola zelenog aja opada sa stupnjem napredovanja bolesti (Adhami i sur., 2009). I iste sastavnice izolirane iz zelenog aja imaju neka kemopreventivna svojstva. Primjerice, EGCG inhibira razvoj tumora pluća u A/J miševa induciranih 4-(metilnitrosamino)-1-(3-piridil)-1-butanonom (NNK) i cisplatinom (Xu i sur., 1992, Mimoto i sur., 2000), te kod $Apc^{Min/+}$ miševa smanjuje pojavu tumora crijeva čak za 51%, dok je polifenon E (PPE) djelotvoran u normalnoj prehrani samo 24,5%, a u AIN93G dijeti čak 70,5% smanjuje učestalost tumora, dok ECG nije pokazivao nikakav učinak (Hao i sur., 2006).

Švicarski albino miševi obrađeni dietilnitrosoaminom (DEN) i polifenolima iz crnog ili zelenog aja, pokazuju manju pojavu alveologenih tumora od kontrolne grupe koja nije primala pripravke polifenola. Mehanizam djelovanja temelji se na smanjenoj ekspresiji Akt signalnog puta, ciklooksigenaze 2, te inaktivaciji nuklearnog faktora kapa B (NF κ B) blokiranjem fosforilacije i razgradnje inhibitora NF κ B (I κ B) (Roy i sur., 2009).

U istraživanjima rađenim sa štakorima, rezultati nisu ovako jednoznačni. U nekim istraživanjima je pokazano inhibitorno djelovanje polifenola na razvoj tumora, dok u drugim nije. Razvoj tumora crijeva u štakora induciranih azoksimetanom (AOM) je bio smanjen, ako su životinjama davani pripravci zelenog aja u tekuci za piće (Kim i sur., 1994), dok kod AOM obrađenih F344 štakora iscrpak zelenog aja pokazuje samo marginalni učinak na formiranje žarišta aberantni kripti (engl. aberant crypti foci ACF) (Challila i sur., 1997). Wargovich i sur. (2000) su u jednom sličnom pokusu AOM obrađenim štakorima davali različite pripravke iz crnog ili zelenog aja, pri čemu je pokazano da EGCG i polifenoli crnog aja inhibiraju formiranje ACF-a dok zeleni čaj ima marginalni učinak, a iscrpki zelenog i crnog aja ne pokazuju nikakav učinak (Wargovich i sur., 2000, Cardeni i sur., 2000). Jedno starije istraživanje s F344 štakorima obrađenim 2-amino-3-metilimidazo[4,5-f]kviinolinom donosi potpuno suprotne rezultate; 2% otopina iscrpka zelenog aja primijenjena oralno, kao tekuci za piće, značajno inhibira ACF formiranje, dok 1% otopina iscrpka crnog aja ne inhibira (Xu i sur., 1996). Nadalje istraživanja Hirosea i sur. (2001) ne dokazuju učinkovito djelovanje zelenog aja u štakora s 1,2-dimethylhydrazine (DMH) induciranim tumorima

(Hirosea i sur., 2001, 2002). Sli ne rezultate je dobio i Weisburger, zato što ni iscrpak zelenog aja, ni iscrpak crnog aja, ni EGCG dodani u prehranu, ne pokazuju inhibitorni u inak na AOM inducirane tumore crijeva (Weisburger i sur., 1998). Istraživanje s NNK induciranim tumorima na F344 štakorima pokazuje kako inhibitorna aktivnost crnog aja poti e od kafeina (Chung i sur., 2003), kao i u karcinogenezi kože (Huang i sur., 1997), gdje je pokazano da su oralno davani zeleni i crni aj u inkovitiji u reduciranju u estalosti i brojnosti UVB induciranih tumora od dekafeiniranog aja. Tako er su djelotvorni i kafein i EGCG ako su nanošeni direktno na kožu SKH-1 bezdlakih miševa. Oni su prije toga obra ivani 2 puta tjedno tijekom 20 tjedana UVB svjetlom. U estalost kožnih tumora je smanjena za 44-70% ili 55-66%, te je u oba slu aja pove an apoptoti ki indeks za 56-92% (Lu i sur., 2002). Sli no istraživanje provedeno s presatcima kože dobrovoljaca pokazuje da iscrpci zelenog i bijelog aja nanesen i na kožu zaštitno djeluju na imunost kože, te smanjuju štetan u inak UV zra enja, ali ne zbog direktne apsorpcije sun evih zraka, zato što je za oba pripravka dokazano da imaju zaštitni imbenik SPF 1 (Camouse i sur., 2009). Najnovija istraživanja na modelima štakora s rakom jetre induciranim dimetilaminoazobenzenom (DAB) potvr uju protutumorsko djelovanje polifenola crnog aja. Grupa štakora koja je pored bazalne prehrane i 0,06% DAB unutar 3 mjeseca primala 0,05% polifenola crnog aja imala je zna ajno manje hepatoma, moduliraju i aktivnost biljega invazije i angiogeneze kao i ekspresiju deacetilaze histona HDAC-1 (Scalbert i sur., 2009).

Stilbeni

Resveratrol je jedan od najzanimljivijih polifenola i brojni radovi temelje se na njegovu djelovanju i utjecaju na razvoj tumora i to moduliraju i nekoliko signalnih puteva.

Usporedbom s nekoliko polifenola, katehini, kvercetin, gali na kiselina prisutnih u crvenom vinu, u dvostadijskom CD-1 mišjem modelu raka kože koriste i DMBA kao inicijator a TPA (forbol 12-miristat 13-acetat) kao promotor, pokazalo se da je *trans*-resveratrol naju inkovitiji protutumorski polifenol prisutan u crvenom vinu (Soleas i sur., 2002).

U usporedbi sa sezamom i sezamovim uljem u drugom istraživanju u *in vivo* DMBA injiciranim i TPA promoviranom mišjem raku kože, se tako er pokazao resveratrol kao najaktivniji (Kapadia i sur., 2002).

Tako su Carbo i sur. pokazali na mužjacima Wister štakora, inokuliranim s brzo rastu im Yoshida AH-130 tumorima jetre, da resveratrol zna ajno smanjuje sadržaj tumorskih stanica i

to zaustavljanjem stani nog ciklusa u G2/M fazi i apoptozom (Carbo i sur., 1999). Sli ni rezultati su dobiveni i s ljudskim oralnim skvamoznim stanicama karcinoma (SCC-25), kod kojih je obrada resveratrolom inhibirala rast i DNA sintezu, te u stanicama akutne promijelocitne leukemije ljudi (HL-60) i to indukcijom apoptoze preko Bcl-2 puta (Surh i sur., 1999). Indukcija apoptoze, inhibicija rasta i modulacija specifi nih gena za prostatu je istraživana na DU-145, PC-3. i JCA-1, i CaP stanicama ljudi. Pokazano je da resveratrol negativno modulira rast stanica i poti e apoptozu (Hsieh i sur., 1999), te op enito da mnogi polifenoli prisutni u vinu inhibiraju proliferaciju ljudskih stanica tumora prostate i to tako što inhibiraju stvaranje NO (Kampa i sur., 2000). Pokazano je da u stanicama tumora prostate izaziva dva suprotna u inka, induciranje S faze i inhibiciju sinteze DNA (Kuwajerwala i sur., 2002); mehanizam inhibicije proliferacije tumorskih stanica još nije sasvim jasan, no pretpostavlja se da ima veze s njegovom sposobnoš u inhibicije ribonukleotid reduktaze i ciklooksigenaze 2 (Fontecave i sur., 1998; Cuendet i sur., 2000). Tako er zamije eno je da je ekspresija proteina specifi nog za prostatu (PSA) bila zna ajno smanjena obradom s resveratrolom putem AR neovisnog mehanizma (Hsieh i sur., 1999, 2000). cDNA mikrotestovi su pokazali razli itu ekspresiju gena u stanicama tumora prostate kao odgovor na 48 satnu izloženost resveratrolu. Androgen senzitivne LNCaP stanice pokazuju ukupno 555 gena koji imaju razli itu ekspresiju, ovisnu o resveratrolu (Narayanan i sur., 2002). Tako er pokazuje direktan antiproliferativni u inak na stanice tumora dojke, neovisno o njihovom statusu estrogenskih receptora (Mgbonyebi i sur., 1998; Gehm i sur., 1997), te inhibiciju rasta 4T1 stanica ovisno o dozi i vremenu djelovanja (Bove i sur., 2002). Još nije jasno dali je resveratrol agonist ili antagonist estrogena, najvjerojatnije inteferira sa estrogenskim receptorom kako bi inhibirao njegovu aktivaciju (Gehm i sur., 1997; Lee i sur., 2001; Lu i sur., 1999). No tako er je inhibirao i proliferaciju stani ne linije ljudskog tumora dojke MDAMB-468 koja nema estrogen receptor, a i zna ajno povisio ekspresiju mRNA TGF- 2 inhibitora rasta, što navodi na zaklju ak da resveratrol inhibira proliferaciju mijenjaju i putove autokrinih modulatora rasta (Basly i sur., 2000; Serrero i sur., 2001).

Moduliraju i razinu H₂O₂, glutationa, te aktivnosti mijeloperoksidaze, oksidirane glutation reduktaze i superoksid dismutaze i smanjuju i razinu ekspresije TGF- 1, predobrada resveratrolom smanjuje štetan u inak TPA obrade, te inhibira de novo sintezu inducibilne nitrooksid sintaze u mišjim makrofagima (Jang i sur., 1999).

She i sur. su radili na stani noj liniji mišje epiderme JB6 i pokazali da resveratrol aktivira ERK, c-Jun NH₂ terminalnu kinazu (JNK) i p38 kinazu te time inducira fosforilaciju serina 15 u p53 (She i sur., 2001). Kao i iscrpci zelenog i bijelog aja, i površinska primjena

resveratrola dokazano inhibira UVB posredovanu indukciju COX i ornitin decarboksilaznu (ODC) enzimsku aktivnost, što je jedan od biljeka za dokazivanje tumora kože (Afaq i sur., 2003). Istraživanja na stani nim kulturama CaCo-2 ljudskih stanica tumora crijeva pokazuju tako er sniženu aktivnost ODC (Schneider i sur., 2000). Obrada ljudskih keratinocita resveratrolom inhibira UVB posredovanu fosforilaciju i degradaciju IκB, aktivaciju IKK, što navodi na zaklju ak da NFκB ima važnu ulogu u kemporevenciji razvoja raka kože (Adhami i sur., 2003).

Što se ti e tumora jetre, pokazano je na stani noj liniji H22 da u ovisnosti o dozi i vremenu primjene resveratrol inhibira rast hepatom induciraju i apoptozu (Sun i sur., 2002), djeluju i ak i pri nižim koncentracijama na hepatom štakora Fao i ljudske hepatoblastoma HepG2 stani ne linije (Delmas i sur., 2000). Pretpostavlja se da resveratrol utje u i na transkripciju CYP1A1 (citokrom P450) preko AHR (aril hidrokarbonskog receptora) djeluje kemopreventivno (Ciolino i Yeh i sur., 2001). Vatikanol C, tetramer resveratrola tako er inducira apoptozu u tri razli ite stani ne linije ljudskih tumorskih stanica crijeva (Ito i sur., 2002). Inhibiciju stani ne proliferacije, akumulaciju stanica u S fazi, te smanjenje ciklina D1, B1 i cdk4 može izazvati i analog resveratrola piceatanol na Caco-2 stani noj liniji (Wolter i sur., 2001, 2002).

U istraživanjima na F344 štakorima sa AOM induciranim tumorima crijeva, je pokazano da resveratrol inhibira zna ajno broj ACF formiranja i njihovu veli inu (Tessitore i sur., 2000). U Min miševa koji imaju geneti ku predispoziciju razvoja intestinalnih tumora, ako im se u vodi za pi e ponudi resveratrol, ak za 70% smanji se pojavnost tumora (Schneider i sur., 2001).

Hecht i sur. su u istraživanju sa A/J miševima induciranim BaP (benzo[a]piren) i NNK tumorima plu a provjeravali utjecaj butiliranog hidroksianizola (BHA), mioinozitola, kurkumina, eskuletina, resveratrola i likopenom oboga enog olerecina raj ice (LTO). Pokazali su da je BHA najbolji inhibitor razvoja tumora, dok je resveratrol neu inkovit (Hecht i sur., 1999). Nešto novija istraživanja sa Lewis miševima nositeljima tumor plu a (LLC) pokazuju inhibiciju metastaziranja oralnim uzimanjem resveratrol 3-O-glukozida i 2,3,4',5-tetrahidroksistilbene-2-O-glukozida, i inhibiciju angiogeneze ljudskih endotelnih stanica umbilikalne vene (HUVECs), bez promjene broja CD8+ i NK1.1+T stanica, što navodi na zaklju ak da se ne radi o aktivaciji prirodno ubila kih stanica ili citotoksi nih T limfocita, nego o inhibiciji DNA sinteze u LLC stanicama i angiogeneze HUVECSa (Kimura i Okuda i sur., 2000, 2001). Nekolicina istraživanja su potvrdila i antiproliferativna i kemopreventivna svojstva resveratrola na leukemije. Pored reduciranja vijabilnosti i smanjene

DNA sinteze u stani noj kulturi ljudskih promijelocitnih stanica leukemije HL-60 (Surh i sur., 1999), zabilježeno je i opadanje ekspresije antiapoptotičnog Bcl-2, te utjecaj kaspaza i CD-95-CD-95 ligand puta (Clement i sur., 1998). Resveratrol inducira smrt stanice aktiviraju i unutarstani ne kaspaze u tumorskim stanicama, no nema utjecaja prvih 72 sata na normalne stanice periferne krvi ljudi, što je povezano sa odsutstvom neobične ekspresije CD95 ili CD95L. Koriste i stani ne linije leukemija dobivenih od pacijenata sa pro-B, pre-B i T stani nim ALL pokazano je da resveratrol i u stanicama otpornim na CD-95 signalne puteve inducira apoptozu bez značajne citotoksičnosti za normalne stanice periferne krvi (Dorrie i sur., 2001). Pretpostavlja se da mehanizam ide preko unutarstani nog mitohondrijima inducirano gubitak stabilnosti mitohondrijske membrane. Zabilježena je i protutumorska aktivnost piceatanola na stanicama BJAB (stanice slične Burkittovom limfomu). On je čak bolje djelovao od resveratrola u *ex vivo* pokusu s limfoblastima 21 pacijenta koji pate od ALL (Wieder i sur., 2001; Potter i sur., 2002). Resveratrol također interagira sa stani nim linijama tumora štitnjače. Dvije linije papilarnih karcinoma, i dvije linije folikularnih karcinoma obrađenih resveratrolom pokazuju aktivaciju i translokaciju u jezgri putem mitogenom aktiviranih (MAPK) kinaza i kinaza reguliranih izvanstani nim signalima (ERK-1 i -2), te DNA fragmentaciju i induciranu apoptozu u oba tipa karcinoma (Shih i sur., 2002).

Propolis i polifenolne sastavnice propolisa

Prema USA odjelu za agrikulturu propolis je smola koju su pčele sakupile s različitih biljaka. Može varirati od svijetlo žute do tamno smeđe, ovisno o izvoru i starosti. Točan sastav propolisa varira ovisno o izvoru, no prosječno sadrži 50% smola i biljnih balzama koji čine polifenolnu frakciju propolisa, 30% voska, 10% esencijalnih i aromatičnih ulja, 5% polena te 5% ostalih sastojaka i organskih otpadaka (Manti i sur., 1983).

Najveći postotak čine flavonoidni pigmenti, identični onima koji se nalaze u biljkama od kojih potječe propolis, a pretpostavlja se da neki mogu biti i modificirani enzimima pčele (Ghisalberti i sur. 1979).

Frenkel i sur. prvi su radili s flavonoidnom sastavnicom propolisa fenetil esterom kafeinske kiseline (CAPE) i zapazili inhibitorno djelovanje na 12-*o*-tetradekanoilforbol-13-acetat-om inducirani edem uha, te inhibiciju ornitin dekarboksilaze (ODC) (Frenkel i sur., 1993).

Rao i sur. (1993) su prvi istraživali djelovanje propolisa i njegovih sastavnica na razvoj tumora crijeva na azoksimetanom induciranim F344 štakorima. Zabilježen je srednji inhibitorni učinak na ODC, a značajan učinak na crijevnu protein-tirozin kinaznu aktivnost

(TPK), te na formiranje formiranje aberantnih kripti debelog crijeva djelovanjem azoksimetana u štakora F344 (Rao i sur., 1993)

Isti tim znanstvenika provjeravao je u inke feniletil-3-metilkafeata (PEMC) na fosfatidil-inozitol specifi nu fosfolipazu C, fosfolipazu A₂, lipooksigenazu (LOX) i ciklooksigenazu (COX) u sluznici crijeva. Nisu zabilježili toksi no djelovanje na jetru, želudac, probavni sustav ili pluća, ali je značajno smanjena pojava invazivnih, neinvazivnih tumora i ukupnog broja adenokarcinoma crijeva, te je smanjena veličina tumora. Smanjena je također i aktivnost fosfolipaze C specifične za fosfatidil-inozitol (PI-PLC) i proizvoda LOX puta iz arahidonske kiseline, ali ne i fosfolipaze A (Rao i sur., 1995).

Do danas je rađeno još mnogo istraživanja s različitim komponentama propolisa te se pokazalo da većina njih između ostalog, pokazuje protutumorska djelovanja (Matsuno i sur., Oršoli i sur. 2007). Najnovija istraživanja rađena su s vodenom otopinom propolisa (engl. water soluble derivatives of propolis; WSDP) fenetil kafeatom (engl. caffeic acid polyphenol esters; CAPE), kafeinskom kiselinom (engl. caffeic acid; CA) i alkoholnom otopinom propolisa (engl. ethanolic ekstrakt of propolis; EEP). WSDP pokazuje nespecifično imunostimulatorno i antimetastatično djelovanje (Oršoli i sur., 2003). Miševi obrađeni s WSDP ili CA prije inokulacije tumora pokazuju produženi životni vijek u usporedbi s neobrađenom skupinom miševa (Oršoli i sur. 2005b). Lokalnom obradom prije inokulacije inhibiran je rast tumora u prisutnosti CA ili CAPE za 29,30% a preživljenje obrađenih životinja je povećano za 51,74% dok je WSDP bio manje djelotvoran. Općenito je formiranje tumora bilo odgođeno što je vjerojatno i glavni razlog povećanog preživljenja (Oršoli i sur., 2005a). CA i CAPE su ovisni o dozi i njihovo protutumorsko djelovanje je jače što je primjenjena doza veća, dok WSDP ne pokazuje takav učinak, te uopće ne utječe na rast tumorskih stanica u *in vitro* istraživanjima. Pokazano je da veće doze CA ili CAPE interagiraju s procesima u stanicama kao što su aktivnost enzima ili razina glutaciona (Oršoli i sur., 2004a, b). Što je najvažnije, u ovim i sličnim istraživanjima dokazana je citotoksičnost na tumorske stanice, dok analogne normalne stanice nisu osjetljive na djelovanje CAPE (Grunberger i sur., 1988). Modificiraju i redoks stanje stanica, CAPE inhibira nekoliko kritičnih signalnih puteva u regulaciji rasta, poput receptora tirozin kinaza i blokira vezanje transkripcijskih regulatora poput NFκB na DNA (Natarajan i sur., 1996). Još se ne zna kojim točno mehanizmom dolazi do ovih zapaženih promjena, no smatra se da je to ključno za njegovo protutumorsko djelovanje.

WSDP u inak posredovan je aktivacijom makrofaga i povećanjem njihove litičke sposobnosti (Oršoli i sur., 2003, 2004b 2008c). Propolisom aktivirani makrofagi pokazuju

izravni (direktni) protutumorski u inak i pojačavaju fagocitozu tumorskih stanica, te indirektni u inak temeljen na topljivim imbenicima koji utječu na metabolizam tumorskih stanica. Potrebna su detaljnija istraživanja na ovom području da bi se razjasnilo ovo indirektno djelovanje makrofaga pod utjecajem polifenolnih sastavnica propolisa (Oršoli i Baši, 2006). *In vitro* istraživanja s papilarnim urotelialnim karcinomom (PUC) obratnim s pripravcima propolisa EEP ili WSDP, te polifenolnim sastavnicama (kafeinskom kiselinom, naringeninom, krizinom i kvercetinom) pokazala su smanjenje vijabilnosti stanica u ovisnosti o dozi, ali i ovisno o vremenu djelovanja do 24 sata; duže izlaganje nije izazvalo dodatni u inak na smrtnost stanica. Navedeni pripravci nisu utjecali na normalne stanice. Najbolje je djelovao EEP i kvercetin (Baši i sur., 2008). Pretpostavlja se da je u inak EEP rezultat sinergisti kog djelovanja različitih polifenola prisutnih u njemu u većoj koncentraciji nego u WSDP (Oršoli i sur., 2006, 2007). Poznato je da flavonoidne komponente propolisa specifično inhibiraju tirozin kinaze i moduliraju signalni put lipidnih kinaza. Inhibiranje tirozin kinaza, fosfatidilinozitol (PI)- i fosfatidilinozitol-4,5- difosfat (PIP₂)-kinaza rezultira u redukciji koncentracije inositol-1,4,5-trifosfata (IP3) što smanjuje otpuštanje Ca²⁺ iz endoplazmatskog retikuluma. To za rezultat ima smanjenje c-myc i Ki-ras onkogeni i pretpostavlja se da je dio antiproliferativnog i proapoptotičkog djelovanja kvercetina (Csokay i sur., 1997; Williams i sur., 2004).

Kemoterapija i djelovanje lijekova na tumore je često umanjena zbog postojanja višestruke otpornosti tumorskih stanica na kemoterapeutike (engl. multi drug resistance MDR) koji iz stanica izbacuju širok spektar hidrofobnih spojeva, uključujući i neke bitne kemoterapeutske spojeve (vinka alkaloidi, antracikline, epipodofilotoksine, taksol, itd) (Germann i sur., 1996). Poznato je da kemoterapeutici induciraju NFκB aktivaciju, koja potiče signale preživljavanja i druge fiziološke procese inhibiraju i apoptozu i potiču rast stanica tumora, te na kraju induciraju i kemorezistenciju tumora. Nadalje, standardna terapija tumora djeluje toksično i na normalne stanice (najugroženije su krvne i krvotvorne stanice) što opet dovodi do problema prilikom njihove kliničke uporabe.

Primjenom propolisa i srodnih flavonoida kao adjuvanasa uz kemoterapeutike, smanjuje se njihova toksičnost za normalne stanice, a povećava za tumorske (Suzuki i sur., 2002; Oršoli i sur., 2008a, 2008c). I to tako što propolis djeluje imunostimulativno na domaćina, a u tumorskim stanicama inhibira aktivnost enzima CYP1A1 i vjerojatno smanjuje djelovanje i funkciju P-glikoproteina (P-gp), jednog od membranskih proteina uključen u MDR (Oršoli i Baši, 2007, Oršoli i sur. 2008c, Leslie i sur., 2001; Sazuka i sur., 1995). Oršoli i sur. su u svom radu iz 2008. godine pokazali da je dodatak propolisa i srodnih flavonoida uz primjenu

irinotekana uzrokovao smanjenje aktivnosti ALT (alanin aminotransferaze) i AST (aspartat aminotransferaze) koji su bili povišeni prilikom upotrebe samog lijeka. Time je dokazan zaštitni učinak flavonoida na normalne stanice. Dok korištenje samog propolisa ili flavonoida, nije mijenjao specifičnu aktivnost AST, ALT ili LDH, što navodi na zaključak da ne utječe direktno na gušteraču i jetru. Nadalje u kombinaciji s cisplatinom i doksorubicinom su spriječili pojavu leukopenije, ako su davani u predobradi (Oršoli i sur., 2008a).

Postavljene su i nove pretpostavke i hipoteze o korištenju propolisa i drugih polifenolnih sastavnica kao radioprotektora. Budući da je već dokazano njihovo djelovanje u hvatanju kisikovih radikala (Chen i sur., 2004; Moreno i sur., 2000), te deaktiviranju slobodnih radikala (Lahouel i sur., 1987; Suzuki i sur. 2002) koji su jedan od glavnih posljedica djelovanja zračenja na organizam. Zaštitno djelovanje propolisa i polifenola od ozračenja cijelog tijela još uvijek nije dobro istraženo (Sankaranarayanan i sur., 2005). Idealni radioprotektori trebaju štiti normalne stanice, dok ne bi imali utjecaj ili čak povećali osjetljivost tumorskih. Kako su najdjelotvorniji radioprotektori sulfhidrilne sastavnice poput cisteina i cisteinamina u dozama kojima djeluju preventivno protiv otrovnih (Hall i sur., 2000), u središte pozornosti su došli propolis i polifenoli, prirodni antioksidansi, dostupni, jeftini i što je najvažnije ne otrovni. Radioprotektivno djelovanje nekih imunomodulatornih spojeva poput bakterijskih lipopolisaharida (LPS), muramil dipeptida, *Mycobacterium bovis* soj BCG i glukan je pokazano (Patchen i sur., 1997). Budući da propolis i njegove polifenolne sastavnice povećavaju hematopoetsku aktivnost, to se može iskoristiti i u radioterapiji (Dimov i sur., 1992; Oršoli i Bašić i sur., 2003). *Ginkgo biloba* propolis i njegove polifenolne sastavnice pokazuju antimitotična svojstva ako se primjene nakon ozračenja (Montoro i sur., Oršoli i Bašić i sur. 2005).

Istraživanje radom s WSDP, EEP, CA, naringinom, krizinom i kvercetinom primjenjeni prije ili poslije ozračenja cijelog tijela miševa (engl. whole body irradiation; WBI), pokazuje da oni imaju radioprotektivna svojstva, slična ili čak bolja od sintetskog radioprotektora amnioetil isothiuree (AET). Prva su oštećenja DNA na limfocitima zato što su oni najosjetljiviji. Predobradom miševa gore navedenim polifenolnim sastavnicama dobiveno je smanjenje oksidativnog oštećenja DNA, pri čemu je najučinkovitija bila WSDP (54,46%), koja je pokazala čak bolje djelovanje od AET (48,52%). Obrada propolisom i polifenolnim sastavnicama nakon WBI pokazuje također redukciju oštećenja; WSDP bio je najdjelotvorniji (89,24%) u usporedbi s AET (75,39%). Međutim predobrada miševa s kvercetinom (QU) pokazala je najbolje preživljenje, dok obrada nakon ozračenja nije bila učinkovita. Nije još sasvim razjašnjen mehanizam kojim djeluju sastavnice (Oršoli i sur., 2008b).

U mnogim animalnim modelima i *in vitro* istraživanjima je pokazana je protutumorska aktivnost polifenolnih sastavnica hrane. No još uvijek nije uvjerljivo dokazano takvo djelovanje i u ljudskim epidemiološkim istraživanjima. Jedan od mogućih razloga je raznolikost ljudske populacije, od različitih genetičkih predispozicija do navika, prehrane i različitih okolišnih uvjeta, dok su animalna istraživanja rađena u strogo kontroliranim laboratorijskim uvjetima. Nadalje, dostupnost komponenti je u organizmu (*in vivo*) mnogo manja od primjenjenih koncentracija u *in vitro* istraživanjima. Bit će potrebno još mnogo *in vivo* istraživanja, po mogućnosti i sa ljudima kako bi se mogli donositi zaključci i preporuke za prehranu kojom bi se smanjio rizik od tumora.

Literatura

1. Adhami V, Siddiqui I, Sarfaraz S, Khwaja S, and Mukhtar H. (2009) Efficacious prostate cancer chemopreventive intervention with green tea polyphenols decreases with increasing tumor grade in the autochthonous transgenic TRAMP model *Clinical Cancer Research* 15, 1947.
2. Adhami VM, Afaq F and Ahmad N (2003) Suppression of ultraviolet B exposure-mediated activation of NF- κ B in normal human keratinocytes by resveratrol. *Neoplasia*. January; 5(1): 74–82.
3. Afaq F, Adhami VM and Ahmad N (2003) Prevention of short-term ultraviolet B radiation-mediated damages by resveratrol in SKH-1 hairless mice. *Toxicol Appl Pharmacol*. Jan 1;186(1):28-37.
4. Bašić I, Oršolić N, Mihaljević Ž and Štajcar D. (2008) Direct cytotoxicity of propolis and its polyphenolic compounds on primary culture of human papillary urothelial carcinoma cells, a possible use in therapy, *Scientific Evidence of the Use of Propolis in Ethnomedicine*,: 377-393
5. Basly JP, Marre-Fournier F, Le Bail JC, Habrioux G and Chulia AJ (2000) Estrogenic/antiestrogenic and scavenging properties of (E)- and (Z)-resveratrol. *Life Sci* 66: 769-777,.
6. Bell JR, Donovan JL, Wong R, et al. (2000) (-)-Catechin in human plasma after ingestion of a single serving of reconstituted red wine. *Am J Clin Nutr*;71:103– 8.
7. Bove K, Lincoln DW and Tsan MF (2002) Effect of resveratrol on growth of 4T1 breast cancer cells *in vitro* and *in vivo*. *Biochem Biophys Res Commun* 291: 1001-1005.
8. Camouse MM, Domingo DS, Swain FR, Conrad EP, Matsui MS, Maes D, Declercq L, Cooper KD, Stevens SR, Baron ED. (2009) Topical application of green and white tea extracts provides protection from solar-simulated ultraviolet light in human skin. *Exp Dermatol*. Jun;18(6):522-6.
9. Cano A, Arnao MB, Williamson G, Garcia-Conesa MT. (2002) Superoxide scavenging by polyphenols: effect of conjugation and dimerization. *Redox Rep*;7:379–83.
10. Carbo N, Costelli P, Baccino FM, Lopez-Soriano FJ and Argiles JM (1999) Resveratrol, a natural product present in wine, decreases tumour growth in a rat tumour model. *Biochem Biophys Res Commun* 254: 739-743.
11. Challa A, Rao DR, Reddy BS, (1997) Interactive suppression of aberrant crypt foci induced by azoxymethane in rat colon by phytic acid and green tea; *Carcinogenesis* 18, 2023–2026.
12. Chen CN, Weng M-S, Wu C-L, Lin J-K. (2004) Comparison of radical scavenging activity, cytotoxic effects and apoptosis induction in human melanoma cells by Taiwanese propolis from different sources. *Evid Based Complement Alternat Med*.;1:175–85.
13. Chinery R, Brockman JA, Peeler Mo, Shyr Y, Beauchamp RD and Coffey RJ (1997) Antioxidants enhance the toxicity of chemotherapeutic agents in colorectal cancer: A p53-independent induction of p21^{Waf1/Cip1} via C/EBP β . *Nature Med* 3: 1233–1241
14. Chung FL, Schwartz J, Herzog CR, Yang YM. (2003) Tea and cancer prevention: studies in animals and humans. *J Nutr*;133:3268S–74S.
15. Ciolino HP and Yeh GC (2001) The effects of resveratrol on CYP1A1 expression and aryl hydrocarbon receptor function *in vitro*. *Adv Exp Med Biol* 492: 183-193.
16. Clement MV, Hirpara JL, Chawdhury and Pervaiz S (1998) Chemopreventive agent resveratrol, a natural product derived from grapes, triggers CD95 signaling-dependent apoptosis in human tumor cells. *Blood* 92: 996-1002.
17. Csokay, B., Prajda, N., Weber, G., and Olah, E. (1997) Molecular mechanisms in the antiproliferative action of quercetin *Life Sci.*, 60(24), 2157-63.
18. Cuendet M and Pezzuto JM (2000) The role of cyclooxygenase and lipoxygenase in cancer chemoprevention. *Drug Metabol Drug Interact* 17: 109-157.
19. Day AJ, Bao Y, Morgan MRA, Williamson G. (2000) Conjugation position of quercetin glucuronides and effect on biological activity. *Free Radic Biol Med*;29:1234–43.
20. Day AJ, Mellon FA, Barron D, Sarrazin G, Morgan MRA, Williamson G. (2001) Human metabolism of dietary flavonoids: identification of plasma metabolites of quercetin. *Free Radic Res*;212:941–52.
21. Delmas D, Jannin B, Malki MC and Latruffe N (2000) Inhibitory effect of resveratrol on the proliferation of human and rat hepatic derived cell lines. *Oncol Rep* 7: 847-852,.
22. Dimov V, Ivanovska N, Bankova N, Popov S (1992) Immunomodulatory action of propolis. IV. Prophylactic activity against Gram-negative infections and adjuvant effect of the water-soluble derivative, *Vaccine* , 10, 817 - 823.
23. Donovan JL, Bell JR, Kasim-Karakas S, et al. (1999) Catechin is present as metabolites in human plasma after consumption of red wine. *J Nutr*;129:1662– 8.
24. Dorrie J, Gerauer H, Wachter Y and Zunino SJ (2001) Resveratrol induces extensive apoptosis by depolarizing mitochondrial membranes and activating caspase-9 in acute lymphoblastic leukemia cells. *Cancer Res* 61: 4731-4739.
25. Fontecave M, Lepoivre M, Elleingand E, Gerez C and Guittet O (1998) Resveratrol, a remarkable inhibitor of ribonucleotide reductase. *FEBS Lett* 421: 277-279.
26. Frenkel K, Wei H, Bhimani R, Ye J, Zadunaisky JA, Huang MT, Ferraro T, Conney AH and Grunberger D (1993) Inhibition of tumor promoter-mediated processes in mouse skin and bovine lens by caffeic acid phenethyl ester. *Cancer Research* 53, 1255± 1261.
27. Gehm BD, McAndrews JM, Chien PY and Jameson JL (1997) Resveratrol, a polyphenolic compound found in grapes and wine, is an agonist for the estrogen receptor. *Proc Natl Acad Sci USA* 94: 14138-14143.

28. Germann UA (1996) P-Glycoprotein – a mediator of multidrug resistance in tumour cells. *Eur J Cancer* 32: 927–944
29. Gescher, A. J. & W. P. Steward (2003) Relationship between mechanisms, bioavailability, and preclinical chemopreventive efficacy of resveratrol: a conundrum. *Cancer Epidemiol Biomarkers Prev*, 12, 953–7
30. Goldberg, D. M., J. Yan & G. J. Soleas (2003) Absorption of three wine-related polyphenols in three different matrices by healthy subjects. *Clin Biochem*, 36, 79–87
31. Grunberger D, Banerjee R, Eisinger K, Oltz EH, Efros L, Caldwell M, Estevez V, Nakanishi K, (1988) Preferential cytotoxicity on tumor cells by caffeic acid phenethyl ester isolated from propolis *Experientia*, 44, 230–232
32. Gupta S, Hastak K, Ahmad N, Lewin JS, Mukhtar H (2001) Inhibition of prostate carcinogenesis in TRAMP mice by oral infusion of green tea polyphenols. *Proc Natl Acad Sci USA*;98:10350–5.
33. Hall, E.J. 2000, Radiobiology for the radiologist. Philadelphia, PA, Lippincott, 136.
34. Han SY, Choung SY, Paik IS, Kang HJ, Choi YH, Kim SJ, and Lee MO (2000) Activation of NF-kappaB determines the sensitivity of human colon cancer cells to TNFalpha-induced apoptosis., *Biol. Pharm. Bull.*, 23, 420.
35. Hao, X., Bose, M., Lambert, J.D., Ju, J., Lee, M.-J., Park, S., Husain, A., Wang, S., Sun, Y., Yang, C.S., 2006. Inhibition of intestinal tumorigenesis in ApcMin/+ mice by (–)-epigallocatechin-3-gallate (EGCG), (–)-epicatechin-3-gallate (ECG), and polyphenon E (PPE). Proceeding Abstract for 2006 AACR Annual Meeting. Abstract # 4894
36. Hecht SS, Kenney PM, Wang M, Trushin N, Agarwal S, Rao AV and Upadhyaya P: (1999) Evaluation of butylated hydroxyanisole, myo-inositol, curcumin, esculetin, resveratrol and lycopene as inhibitors of benzo[a]pyrene plus 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis in A/J mice. *Cancer Lett* 137: 123–130.
37. Hirose, M., et al., (2002) Lack of inhibitory effects of green tea catechins in 1,2-dimethylhydrazine-induced rat intestinal carcinogenesis model: comparison of the different formulations, administration routes and doses. *Cancer Lett.*188, 163–170.
38. Hirose, M., et al., Hoshiya, T., Mizoguchi, Y., Nakamura, A., Akagi, K., Shirai, T., (2001) Green tea catechins enhance tumor development in the colon without effects in the lung or thyroid after pretreatment with 1,2-dimethylhydrazine or 2,2-dihydroxy-di-n-propylnitrosamine in male F344 rats. *Cancer Lett.* 168, 23–29.
39. Holt RR, Lazarus SA, Sullards MC, et al. (2002) Procyanidin dimer B2 [epicatechin-(4_β-8)-epicatechin] in human plasma after the consumption of a flavanol-rich cocoa. *Am J Clin Nutr*;76:798–804.
40. Hsieh TC and Wu JM (1999) Differential effects on growth, cell cycle arrest, and induction of apoptosis by resveratrol in human prostate cancer cell lines. *Exp Cell Res* 249: 109–115.
41. Hsieh TC and Wu JM (2000) Grape-derived chemopreventive agent resveratrol decreases prostate-specific antigen (PSA) expression in LNCaP cells by an androgen receptor (AR)-independent mechanism. *Anticancer Res* 20: 225–228.
42. Huang MT, Xie JG, Wang ZY, et al. (1997) Effects of tea, decaffeinated tea, and caffeine on UVBlight-induced complete carcinogenesis in SKH-1 mice: demonstration of caffeine as a biologically important constituent of tea. *Cancer Res*;57:2623–9.
43. Ito T, Akao Y, Tanaka T, Iinuma M, Nozawa Y and Vaticanol C (2002) A novel resveratrol tetramer, inhibits cell growth through induction of apoptosis in colon cancer cell lines. *Biol Pharm Bull* 25: 147–148.
44. Jang M and Pezzuto JM (1999) Cancer chemopreventive activity of resveratrol. *Drugs Exp Clin Res* 25: 65–77.
45. Kampa M, Hatzoglou A, Notas G, Damianaki A, Bakogeorgou E, Gemetzi C, Kouroumalis E, Martin PM and Castanas E (2000) Wine antioxidant polyphenols inhibit the proliferation of human prostate cancer cell lines. *Nutr Cancer* 37: 223–233.
46. Kapadia GJ, Azuine MA, Tokuda H, Takasaki M, Mukainaka T, Konoshima T and Nishino H (2002) Chemopreventive effect of resveratrol, sesamol, sesame oil and sunflower oil in the Epstein-Barr virus early antigen activation assay and the mouse skin two-stage carcinogenesis. *Pharmacol Res* 45: 499–505.
47. Kim M, Hagiwara N, Smith SJ, Yamamoto T, Yamane T, Takahashi T, (1994) Preventive effect of green tea polyphenols colon carcinogenesis. In: Huang, M.-T., Osaswa, T., Ho, C-T., Rosen, R.T. (Eds.), *Food Phytochemicals in Cancer Prevention*, pp. 51–55. ACS Symposium Series 546.
48. Kimura Y and Okuda H (2000) Effects of naturally occurring stilbene glucosides from medicinal plants and wine, on tumour growth and lung metastasis in Lewis lung carcinoma-bearing mice. *J Pharm Pharmacol* 52: 1287–1295.
49. Kimura Y and Okuda H (2001) Resveratrol isolated from *Polygonum cuspidatum* root prevents tumor growth and metastasis to lung and tumor-induced neovascularization in Lewis lung carcinomabearing mice. *J Nutr* 131: 1844–1849.
50. Kopp P (1998) Resveratrol, a phytoestrogen found in red wine. A possible explanation for the conundrum of the 'French paradox'? *Eur J Endocrinol*, 138, 619–20
51. Kuhnle G, Spencer JP, Chowrimootoo G, Schroeter H, Debnam ES, Srai SK, Rice-Evans C, Hahn U (2000) Resveratrol is absorbed in the small intestine as resveratrol glucuronide. *Biochem Biophys Res Commun*, 272, 212–7
52. Kuwajerwala N, Cifuentes E, Gautam S, Menon M, Barrack ER and Reddy GP (2002) Resveratrol induces prostate cancer cell entry into s phase and inhibits DNA synthesis. *Cancer Res* 62: 2488–2492.
53. Lahouel M, Viotte G, Sumereau E, Morin JP, and Fillastre JP (1987) Haematotoxicity of doxorubicin and 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea (CCNU) and of their association in rats. *Drugs Exp Clin Res.*;13(10):593–9.
54. Landau JM, Wang ZY, Yang GY, Ding W, Yang CS. (1998) Inhibition of spontaneous formation of lung tumors and rhabdomyosarcomas in A/J mice by black and green tea. *Carcinogenesis* 19:501–7.
55. Lee JE and Safe S (2001) Involvement of a post-transcriptional mechanism in the inhibition of CYP1A1 expression by resveratrol in breast cancer cells. *Biochem Pharmacol* 62: 1113–1124.

56. Lee MJ, Maliakal P, Chen LS, et al. (2002) Pharmacokinetics of tea catechins after ingestion of green tea and (–)-epigallocatechin-3-gallate by humans: formation of different metabolites and individual variability. *Cancer Epidemiol Biomarkers Prev*;11:1025–32.
57. Leslie EM, Mao Q, Oleschuk CJ, Deeley RG, and Cole SP (2001) Modulation of multidrug resistance protein 1 (MRP1/ABCC1) transport and ATPase activities by interaction with dietary flavonoids. *Mol Pharmacol*. May;59(5):1171-80.
58. Li N, Chen X, Liao J, et al. (2002) Inhibition of 7,12-dimethylbenz[*a*]anthracene (DMBA)-induced oral carcinogenesis in hamsters by tea and curcumin. *Carcinogenesis*;23:1307–13.
59. Liu JD, Chen SH, Lin CL, Tsai SH, Liang YC. (2001) Inhibition of melanoma growth and metastasis by combination with (–)-epigallocatechin-3-gallate and dacarbazine in mice. *J Cell Biochem*;83:631–42.
60. Lu R and Serrero G (1999) Resveratrol, a natural product derived from grape, exhibits antiestrogenic activity and inhibits the growth of human breast cancer cells. *J Cell Physiol* 179: 297-304.
61. Lu YP, Lou YR, Xie JG, et al. (2002) Topical applications of caffeine or (–)-epigallocatechin gallate (EGCG) inhibit carcinogenesis and selectively increase apoptosis in UVB-induced skin tumors in mice. *Proc Natl Acad Sci USA*;99:12455–60.
62. Manach C, Scalbert A, Morand C, Rémésy C, Jimenez L. Polyphenols: food sources and bioavailability. *Am J Clin Nutr* 2004;79:727–47
63. Matsuno T, (1995) A new clerodane diterpenoid isolated from propolis. *Z. Naturforsch.*, 50, 93–97.
64. Mgbonyebi OP, Russo J and Russo IH (1998) Antiproliferative effect of synthetic resveratrol on human breast epithelial cells. *Int J Oncol* 12: 865-869.
65. Middleton EJr, Kandaswami C, and Theoharides T (2000) The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer; *pharmacol rev* 52:673–751,
66. Mimoto J, Kiura K, Matsuo K, et al. (2000) (–)-Epigallocatechin gallate can prevent cisplatin-induced lung tumorigenesis in A/J mice. *Carcinogenesis*;21:915–9.
67. Montoro A, Almonacid M, Serrano J, Saiz M, Barquinero JF, Barrios L, Verdu G, Perez J, and Villaescusa JI (2005), Assessment by cytogenetic analysis of the radioprotection properties of propolis extract *Radiation Protection Dosimetry*;115:461-464.
68. Moreno M, Isla M, Sampietro A and Vattuone M (2000). Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. *J. Ethnopharmacol.*, **71**: 109-114.
69. Murugan RS, Vinothini G, Hara Y, Nagini S. (2009) Black Tea Polyphenols Target Matrix Metalloproteinases, RECK, Proangiogenic Molecules and Histone Deacetylase in a Rat Hepatocarcinogenesis Model. *Anticancer Res*. Jun;29(6):2301-5.
70. Narayanan BA, Narayanan NK, Stoner GD and Bullock BP (2002) Interactive gene expression pattern in prostate cancer cells exposed to phenolic antioxidants. *Life Sci* 70: 1821-1839.
71. Natarajan K, Singh S, Burke TR, Jr Grunberger D, Aggarwal BB (1996). Caffeic acid phenethyl ester is a potent and specific inhibitor of activation of nuclear transcription factor NF-kappa B. *Proc. Natl. Acad. Sci., U.S.A.*, 93, 9090–9095.
72. Oršoli N and Baši I. (2005) Antitumor, hematostimulative and radioprotective action of water - soluble derivative of propolis (WSDP). *Biomed Pharmacother*;59:561-570.
73. Oršoli N and Baši I., Cancer Chemoprevention by Propolis and its Polyphenolic Compounds in Experimental Animals, *RPMP Vol.17 – Phytochemistry & Pharmacology-III*, Date : 7 // 2006.
74. Oršoli N, Baši I, (2003) Immunomodulation by water-soluble derivative of propolis (WSDP) a factor of antitumor reactivity. *J. Ethnopharmacol.*, 84, 265–273.
75. Oršoli N, Bendelja K, Brbot-Šaranovi A, Baši I (2004a) Antitumor, hematostimulative and radioprotective action of water-soluble derivative of propolis (WSDP) *Period. Biol.*, 106, 367–372.
76. Oršoli N, Bevanda M, Bendelja K, Horvat-Knežević A, Benković V, Baši I (2008c) Propolis and related polyphenolic compounds; their relevance to host resistance and interaction with chemotherapy, *Scientific Evidence of the Use of Propolis in Ethnomedicine*,: 223-250.
77. Oršoli N, Horvat-Knežević A, Benković V, Baši I, (2008a) Benefits of use of propolis and related flavonoids against the toxicity of chemotherapeutic agents *Scientific Evidence of the Use of Propolis in Ethnomedicine*,: 195-222.
78. Oršoli N, Knežević AH, Sver L, Terzić S, Basic I. (2004b) Immunomodulatory and antimetastatic action of propolis and related polyphenolic compounds. *J Ethnopharmacol* 2004; **94**: 307-315.
79. Oršoli N, Terzić S, Mihaljević Ž, Šver L, Baši I. (2005a) Effects of Local Administration of Propolis and Its Polyphenolic Compounds on Tumor Formation and Growth *Biol. Pharm. Bull.* 28(10) 1928–1933
80. Oršoli N, Terzić S, Šver L, Baši I, (2005b) Honey-bee products in prevention and/or therapy of murine transplantable tumours *J. Sci. Food Agric.*, 85, 363–370.
81. Oršoli N., Benković V., Horvat-Knežević A., Baši I., Scientific base for using propolis and its polyphenolic/ flavonoid compounds as an adjunct to radiation therapy *Scientific Evidence of the Use of Propolis in Ethnomedicine*, 2008b: 251-310.
82. Patchen ML Bump EA, and Malaker K (Ed.), (1997) *Radioprotectors: Chemical, Biological, and Clinical Perspectives*, CRC Press, New York, 213.
83. Potter GA, Patterson LH, Wanogho E, Perry PJ, Butler PC, Ijaz T, Ruparelia KC, Lamb JH, Farmer PB, Stanley LA and Burke MD (2002) The cancer preventive agent resveratrol is converted to the anti-cancer agent piceatannol by the cytochrome P450 enzyme CYP1B1. *Br J Cancer* 86: 774-778.
84. Rao CV, Desai D, Rivenson A, Simi B, Amin S and Reddy BS (1995) Chemoprevention of colon carcinogenesis by phenylethyl-3-methylcatechol. *Cancer Research* 55, 2310±2315.

85. Rao CV, Desai D, Simi B, Kulkarni N, Amin S and Reddy BS (1993) Inhibitory effect of caffeic acid esters on azoxymethane-induced biochemical changes and aberrant crypt foci formation in rat colon. *Cancer Research* 53, 4182-4188.
86. Rios LY, Gonthier MP, Remesy C, et al. (2003) Chocolate intake increases urinary excretion of polyphenol-derived phenolic acids in healthy human subjects. *Am J Clin Nutr*;77:912– 8.
87. Roy P, Nigam N, Singh M, George J, Srivastava S, Naqvi H, Shukla Y. (2009) Tea polyphenols inhibit cyclooxygenase-2 expression and block activation of nuclear factor-kappa B and Akt in diethylnitrosoamine induced lung tumors in Swiss mice. *Invest New Drugs*.
88. Sankaranarayanan K, and Wassom JS (2005) Ionizing radiation and genetic risks XIV. Potential research directions in the post-genome era based on knowledge of repair of radiation-induced DNA double-strand breaks in mammalian somatic cells and the origin of deletions associated with human genomic disorders. *Mutat Res*.;578(1-2):333-70. Review.
89. Sazuka M, Murakami S, Isemura M, Satoh K, Nukiwa T. (1995) Inhibitory effects of green tea infusion on in vitro invasion and in vivo metastasis of mouse lung carcinoma cells. *Cancer Lett*;98:27–31.
90. Scalbert A, Williamson G. (2000) Dietary intake and bioavailability of polyphenols. *J Nutr*;130:2073S– 85S
91. Schneider Y, Duranton B, Gosse F, Schleiffer R, Seiler N and Raul F: (2001) Resveratrol inhibits intestinal tumorigenesis and modulates host-defense-related gene expression in an animal model of human familial adenomatous polyposis. *Nutr Cancer* 39: 102-107.
92. Schneider Y, Vincent F, Duranton B, Badolo L, Gosse F, Bergmann C, Seiler N and Raul F (2000) Anti-proliferative effect of resveratrol, a natural component of grapes and wine, on human colonic cancer cells. *Cancer Lett* 158: 85-91.
93. Serrero G and Lu R (2001) Effect of resveratrol on the expression of autocrine growth modulators in human breast cancer cells. *Antioxid Redox Signal* 3: 969-979.
94. Setchell KD, Brown NM, Desai P, et al. (2001) Bioavailability of pure isoflavones in healthy humans and analysis of commercial soy isoflavone supplements. *J Nutr*;131:1362S–75S.
95. Setchell KD, Brown NM, Lydeking-Olsen E. (2002) The clinical importance of the metabolite equol: a clue to the effectiveness of soy and its isoflavones. *J Nutr*;132:3577– 84.
96. Setchell KD, Cassidy A. (1999) Dietary isoflavones: biological effects and relevance to human health. *J Nutr*;129:758S– 67S.
97. She QB, Bode AM, Ma WY, Chen NY and Dong Z (2001) Resveratrol-induced activation of p53 and apoptosis is mediated by extracellular-signal-regulated protein kinases and p38 kinase. *Cancer Res* 61: 1604-1610.
98. Shih A, Davis FB, Lin HY and Davis PJ (2002) Resveratrol induces apoptosis in thyroid cancer cell lines via a MAPK- and p53- dependent mechanism. *J Clin Endocrinol Metab* 87: 1223-1232.
99. Signorelli, P. & R. Ghidoni (2005) Resveratrol as an anticancer nutrient: molecular basis, open questions and promises. *J Nutr Biochem*, 16, 449-66
100. Soleas GJ, Grass L, Joseph PD, Goldberg DM and Diamandis EP (2002) A comparison of the anticarcinogenic properties of four red wine polyphenols. *Clin Biochem* 35: 119-124.
101. Strack D. 1997. Phenolic metabolism. In *Plant Biochemistry*, ed. PM Dey, JB Harbourne, pp. 387–416.
102. Sun ZJ, Pan CE, Liu HS and Wang GJ (2002) Anti-hepatoma activity of resveratrol *in vitro*. *World J Gastroenterol* 8: 79-81.
103. Surh YJ, Hurh YJ, Kang JY, Lee E, Kong G and Lee SJ (1999) Resveratrol, an antioxidant present in red wine, induces apoptosis in human promyelocytic leukemia (HL-60) cells. *Cancer Lett* 140: 1-10.
104. Suzuki, K., Koike, H., Matsui, H., Ono, Y., Hasumi, M., Nakazato, H., Okugi, H., Sekine, Y., Oki, K., Ito, K., Yamamoto, T., Fukabori, Y., Kurokawa, K., and Yamanaka, H. (2002), Genistein, a soy isoflavone, induces glutathione peroxidase in the human prostate cancer cell lines LNCaP and PC-3. *Int. J. Cancer*, 99, 846.
105. Tessitore L, Davit A, Sarotto and Caderni G (2000) Resveratrol depresses the growth of colorectal aberrant crypt foci by affecting bax and p21(CIP) expression. *Carcinogenesis* 21: 1619-1622.
106. Walle T, Hsieh F, DeLegge MH, Oatis JE Jr. & Walle UK (2004) High absorption but very low bioavailability of oral resveratrol in humans. *Drug Metab Dispos*, 32, 1377-82
107. Wargovich MJ, Jimenez A, McKee K, Steele VE, Velasco M, Woods J, Price R, Gray K, Kelloff GJ (2000). Efficacy of potential chemopreventive agents on rat colon aberrant crypt formation and progression. *Carcinogenesis* 21, 1149–1155.
108. Weisburger JH, et al. (1998). Effect of tea extracts, polyphenols, and epigallocatechin gallate on azoxymethane-induced colon cancer. *Proc.Soc. Exp. Biol. Med.* 217, 104–108.
109. Wieder T, Prokop A, Bagci B, Essmann F, Bernicke D, Schulze-Osthoff K, Dorken B, Schmalz HG, Daniel PT and Henze G (2001) Piceatannol, a hydroxylated analog of the chemopreventive agent resveratrol, is a potent inducer of apoptosis in the lymphoma cell line BJAB and in primary, leukemic lymphoblasts. *Leukemia* 15: 1735-1742.
110. Williams RJ, Spencer JP, and Rice-Evans C (2004) Flavonoids: antioxidants or signalling molecules? *Free Radic. Biol. Med.*,36(7), 838-49. Review.
111. Witschi H, Espiritu I, Yu M, Willits NH. (1998) The effects of phenethyl isothiocyanate, N-acetylcysteine and green tea on tobacco smoke-induced lung tumors in strain A/J mice. *Carcinogenesis*;19: 1789 –94.
112. Wolter F, Akoglu B, Clausnitzer A and Stein J (2001) Downregulation of the cyclin D1/Cdk4 complex occurs during resveratrol-induced cell cycle arrest in colon cancer cell lines. *J Nutr* 131: 2197-2203.
113. Wolter F, Clausnitzer A, Akoglu B and Stein J (2002) Piceatannol, a natural analog of resveratrol, inhibits progression through the S phase of the cell cycle in colorectal cancer cell lines. *J Nutr* 132: 298-302.
114. Xu Y, Ho CT, Amin SG, Han C & Chung FL (1992) Inhibition of tobacco-specific nitrosamine-induced lung tumorigenesis in A/J mice by green tea and its major polyphenol as antioxidants. *Cancer Res.* 52: 3875–3879.

115. Xu Y, Ho CT, Amin SG, Han C, Chung FL. (1992) Inhibition of tobaccospecific nitrosamine-induced lung tumorigenesis in A/J mice by green tea and its major polyphenol as antioxidants. *Cancer Res*; 52:3875–9.
116. Xu, M., Bailey, A.C., Hernaez, J.f., Taoka, C.R., Schut, H.A.J., Dashwood, R. H., (1996) Protection by green tea, black tea, and indole-3-carbinol against 2-amino-3-methylimidazo[4,5-f]quinoline-induced DNA adducts and colonic aberrant crypts in the F344 rat. *Carcinogenesis* 17, 1429–1434.
117. Yang CS, Chen LS, Lee MJ, Balentine D, Kuo MC, Schantz SP (1998) Blood and urine levels of tea catechins after ingestion of different amounts of green tea by human volunteers. *Cancer Epidemiol Biomarkers Prev*;7:351–4.
118. Yang CS, Lambert JD, Ju J, Lu G, Sang S (2007) Tea and cancer prevention: Molecular mechanisms and human relevance *Toxicology and Applied Pharmacology* 224 265–273
119. Yang CS, Liao J, Yang GY, Lu G. (2005). Inhibition of lung tumorigenesis by tea. *Exp. Lung Res.* 31, 135–144.
120. Yang CS, Maliakal P, Meng X (2002) Inhibition of carcinogenesis by tea. *Annu. Rev. Pharmacol. Toxicol.* 42, 25–54.
121. Yang G, Wang ZY, Kim S, et al. (1997) Characterization of early pulmonary hyperproliferation and tumor progression and their inhibition by black tea in a 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis model with A/J mice. *Cancer Res*;57:1889 –94.
122. Zhang Y, Song TT, Cunnick JE, Murphy PA, Hendrich S. (1999) Daidzein and genistein glucuronides in vitro are weakly estrogenic and activate human natural killer cells at nutritionally relevant concentrations. *J Nutr*;129:399–405.

Sažetak

Polifenoli su jedan od najšire rasprostranjenih sekundarnih metabolita u biljnom svijetu i kao takvi su prisutni i u ljudskoj prehrani. To su niskomolekularni kemijski spojevi sastavljeni od tri prstena sa različitim supstituentima. Biljkama služe kao zaštita od herbivora, zaraza, ozljeda a osim toga pokazuju i razna antibakterijska, antiviralna, antioksidativna, protutumorska, protuupalna, antialergijska, antimutagena i imunomodulatorna svojstva i aktivnosti u animalnim organizmima i *in vitro*.

Zbog svih tih svojstava su u zadnja dva desetljeća predmet mnogih istraživanja, sa naglaskom na protutumorsku aktivnost. U ovom radu smo se osvrnuli samo na neke polifenole, njihovu dostupnost u animalnim organizmima, njihovo djelovanje i moguće kliničke primjene. Protutumorska aktivnost je u *in vitro* istraživanjima pokazana za gotovo sve istraživane komponente, dok *in vivo* istraživanja nisu bila tako jednoznačna. Najveća prepreka predstavlja ograničena biološka dostupnost. Zbog toga se pretpostavlja da bi najveći i u inak u organizmu mogli imati u probavnom sustavu, zato što je u izravnom doticaju s oralno primjenjenim polifenolima, što je zapravo i cilj ovakvih istraživanja, pokazati učinkovito inhibiranje rasta i razvoja tumora primjenom prirodnih komponenti hrane.

Summary

Polyphenols are one of the most abundant secondary metabolites in plants and as such are present in the human diet. These are low molecular weight chemical compounds composed of a three ring constitution with different substituents. The plants use them as a protection against herbivore, infection, injury and they show a variety of antibacterial, antiviral, antioxidative, antitumor, anti-allergy, antimutagenesis and immunomodulatory properties and activities in animal organisms and *in vitro*.

Because of all these properties they are in the last two decades the subject of many investigations, with emphasis on the antitumor activity. In this paper we have reflected only some of the polyphenols, their presence in animal organisms, their activity and potential clinical application. Antitumor activity in *in vitro* studies was shown for almost all used components, while *in vivo* investigations didn't show such a uniformity. The biggest obstacle is the limited biological availability. Therefore, it is assumed that the largest effect in the organism may have been done in the alimentary tract, because it is in direct contact with the oral applied polyphenols, which is actually the goal of such research, to demonstrate effective inhibition of growth and development of tumors using natural components of food.