

Generalized Jordan-Wigner transformation and number operators

Dorešić, Miroslav; Meljanac, Stjepan; Mileković, Marijan

Source / Izvornik: **Fizika B**, 1994, 3, 57 - 65

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:654404>

Rights / Prava: [In copyright](#)

Download date / Datum preuzimanja: **2021-12-08**



Repository / Repozitorij:

[Repository of Faculty of Science - University of Zagreb](#)



GENERALIZED JORDAN-WIGNER TRANSFORMATION AND NUMBER
OPERATORS

MIROSLAV DOREŠIĆ, STJEPAN MELJANAC and MARIJAN MILEKOVIĆ*

Rudjer Bošković Institute, Bijenička c.54, 41001 Zagreb, Croatia
**Prirodoslovno-Matematički Fakultet, Department of Theoretical Physics*
Bijenička c.32, 41000 Zagreb, Croatia

Received 18 May 1994

UDC 53.01 : 512

PACS 02.20+b

We define a class of deformed multimode oscillator algebras which possess number operators and can be mapped to multimode Bose algebra by generalized Jordan-Wigner transformation. We construct and discuss the states in the Fock space and the corresponding number operators.

1. Introduction

Recently, much interest has been devoted to the study of quantum groups [1] and to generalizations (deformations) of oscillator algebras [2-6]. Any type of single-mode deformed oscillator can be related (mapped) to a single-mode Bose oscillator [7]. Deformations of multimode oscillators have also been studied [8]. For Pusz-Woronowicz oscillators, covariant under the $SU_q(n)(SU_q(n|m))$ quantum algebra (superalgebra) [2,8] and anyonic-type algebras [9], there exist mappings to multimode Bose algebra. However, deformed generalized quon algebras are known [10] for which such mappings do not exist. All these algebras are associative and the R -matrix approach to them has also been pursued [11]. There exist multimode associative deformed-oscillator algebras that cannot be represented by the R -matrix

approach, but can be naturally mapped to the corresponding Bose algebra. We mention that multimode deformed-oscillator algebras are important for possible physical applications, e.g. in q -deformed field theory [9] and generalized statistics [13].

Our aim is to define and analyze a general class of deformed multimode oscillator-algebras which possess well-defined number operators for each type of oscillator and can be mapped to a multimode Bose algebra. These mappings can be viewed as generalized Jordan-Wigner transformations. We discuss the corresponding algebras. For special values of parameters, one recovers all known such algebras [2,3,5,6,8-10]. Finally, we construct and discuss the states in the Fock space and the corresponding number operators for deformed oscillators.

2. Mappings of the Bose algebra and their Fock-space representation

Let us define the annihilation and creation operators b_i, b_i^+ , ($i \in S$) satisfying the Bose algebra

$$\begin{aligned} [b_i, b_j^+] &= \delta_{ij}, \quad \forall i, j \in S \\ [b_i, b_j] &= 0 \end{aligned} \tag{1}$$

where $S = 1, 2, \dots, n$ or S is a set of sites on a lattice. The number operators N_i satisfy

$$\begin{aligned} [N_i, b_j] &= -b_i \delta_{ij}, \quad \forall i, j \in S \\ [N_i, b_j^+] &= b_i^+ \delta_{ij} \\ N_i &= b_i^+ b_i, \quad \forall i, j \in S. \end{aligned} \tag{2}$$

Now we define generalized Jordan-Wigner transformations of the above Bose algebra, Eqs. (1) and (2), as

$$a_i = b_i e^{\sum_j c_{ij} N_j} \sqrt{\frac{\varphi_i(N_i)}{N_i}} \tag{3}$$

where c_{ij} are complex numbers and $\varphi_i(N_i)$ are arbitrary (complex) functions with the properties $\varphi_i(0) = 0$, $\lim_{\epsilon \rightarrow 0} \frac{\varphi_i(\epsilon)}{\epsilon} = 1$, $|\varphi_i(1)| = 1, \forall i \in S$. We further assume that $|\varphi_i(N)|$ are bijective, monotonically increasing functions or that $\varphi_i(N) = \frac{1-(-)^N}{2}$, implying $a_i^2 = 0$. The mappings (3) generalize the mappings considered in Refs. 8 and 11. It is important to note that the number operators are preserved,

i.e.

$$N_i^{(a)} = N_i^{(b)} \equiv N_i, \quad \forall i \in S. \tag{4}$$

Then it is easy to find the corresponding deformed-oscillator algebra:

$$\begin{aligned} a_i a_j &= e^{c_{ji} - c_{ij}} a_j a_i \quad i \neq j \\ a_i a_j^+ &= e^{c_{ij} + c_{ji}^*} a_j^+ a_i, \quad i \neq j \\ a_i a_i^+ &= |\varphi_i(N_i + 1)| e^{\sum_j (c_{ij} + c_{ij}^*) N_j} e^{(c_{ii} + c_{ii}^*)} \\ a_i^+ a_i &= |\varphi_i(N_i)| e^{\sum_j (c_{ij} + c_{ij}^*) N_j}. \end{aligned} \tag{5}$$

Note that $a_i^2 \neq 0$, unless $b_i \varphi_i(N_i) b_i \varphi_i(N_i) = 0$. For example, if $\varphi_i(N_i) = \frac{1 - (-)^{N_i}}{2}$, then $a_i^2 = 0$, implying the hard-core condition for the i^{th} oscillator. Generally, there are other mappings of Bose algebra (Eqs. (1) and (2)), but, in general, they do not have the number operators $N_i^{(a)}$, and Eq. (4) does not hold for mappings other than those in Eq. (3).

We point out that the complete deformed-oscillator algebra is associative owing to the mapping of Bose algebra. The Fock space for the deformed-oscillator algebra is spanned by powers of the creation operators a_i^+ , $i \in S$, acting on the vacuum $|0 \rangle^{(a)} = |0 \rangle^{(b)} \equiv |0 \rangle$. The states in the Fock space are specified by the eigenvalues of the number operators N_i , namely $|n_1, n_2, \dots, n_i, \dots \rangle^{(a)} = |n_1, n_2, \dots, n_i, \dots \rangle^{(b)}$. (If there exists a number $n_i^{(0)} \in \mathbf{N}$, such that $\varphi_i(n_i^{(0)}) = 0$, then $N_i = 0, 1, \dots, (n_i^{(0)} - 1)$.) States with unit norm are

$$\begin{aligned} |n_1, \dots, n_n \rangle &= \frac{(a_1^+)^{n_1} \dots (a_n^+)^{n_n}}{\sqrt{[\tilde{\varphi}_1(n_1)]! \dots [\tilde{\varphi}_n(n_n)]!}} e^{-\frac{1}{2} \sum_j \theta_{ji} (c_{ij} + c_{ij}^*) n_i n_j} |0, 0, \dots, 0 \rangle \\ [\tilde{\varphi}(n)]! &= \tilde{\varphi}(n) \tilde{\varphi}(n-1) \dots \tilde{\varphi}(1) \\ \tilde{\varphi}_i(n_i) &= |\varphi_i(n_i)| e^{(c_{ii} + c_{ii}^*) n_i}, \end{aligned} \tag{6}$$

where θ_{ij} is the step function. (For anyons in $(2 + 1)$ -dimension space [9], θ is the angle function.)

Furthermore, the matrix elements of the operators a_i, a_i^+ , $i \in S$, are

$$\begin{aligned} \langle \dots (n_i - 1) \dots | a_i | \dots n_i \dots \rangle &= \langle \dots n_i \dots | a_i^+ | \dots (n_i - 1) \dots \rangle^* \\ &= \sqrt{\varphi_i(n_i)} e^{\frac{1}{2} \sum_j (c_{ij} + c_{ij}^*) n_j} \prod_{j \neq i} \delta_{n_j, n'_j}. \end{aligned} \tag{7}$$

We also find for any $k = 0, 1, 2, \dots$ that

$$(a_j^+)^k (a_j)^k = \frac{[\tilde{\varphi}_j(N_j)]!}{[\tilde{\varphi}_j(N_j - k)]!} e^{k \sum_{l \neq j} (c_{jl} + c_{jl}^*) N_l}$$

$$(a_j)^k (a_j^+)^k = \frac{[\tilde{\varphi}_j(N_j + k)]!}{[\tilde{\varphi}_j(N_j)]!} e^{k \sum_{l \neq j} (c_{jl} + c_{jl}^*) N_l}. \quad (8)$$

The norms of arbitrary linear combinations of the states in Eq. (6) in the Fock space, corresponding to deformed-oscillator algebra, are positive by definite owing to the mapping of Bose algebra (Eqs. (1) and (2)). Namely, $|n_1, n_2, \dots, n_i \dots \rangle^{(a)} = |n_1, n_2, \dots, n_i \dots \rangle^{(b)} \equiv |n_1, n_2, \dots, n_i \dots \rangle$.

This class of deformed multimode oscillator algebras comprises multimode Biedenharn-Macfarlane [3], Aric-Coon [6], two- (p, q) parameter [12], Fermi, generalized Green's [13,14], as well as anyonic [9] and Pusz-Woronowicz (with or without the hard-core condition) oscillators covariant under the $SU_q(n)$ ($SU_q(n|m)$) algebra (superalgebra) [8].

Non-isomorphic (non-equivalent) algebras are classified by different matrix elements given by (7), i.e. with the functions $g_i(n_1, n_2, \dots, n_n) = |\varphi_i(n_i)| e^{\sum_j (c_{ij} + c_{ij}^*) n_j}$. It is important to mention that there are mappings of Bose algebra which do not preserve the relation $N_i^{(a)} = N_i^{(b)}$, given by (4). Moreover, there are mappings of Bose algebra for which the number operators $N_i^{(a)}$ do not even exist. Such an example is the exchange algebra presented in Ref. 15.

3. Number operators

Finally, we construct and discuss the number operators $N_i^{(a)}$. Let $\varphi_i(n_i)$ be bijective mappings. Then, using Eq. (3), we have

$$b_i = a_i e^{-\sum_j c_{ij} N_j} \sqrt{\frac{N_i}{\varphi_i(n_i)}}, \quad \forall i \in S$$

$$N_i = b_i^+ b_i. \quad (9)$$

The spectra of $N_i^{(a)}$ and $N_i^{(b)}$ coincide. In this case,

$$a_i^+ a_i = g_i(N_1, N_2, \dots, N_i \dots) = |\tilde{\varphi}_i(N_i)| e^{\sum_{j \neq i} (c_{ij} + c_{ij}^*) N_j}. \quad (10)$$

Let us denote $a_i^+ a_i = x_i$, and remark that x_i commute with any N_j and among themselves. Then, the N_i operators can be written as

$$N_i = N_i(x_1, \dots, x_n) = \sum_{k=0}^{\infty} \sum_{(i_1 \dots i_k)} \frac{1}{k!} \left(\frac{\partial^k N_i}{\partial x_{i_1} \dots \partial x_{i_k}} \right)_{x=0} (x_{i_1} x_{i_2} \dots x_{i_k})$$

$$N_i(0, 0, \dots, 0) = 0, \quad \forall i \in S. \tag{11}$$

The coefficients in the Taylor expansion can be obtained from (10)

$$g_i(N_1, N_2, \dots, N_i \dots) = x_i, \quad \forall i \in S \tag{12}$$

namely, from

$$\left(\frac{\partial^k g_i(N_1, N_2, \dots, N_i \dots)}{\partial x_{j_1} \dots \partial x_{j_k}} \right)_{x=0} = \delta_{k1} \delta_{ji}. \tag{13}$$

These equations give a set of recurrence relations for the coefficients in the Taylor expansion of N_i as a function of the variables x_j , $j \in S$. For example,

$$c_0^{(i)} \equiv (N_i)_{x=0} = 0$$

$$c_j^{(i)} \equiv \left(\frac{\partial N_i}{\partial x_j} \right)_{x=0} = \frac{1}{\left(\frac{\partial g_i}{\partial N_i} \right)_0} \delta_{ij} \tag{14}$$

$$c_{jk}^{(i)} \equiv \frac{1}{2} \left(\frac{\partial^2 N_i}{\partial x_j \partial x_k} \right)_{x=0} = -\frac{1}{2} \left(\frac{\partial^2 g_i}{\partial N_j \partial N_k} \right)_0 c_i^{(i)} c_j^{(j)} c_k^{(k)}.$$

Specially,

$$c_j^{(i)} = \frac{1}{\left(\frac{d\varphi_i}{dN_i} \right)_0} \delta_{ij}$$

$$c_{ij}^{(i)} = -\frac{1}{2} (c_{ij} + c_{ji}^*) c_i^{(i)} c_j^{(j)} \tag{15}$$

$$c_{jk}^{(i)} = 0, \quad j \neq i, \quad k \neq i$$

where $g_i(N_1, N_2, \dots, N_i \dots)$ is given by Eq. (10).

We also mention the following result. If there exists a continuous mapping defined by (3) from Bose oscillators to deformed oscillators, then the number operators exist and can be written in the form

$$N_i = x_i [c_i^{(i)} + \sum_j c_j^{(i)} x_j + \dots] \tag{16}$$

where $x_j = a_j^+ a_j$.

For example, let us consider the n -mode Pusz-Woronowicz oscillator algebra [2] (of Bose type, covariant under the $SU_q(n)$ quantum algebra), $q \in \mathbf{R}$:

$$\begin{aligned} a_i a_j &= q^{sgn(j-i)} a_j a_i \\ a_i a_j^+ &= q a_j^+ a_i, \quad i \neq j \\ a_i a_i^+ &= 1 + (q^2 - 1) \sum_j \theta_{ij} a_j^+ a_j + q^2 a_i^+ a_i. \end{aligned} \tag{17}$$

There exist mappings to the Bose algebra (1) and (2) with $c_{ij} = \theta_{ij} \ln q$, $\varphi(N) = \frac{q^{2N} - 1}{q^2 - 1}$ (see Table 1). Hence, we can calculate the coefficients in the expansion (16). Using Eq. (15), we obtain

$$\begin{aligned} c_j^{(i)} &= \frac{q^2 - 1}{\ln q^2} \delta_{ij} \\ c_{ij}^{(i)} &= -\frac{1}{2} \frac{(q^2 - 1)^2}{\ln q^2} (\delta_{ij} + \theta_{ij}). \end{aligned} \tag{18}$$

When $q^2 \rightarrow 1$, then $c_i^{(i)} \rightarrow 1$, whereas $c_{ij}^{(i)} \rightarrow 0$, etc., and $N_i \rightarrow a_i^+ a_i$. However, when $q^2 \rightarrow 0$, all coefficients diverge and the expansion (16) is not valid. Note that when $q = 0$, the mapping (9) becomes singular, but one can still define the corresponding number operators. For $q = 0$, the Pusz-Woronowicz algebra (17) reduces to

$$\begin{aligned} a_i a_j &= 0, \quad i < j \\ a_i a_j^+ &= 0, \quad i \neq j \\ a_i a_i^+ &= 1 - \sum_j \theta_{ij} a_j^+ a_j. \end{aligned} \tag{19}$$

Therefore, we also present the number operators in another form [13] that holds for an arbitrary algebra having number operators, and holds even for $q = 0$:

$$N_i = a_i^+ a_i + \sum_{k=1}^{\infty} \sum_{\pi \in S_k} \sum_{j_1 \dots j_k} d_{\pi(j_1 \dots j_k), j_1 \dots j_k} a_{\pi(j_k)}^+ \dots a_{\pi(j_1)}^+ a_i a_{j_1} \dots a_{j_k} \tag{20}$$

where S_k denotes the permutation group and $d_{\pi(j_1 \dots j_k), j_1 \dots j_k}$ are the coefficients of the expansion.

For the Pusz-Woronowicz algebra, the number operators in the above form (see also Ref. 8) are given by

$$N_i = \sum_{k=0}^{\infty} \sum_{j=0}^k c_{k,j} \sum_{p_1 \dots p_j} \theta_{i p_1} \dots \theta_{i p_j} a_{p_j}^+ \dots a_{p_1}^+ (a_i^+)^{k-j+1} (a_i)^{k-j+1} a_{p_1} \dots a_{p_j} \quad (21)$$

with the conditions $c_{0,0} = 1$, $c_{0,-j} = 0$, $j > 0$, and with the recurrence relations

$$c_{k+1, k+1-j} = \frac{(1-q^2)(1-q^{2j})}{1-q^{2(j+1)}} c_{k, k+1-j} + q^{2j}(1-q^2)c_{k, k-j}. \quad (22)$$

Starting with $c_{0,0} = 1$, we find that

$$c_{k,1} = c_{k,k} = (1-q^2)^k$$

$$c_{k,0} = \frac{(1-q^2)^{k+1}}{1-q^{2(k+1)}}. \quad (23)$$

In the limit $q^2 \rightarrow 1$, the number operators are $N_i = a_i^+ a_i$. In the limit $q = 0$, all coefficients are $c_{k,j} = 1$, $\forall k, j$. This result is similar to that found by Greenberg in Ref.10 for quons with $q = 0$. Finally, for $\varphi_i(N_i) = \frac{1-(-)^{N_i}}{2}$, the number operators, $N_i^{(a)} \neq N_i^{(b)}$, are simply

$$N_i^{(a)} = a_i^+ a_i = b_i^+ b_i [1 - \theta(n_i - 1)]. \quad (24)$$

We conclude with a few remarks. Using the general Jordan-Wigner transformation, Eq. (3), we unify and generalize the results for the states in the Fock space for the mappings of multimode Bose algebra presented in Table 1. We point out that this transformation represents the most general class of multimode deformed oscillator algebras preserving number operators. We show that for $q \neq 0$ the number operators can be expanded in Taylor series in commuting operators $x_j = a_j^\dagger a_j$, Eq. (11), which is complementary to the results of Jagannathan et al. in Ref. 8. However, this expansion (as well as the expansion used in Ref. 8) diverges for $q = 0$. Hence, we give another form of the number operators, Eq. (20), which is valid for arbitrary q and particularly useful for $q = 0$. This expression, Eq. (20), holds for an arbitrary algebra having number operators.

TABLE 1.
Parameters c_{ij} and $\varphi_i(N_i)$ for the various deformed-oscillator algebras. We use definition $[x]_p = \frac{p^x - 1}{p - 1}$.

type of oscillators	c_{ij}	$\varphi_i(N_i)$
Bose	$i\pi\lambda_{ij}, \lambda_{ij} - \lambda_{ji} \in 2\mathbf{Z}, \forall i, j$	N_i
Fermi	$i\pi\lambda_{ij}, \lambda_{ij} - \lambda_{ji} \in 2\mathbf{Z} + 1, \forall i, j$	$[N_i]_{-1}$
Green's oscillators [13,14]	$i\pi\lambda_{ij}, \lambda_{ij} - \lambda_{ji} \in \mathbf{Z}, \forall i, j$	$[N_i]_{\pm 1}$
Anyonic-type [11]	$i\pi\lambda_{ij}, \lambda_{ij} - \lambda_{ji} \in \mathbf{R}, \forall i, j$	$[N_i]_{\pm 1}$
Anyons [9]	$i\lambda\theta_{ij}, \lambda \in \mathbf{R}, \theta_{ij}$ is angle	$[N_i]_{\pm 1}$
Pusz-Woronowicz (Bose) [2]	$\theta_{ij} \ln q, q \in \mathbf{R}$	$[N_i]_{q^2}$
Pusz-Woronowicz (Fermi) [2]	$\theta_{ij} \ln(-q), q \in \mathbf{R}$	$[N_i]_{-1}$
$SU_q(n m)$ -covariant (Bose) oscillators [8]	$\theta_{ij} \ln q, q \in \mathbf{R}, i \leq n$	$[N_i]_{q^2}$
$SU_q(n m)$ -covariant (Fermi) oscillators [8]	$\theta_{ij} \ln(-q), q \in \mathbf{R}, n + 1 \leq i$	$[N_i]_{-1}$
Biedenharn-Macfarlane [3]	$-\frac{1}{2}\delta_{ij} \ln q_i, q_i \in \mathbf{C}$	$[N_i]_{q_i^2}$
Arik-Coon [6]	$0, q_i \in \mathbf{C}$	$[N_i]_{q_i}$

Acknowledgements

This work was supported by the joint Croatian-American contract NSF JF 999 and the Scientific Fund of the Republic of Croatia.

References

- 1) V. G. Drinfeld, in *Proceedings of the ICM (Berkeley, CA)* p.798, (1986);
M. Jimbo, *Lett. Math. Phys.* **10** (1985) 63;
- 2) W. Pusz and S. L. Woronowicz, *Rep. Math. Phys.* **27** (1989) 231;
- 3) L. C. Biedenharn, *J. Phys. A: Math. Gen.* **22** (1989) L873;
A. J. Macfarlane, *J. Phys. A: Math. Gen.* **22** (1989) L983;
- 4) T. Hayashi, *Comm. Math. Phys.* **127** (1990) 129;
- 5) K. Odaka, T. Kishi and S. Kamefuchi, *J. Phys. A: Math. Gen.* **24** (1991) L591;
S. Chaturvedi and V. Srinivasan, *Phys. Rev. A* **44** (1992) 8020;
A. J. Macfarlane, *J. Math. Phys.* **35** (1994) 1054;
- 6) M. Arik and D. D. Coon, *J. Math. Phys.* **17** (1976) 524;
- 7) S. Meljanac, M. Mileković and S. Pallua, *Unified View of Deformed Single-Mode Oscillator Algebras*, preprint RBI-TH-3/94, PMF-ZTF-3/94, HEP-TH-9404039, (to appear in *Phys. Lett. B*);

- 8) R. Chakrabarti and R. Jagannathan, *J. Phys. A: Math. Gen.* **25** (1992) 6393;
R. Jagannathan, R. Sridhar, R. Vasudevan, S. Chaturvedi, M. Krishnakumari, P. Shanta and V. Srinivasan, *J. Phys. A: Math. Gen.* **25** (1992) 6429 and references therein;
- 9) A. Lerda and S. Sciuto, *Nucl. Phys. B* **401** (1993) 613;
V. Bardek, M. Dorešić and S. Meljanac, *Example of q-deformed field theory*, preprint HEP-TH-9310171 (to appear in *Int. J. Mod. Phys.*);
V. Bardek, M. Dorešić and S. Meljanac, *Anyons as Quons*, (to appear in *Phys. Rev. D*);
- 10) O. W. Greenberg, *Phys. Rev. Lett.* **64** (1990) 705;
Phys. Rev. D **43** (1991) 4111;
- 11) D. Fairlie and C. Zachos, *Phys. Lett.* **256B** (1991) 43;
P. P. Kulish, *Phys. Lett.* **161A** (1991) 50;
A. Kempf, *J. Math. Phys.* **34** (1993) 969;
J. Van der Jeugt, *J. Phys. A: Math. Gen.* **26** (1993) L405;
- 12) R. Chakrabarti and R. Jagannathan, *J. Phys. A: Math. Gen.* **24** (1991) L711;
M. R. Kibler, *Introduction to quantum algebras*, preprint LYCEN 9234;
- 13) S. Meljanac and A. Perica, *Generalized quon statistics*, preprint RBI-TH-10/93, *Number operator in a general quon algebra*, preprint RBI-TH-4/94 (accepted for publication in *J. Phys. A: Math. Gen.*);
- 14) Y. Ohnuki and S. Kamefuchi, *Quantum Field Theory and Parastatistic* (University of Tokio Press, Tokio, Springer, Berlin, 1982);
- 15) L. Brink, T. H. Hansson, S. Konstein and M. A. Vasiliev, *Nucl. Phys.* **401B** (1993) 591;
A. P. Polychronakos, *Phys. Rev. Lett.* **69** (1992) 703.

GENERALIZIRANA JORDAN-WIGNEROVA TRANSFORMACIJA I OPERATORI BROJA ČESTICA

MIROSLAV DOREŠIĆ, STJEPAN MELJANAC i MARIJAN MILEKOVIĆ*

*Institut Ruđer Bošković, Bijenička c.54, 41001 Zagreb,
* Prirodoslovno-Matematički Fakultet, Zavod za teorijsku fiziku
Bijenička c.32, 41000 Zagreb,*

UDK 53.01 :512

PACS 02.20+b

Definirana je klasa deformiranih oscilatornih algebri sa operatorima broja čestica i poopćenim Jordan-Wignerovim preslikavanjem na Bose algebre. Konstruirana su i diskutirana stanja u Fockovom prostoru te pripadajući operatori broja čestica.