

Laser assisted tunneling in a Tonks-Girardeau gas

Lelas, Karlo; Drpić, Nikola; Dubček, Tena; Jukić, Dario; Pezer, Robert; Buljan, Hrvoje

Source / Izvornik: **New Journal of Physics, 2016, 18**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1088/1367-2630/18/9/095002>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:085890>

Rights / Prava: [Attribution 3.0 Unported](#)/[Imenovanje 3.0](#)

Download date / Datum preuzimanja: **2025-04-02**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)





PAPER

Laser assisted tunneling in a Tonks–Girardeau gas

Karlo Lelas¹, Nikola Drpić², Tena Dubček², Dario Jukić³, Robert Pezer⁴ and Hrvoje Buljan²¹ Faculty of Textile Technology, University of Zagreb, Prilaz baruna Filipovića 28a, 10000 Zagreb, Croatia² Department of Physics, University of Zagreb, Bijenička c. 32, 10000 Zagreb, Croatia³ Faculty of Civil Engineering, University of Zagreb, Fra Andrije Kačića-Miošića 26, 10000 Zagreb, Croatia⁴ Faculty of Metallurgy, University of Zagreb, Aleja narodnih heroja 3, HR 44103 Sisak, CroatiaE-mail: hbuljan@phy.hr

Keywords: Tonks–Girardeau gas, laser assisted tunneling, synthetic gauge fields

OPEN ACCESS

RECEIVED
5 June 2016REVISED
18 July 2016ACCEPTED FOR PUBLICATION
2 August 2016PUBLISHED
6 September 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

We investigate the applicability of laser assisted tunneling in a strongly interacting one-dimensional (1D) Bose gas (the Tonks–Girardeau gas) in optical lattices. We find that the stroboscopic dynamics of the Tonks–Girardeau gas in a *continuous* Wannier–Stark-ladder potential, supplemented with laser assisted tunneling, effectively realizes the ground state of 1D hard-core bosons in a *discrete* lattice with nontrivial hopping phases. We compare observables that are affected by the interactions, such as the momentum distribution, natural orbitals and their occupancies, in the time-dependent *continuous* system, to those of the ground state of the *discrete* system. Stroboscopically, we find an excellent agreement, indicating that laser assisted tunneling is a viable technique for realizing novel ground states and phases with hard-core 1D Bose gases.

1. Introduction

The fractional quantum Hall (FQH) state emerges in a system of strongly interacting charged particles in a strong magnetic field and confined in two-dimensions [1, 2]. The richness of this system motivates the quest for novel topological states of matter in other systems [3]. Ultracold atomic gases are an ideal playground for a controlled preparation, manipulation, and detection of quantum many-body states [4]. However, to achieve topological states such as the FQH state in ultracold atomic gases, one must create a synthetic (artificial) magnetic field, wherein atoms behave as charged particles in magnetic fields [5–7].

A variety of methods for the creation of synthetic magnetic fields have been implemented over the years [8–14], including the Coriolis force method in rapidly rotating BECs [8, 9], and methods based on the Berry phase, which plays the role of the Aharon–Bohm phase (see [10] for bulk BECs). In optical lattices, one engineers the amplitude and the phase of the tunneling matrix elements (hopping parameters) [11–14], for example, by shaking the lattice [12] or using laser assisted tunneling [11, 13–16]. This has led to the experimental realization of paradigmatic condensed-matter Hamiltonians such as the Harper–Hofstadter Hamiltonian [13, 14] and the Haldane Hamiltonian [17]. However, most of the efforts regarding synthetic magnetic fields were focused on single particle effects.

Recently, strong interactions were used in the physics of gauge fields, in the first observation of BEC (i.e. the ground state) in the Harper–Hofstadter Hamiltonian [18]. As synthetic magnetic fields in optical lattices are essentially obtained by periodic driving [11–13], an important question in this context is to understand the behavior of periodically driven interacting quantum systems [19–23]. From the eigenstate thermalization hypothesis it follows that driven interacting systems will, after sufficiently long time, heat up to an infinite temperature [24]. However, in some regimes, the system can approach a prethermalized Floquet steady state before heating up [21, 22], implying that the method can be used (in some regimes) with interactions present. Next, laser assisted tunneling was suggested as a scheme to engineer and promote three-body interactions in atomic gases [25]. An interplay of on-site Hubbard interactions and laser assisted tunneling was recently suggested for realizing versatile Hamiltonians in optical lattices [26]. Following the realization of the Harper–Hofstadter Hamiltonian with bosonic atoms [13, 14], two-dimensional strongly correlated lattice bosons in a

strong magnetic field were recently studied [27]. In one-dimensional (1D) fermionic systems, the classification of topological phases was shown to depend on the presence/absence of interactions [28].

In the quest for strongly correlated topological states, the applicability of methods for synthetic magnetic fields should be scrutinized in the presence of interactions. Here we examine the applicability of laser assisted tunneling [11, 13, 14] for a strongly interacting 1D Bose gas (the Tonks–Girardeau gas [29]) in an optical lattice. The Tonks–Girardeau model is exactly solvable via the Fermi–Bose mapping, i.e., by mapping a wave function for noninteracting spinless 1D fermions to that of impenetrable core 1D bosons [29]. The experimental realization of the Tonks–Girardeau gas in atomic waveguides, proposed by Olshanii [30], has been accomplished more than a decade ago [31–33]. Impenetrable core interactions for bosons mimic the Pauli exclusion principle in x -space, thus, the single particle density is identical for the Tonks–Girardeau gas and noninteracting spinless fermions [29]. However, the two systems considerably differ in momentum space [34]. The laser assisted tunneling should work well for noninteracting spinless fermions (on the Fermi side of the Fermi–Bose mapping [29]), but it is not immediately clear how will the interplay of this method and impenetrable core interactions affect the momentum distribution, and the other observables depending on phase coherence.

Here we demonstrate, by numerical calculations, that the stroboscopic dynamics of a Tonks–Girardeau gas in a *continuous* Wannier–Stark-ladder potential, supplemented with periodic driving, which simulates laser assisted tunneling, effectively realizes the ground state of hard-core bosons (HCB) on the *discrete* lattice with nontrivial hopping phases (i.e. complex hopping parameters). We calculate the momentum distribution, natural orbitals and their occupancies for the ground state of HCB on such a discrete lattice, and find excellent agreement between these results, and observables calculated for a series of stroboscopic moments of the Tonks–Girardeau gas in continuous periodically driven Wannier–Stark-ladders.

Before presenting our results, we further discuss the motivation for studying interacting 1D Bose gases in synthetic gauge fields. Quite generally, in addition to being exactly solvable (in some situations) and experimentally accessible, interacting 1D Bose gases present a many-body system with enhanced quantum effects due to the reduced dimensionality. More specifically, in discrete lattices with phase dependent hopping amplitudes, one may explore strongly correlated ground states and excitations with potentially intriguing many-body properties. It should be mentioned that 1D spin-polarized fermions, 1D HCB, and 1D hard-core anyons, are related through the Bose–Fermi and anyon-fermion mapping [35]. Free expansion of 1D hard core anyons has been studied in [36]. In [37], the multi-particle tunneling decay (in 1D) was studied in dependence on interactions and statistics, by addressing it for these three types of 1D particles. Furthermore, laser-assisted tunneling addressed here, plays the key role in a recent proposal for the experimental realization of anyons in 1D optical lattices [38].

2. The Tonks–Girardeau model

We consider a gas of N identical bosons in 1D, which interact via pointlike interactions, described by the Hamiltonian

$$\hat{H} = \sum_{i=1}^N \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_i^2} + V(x_i, t) \right] + g_{1D} \sum_{1 \leq i < j \leq N} \delta(x_i - x_j). \quad (1)$$

Such a system can be realized with ultracold bosonic atoms trapped in effectively 1D atomic waveguides [31–33], where $V(x)$ is the axial trapping potential, and $g_{1D} = 2\hbar^2 a_{3D} [ma_{\perp}^2 (1 - Ca_{3D}/\sqrt{2}a_{\perp})]^{-1}$ is the effective 1D coupling strength; a_{3D} stands for the three-dimensional s -wave scattering length, $a_{\perp} = \sqrt{\hbar/m\omega_{\perp}}$ is the transverse width of the trap, and $C = 1.4603$ [30]. By varying ω_{\perp} , the system can be tuned from the mean field regime up to the strongly interacting Tonks–Girardeau regime [29] with infinitely repulsive contact interaction $g_{1D} \rightarrow \infty$. For the Tonks–Girardeau regime, the interaction term of the Hamiltonian (1) can be replaced by a boundary condition on the many-body wave function [29]:

$$\Psi_B(x_1, x_2, \dots, x_N, t) = 0 \text{ if } x_i = x_j \quad (2)$$

for any $i \neq j$. With this boundary condition, the Hamiltonian becomes

$$\hat{H} = \sum_{i=1}^N \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_i^2} + V(x_i, t) \right]. \quad (3)$$

The boundary condition (2) and the Schrödinger equation for (3) are satisfied by an antisymmetric many-body wave function Ψ_F describing a system of noninteracting spinless fermions in 1D [29]. Because the system is 1D, an exact (static and time-dependent) solution of the Tonks–Girardeau model can be written via the famous Fermi–Bose mapping [29]:

$$\Psi_B(x_1, x_2, \dots, x_N, t) = \prod_{1 \leq i < j \leq N} \text{sgn}(x_i - x_j) \Psi_F(x_1, x_2, \dots, x_N, t). \quad (4)$$

The fermionic wave function Ψ_F can be written in the form of a Slater determinant (or generally as a superposition of such determinants)

$$\Psi_F(x_1, \dots, x_N, t) = \frac{1}{\sqrt{N!}} \det_{m,j=1}^N [\psi_m(x_j, t)], \quad (5)$$

where $\psi_m(x, t)$ denotes N orthonormal single particle wave functions obeying a set of uncoupled single-particle Schrödinger equations:

$$i\hbar \frac{\partial \psi_m}{\partial t} = \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x, t) \right] \psi_m(x, t), \quad m = 1, \dots, N. \quad (6)$$

Equations (4)–(6) prescribe the construction of the many-body wave function describing the Tonks–Girardeau gas in an external potential $V(x, t)$. The mapping is applicable both in the stationary [29] and the time-dependent case [39].

The expectation values of the one-body observables are obtained from the reduced single particle density matrix (RSPDM), defined as

$$\rho_B(x, y, t) = N \int dx_2 \dots dx_N \Psi_B^*(x, x_2, \dots, x_N, t) \Psi_B(y, x_2, \dots, x_N, t). \quad (7)$$

Observables of interest here are the single particle x -density $\rho_B(x, x, t) = \sum_{m=1}^N |\psi_m(x, t)|^2$, and the momentum distribution [34]:

$$n_B(k, t) = \frac{1}{2\pi} \int dx dy e^{ik(x-y)} \rho_B(x, y, t). \quad (8)$$

A concept that is very useful for the understanding of the bosonic many-body systems is that of natural orbitals. The natural orbitals $\Phi_i(x, t)$ are eigenfunctions of the RSPDM

$$\int dx \rho_B(x, y, t) \Phi_i(x, t) = \lambda_i(t) \Phi_i(y, t), \quad i = 1, 2, \dots, \quad (9)$$

where λ_i are the corresponding eigenvalues; the RSPDM is diagonal in the basis of natural orbitals

$$\rho_B(x, y, t) = \sum_{i=1}^{\infty} \lambda_i(t) \Phi_i^*(x, t) \Phi_i(y, t). \quad (10)$$

The natural orbitals can be interpreted as effective single-particle states occupied by the bosons, where λ_i represents the occupancy of the corresponding orbital [40]. The fermionic RSPDM $\rho_F(x, y, t)$ and the momentum distribution $n_F(k)$ are defined by equations (7) and (8) with $\Psi_B \rightarrow \Psi_F$. The single particle density $\rho_B(x, x, t)$ is identical for the Tonks–Girardeau gas and the noninteracting Fermi gas [29]. However, the momentum distributions of the two systems on the two sides of the mapping considerably differ [34]. The momentum distribution and $\rho_B(x, y, t)$ for the continuous Tonks–Girardeau model (equations (2) and (3)) can be efficiently calculated by using the procedure outlined in [41].


3. Laser assisted tunneling in a Tonks–Girardeau gas

Our strategy is as follows: we study the ground state of hard core bosons (HCB) on a discrete lattice, with nontrivial phases of the hopping parameters (i.e. with complex hopping parameters). Then, we examine in detail the quantum dynamics of the Tonks–Girardeau gas in a continuous Wannier–Stark-ladder potential with periodic driving, which corresponds to laser assisted tunneling [11, 13, 14]. Parameters of the periodic drive are tuned to correspond to the phases of the hopping parameters in the discrete system. More specifically, we observe the (stroboscopic) quantum dynamics of the single particle density in x - and in k -space, and compare it with the ground state properties of HCB on the discrete lattice.

In our simulations of the Tonks–Girardeau gas with periodic driving, the gas is initially (at $t = 0$) in the ground state of the optical lattice potential $V_L(x)$, which has $M = 40$ lattice sites, and infinite wall boundary conditions:

$$V_L(x) = \begin{cases} V_L \cos^2(\pi x/D), & \text{if } -20D \leq x \leq 20D, \\ \infty, & \text{otherwise.} \end{cases} \quad (11)$$

Here $V_L = 10E_R$ is the amplitude of the optical lattice, $E_R = \hbar^2/(2m\lambda^2)$ is the recoil energy, $\lambda = 1.064 \mu\text{m}$, and $D = \lambda/2$ is the period of the optical lattice. The lattice is loaded with ^{87}Rb atoms, $m = 1.455 \times 10^{-25} \text{ kg}$. For such a deep lattice, the Tonks–Girardeau gas (described with Hamiltonian (3) together with condition (2)) can be approximated by the model of HCB on a discrete lattice [43, 44]:



$$\hat{H} = -J \sum_{m=1}^M [\hat{b}_{m+1}^\dagger \hat{b}_m + \text{h.c.}], \quad \hat{b}_m^{\dagger 2} = \hat{b}_m^2 = 0, \quad \{\hat{b}_m, \hat{b}_m^\dagger\} = 1 \quad (12)$$

Here, the bosonic creation and annihilation operators at site m are denoted by \hat{b}_m^\dagger and \hat{b}_m respectively, and J is the hopping parameter. The hard core constraint $\hat{b}_m^{\dagger 2} = \hat{b}_m^2 = 0$ precludes multiple occupancy of one lattice site, and the brackets in equation (12) apply only to on-site anticommutation relations, for $m \neq n$ these operators commute as usual for bosons $[\hat{b}_m, \hat{b}_n^\dagger] = 0$. We calculate the effective hopping parameter $J = 0.019E_R$ from the Wannier states of the optical lattice by using the MLGWS code [45]. The discrete lattice of the Hamiltonian (12) is sketched in figure 1(a).


In order to obtain the HCB discrete lattice model with tunable hopping amplitudes and phases, one can employ the laser assisted tunneling method. The scheme [11, 13, 14] utilizes far off-resonant lasers and a single atomic internal state, which minimizes heating by spontaneous emission. An early theoretical proposal related to this scheme was based on coupling of different internal states [15]. The theoretical proposal in [16] was later modified to obtain a homogeneous synthetic magnetic field [13, 14]. In order to simulate the scheme numerically, at $t = 0$ we introduce the tilt potential $V_T(x) = \alpha x/D$, and simultaneously the time- and space-periodic potential $V_R(x, t) = V_R \cos^2[(qx - \omega t)/2]$. The periodic potential with the tilt, $V_L(x) + V_T(x)$, is the continuous Wannier–Stark-ladder potential; for a sufficiently large tilt, tunneling between neighboring lattice sites is suppressed [13, 14]. The periodic drive potential $V_R(x, t)$ simulates two-photon Raman transitions used to restore the tunneling [13, 14], and introduce nontrivial phases in the hopping parameters. Thus, for $t \geq 0$, the Tonks–Girardeau gas evolves in the time-periodic potential

$$V(x, t) = \begin{cases} V_L(x) + V_T(x) + V_R(x, t), & \text{if } -20D \leq x \leq 20D, \\ \infty, & \text{otherwise.} \end{cases} \quad (13)$$

The strength of the tilt and the drive are set by $\alpha = 0.1V_L$ and $V_R = 0.17V_L$, respectively. The frequency of driving is in resonance with the energy offset between neighboring sites of the tilted potential, that is, $\omega = \alpha/\hbar$.

For a deep optical lattice, our continuous model with potential (13) can be approximated by a discrete Hamiltonian with the kinetic (hopping) term, tilt, drive, and on-site interactions. Such a discrete Hamiltonian is a starting point in [46, 47], for deriving a discrete model with complex hopping amplitudes and interactions. More specifically, in the case of resonant driving ($\omega = \alpha/\hbar$), a unitary transformation can cast this discrete Hamiltonian into a rotating frame, such that the kinetic term, together with the tilt and drive terms, become an effective kinetic term with complex hopping amplitudes [46, 47]. The on-site interaction term is not affected by this unitary transformation [46, 47]. The derivation is applicable for any strength of the interaction U/J [46, 47], where U stands for the on-site interaction energy. Even though the derivation in [46, 47] is for 2D lattices, it is applicable for 1D lattices as well; it is also valid in the strongly interacting limit $U/J \rightarrow \infty$ studied here. In other words, the Tonks–Girardeau gas in the continuous potential (13) can be approximated with the Hamiltonian of HCB on a discrete lattice with nontrivial hopping phases:

$$\hat{H} = -K \sum_{m=1}^M [e^{i\phi_m} \hat{b}_{m+1}^\dagger \hat{b}_m + \text{h.c.}], \quad \hat{b}_m^{\dagger 2} = \hat{b}_m^2 = 0, \quad \{\hat{b}_m, \hat{b}_m^\dagger\} = 1, \quad (14)$$



where the hopping phase is given with $\phi_m = qDm$ and K is the effective hopping amplitude. By choosing a different gauge, the phases in equation (14) can be eliminated. Nevertheless, a comparison of the continuous Tonks–Girardeau system in the time-dependent potential (13), with the discrete model (14), provides valuable information on the applicability of laser assisted tunneling in the presence of strong interactions. Moreover, these results have implications for interacting systems where nontrivial phases cannot be eliminated by a gauge transformation, and in systems with gauge dependent observables. We consider a discrete lattice with $M = 40$ sites, corresponding to $M = 40$ lattice sites of the optical lattice potential (11). The discrete lattice of the Hamiltonian (14) is sketched in figure 1(b).

In what follows, we make a particular choice of the phases of the hopping parameters. In experiments, this is set by choosing the angle between the Raman beams [13, 14], and here we set it by choosing $q = \pi/D$ in the time-dependent potential $V_R(x, t)$; thus the hopping phase is $\phi_m = m\pi$. For this choice of the hopping phase, the discrete lattice has alternating hopping matrix elements, $K(-1)^m$, for tunneling from site m to site $m + 1$. We estimate the effective hopping amplitude to be $K = 0.012E_R$. This is obtained by comparing the expansion of the initially localized single particle Gaussian wave packet in the total potential (13), with the expansion in the discrete lattice (14), and adjusting K until the two patterns coincide; this method was adopted from [48].

In order to obtain the ground state properties of the HCB Hamiltonian (14), we use the Jordan–Wigner transformation [42–44]


$$\hat{b}_m^\dagger = \hat{f}_m^\dagger \prod_{\beta=1}^{m-1} e^{-i\pi\hat{f}_\beta^\dagger\hat{f}_\beta}, \quad \hat{b}_m = \prod_{\beta=1}^{m-1} e^{i\pi\hat{f}_\beta^\dagger\hat{f}_\beta} \hat{f}_m, \quad (15)$$

which maps the HCB Hamiltonian (14), to the Hamiltonian for discrete noninteracting spinless fermions:

$$\hat{H} = - \sum_{m=1}^M [K e^{i\phi_m} \hat{f}_{m+1}^\dagger \hat{f}_m + \text{h.c.}], \quad (16)$$

where \hat{f}_m^\dagger and \hat{f}_m are the creation and annihilation operators for spinless fermions. We calculate the ground state momentum distribution $N_B(k)$ of the HCB Hamiltonian (14), and the ground state momentum distribution $N_F(k)$ of the noninteracting spinless fermions Hamiltonian (16), by using the procedure outlined by Rigol and Muramatsu [43, 44].

In figures 2(a)–(f) we show $N_B(k)$ and $N_F(k)$ in dependence on the number of particles N . The figure can be understood by considering the single-particle energy bands of the discrete lattice with $\phi_m = m\pi$, illustrated in figure 1(c). There are two bands, $E_1(k) = -2K|\sin(kD)|$ and $E_2(k) = 2K|\sin(kD)|$, which touch at a 1D Dirac point at $k = 0$. The ground state of N HCB is constructed by using the first N single particle states [43, 44], which on the Fermi side of the mapping fill the states up to the Fermi level. Note that for $N = 20$, the band $E_1(k)$ is filled and $E_2(k)$ empty, while for $N = 40$ they are both full. For $N < M/2 = 20$, the single particle states partially fill the first band $E_1(k)$ which has minima at the edges of the first Brillouine zone at $\pm\pi/2D$; thus both $N_B(k)$ and $N_F(k)$ are centered at these values. The fermionic distribution has the characteristic plateau(s) with Fermi edges, whereas the bosonic distribution has a spike at the maxima; the spike is a consequence of the fact that bosons





tend to occupy the same single particle state of lowest energy. For filling $M/2 < N < M = 40$, the bosonic momentum distribution retains its peak at the edges of the Brillouin zone $\pm\pi/2D$, whereas the fermionic plateau is inverted because the Fermi level is now at the 2nd band $E_2(k)$. In figure 2(f) we show the results for filled bands $N = M = 40$, i.e. both bosons and fermions are in the Mott insulating state with one particle per lattice site and the two momentum distributions overlap, $N_B(k) = N_F(k)$.

Next, we explore the quantum dynamics of the Tonks–Girardeau gas in the *continuous* Wannier–Stark-ladder potential $V_I(x) + V_T(x)$ with periodic driving $V_R(x, t)$ (simulating laser assisted tunneling). The initial state is the ground state of the optical lattice potential $V_I(x)$ (11). Note that we use capital letters ($N_B(k)$) to describe momentum distributions of the discrete model, and lower case ($n_B(k)$) for the momentum distribution of the continuous model. We find that the time dependent momentum distribution $n_B(k, t)$ for Tonks–Girardeau bosons, and $n_F(k, t)$ for free fermions, are *stroboscopically* [46] in excellent agreement with the momentum distributions $N_B(k)$ and $N_F(k)$ of the discrete lattice models (Hamiltonians (14) and (16)). More specifically, at times $t = (4n - 1)T/4$, $n = \{1, 2, 3 \dots\}$, where $T = 2\pi/\omega = 0.497$ ms is the period of the periodic driving, the *continuous* and *discrete* model momentum distributions coincide. The $N_B(k)$ of the discrete model has its domain in the first Brillouin zone $k \in [-\pi/2D, \pi/2D]$, whereas for $n_B(k, t)$ of the continuous model $k \in (-\infty, \infty)$. Apart from the main peaks in the 1st Brillouin zone, the continuous momentum distributions $n_B(k, t)$ have additional (expected) peaks at positions shifted by an integer number of reciprocal lattice vectors.

In figures 3(a) and (b) we show $n_B(k, t)$ and $n_F(k, t)$ for $N = 10$ particles at the first three stroboscopic times $t = \{3T/4, 7T/4, 11T/4\}$; the lines overlap indicating that a Floquet steady state is reached. We see excellent agreement of the continuous momentum distributions $n_B(k, t)$ and $n_F(k, t)$, presented in figures 3(a) and (b), with the momentum distributions $N_B(k)$ and $N_F(k)$ of the discrete model for $N = 10$ particles, figure 2(a).

In figures 3(c) and (d) we show the momentum distributions $n_B(k, t)$ and $n_F(k, t)$ for non-stroboscopic moments for $N = 10$ particles. At $t = 0$, the Tonks–Girardeau gas is in the ground state of the optical lattice (11), which can be approximated with the discrete model (12) sketched in figure 1(a). The discrete model (12) has one energy band $E(k) = -2J \cos(kD)$. In figure 3(c) we show the momentum distributions $n_B(k, t)$ of the initial state at $t = 0$ (black dotted line) and at $t = T$ (green dotted line); we see that the maxima of $n_B(k, t)$ are at $k = 0, \pm 4\pi/2D$, which is consistent with the minima of the band $E(k)$. The same reasoning holds for $n_F(k, t)$ at times $t = \{0, T\}$ in figure 3(d). For completeness, in figure 3(c), we also show $n_B(k, t)$ for non-stroboscopic times $t = \{0.18T, 0.83T\}$ (black and red solid lines respectively); at these times momentum distributions $n_B(k, t)$ have one dominant maximum and several smaller peaks at various k .

In figures 4(a)–(d), we show the momentum distributions $n_B(k, t)$ and $n_F(k, t)$ at the 10th stroboscopic appearance $t = 39T/4 \approx 5$ ms for different numbers of particles, $N = \{20, 21, 30, 40\}$. We find excellent agreement with the momentum distributions $N_B(k)$ and $N_F(k)$ (shown in figures 2(c)–(e)) of the discrete Hamiltonians (14) and (16), respectively. For $N = 40$, the momentum distributions $n_B(k, t)$ and $n_F(k, t)$ do not have sharp peaks, and they slowly decay as $k \rightarrow \infty$; this shape differs from the uniform $N_B(k)$ and $N_F(k)$



corresponding to the Mott state shown in figure 2(f), which is expected for the lattice of finite depth. However, the bosonic and fermionic distributions coincide both in the continuous and the discrete model.

Besides momentum distributions, it is instructive to study the dynamics of natural orbitals $\Phi_n^{\text{TG}}(x)$ defined in equation (9), and their occupancies λ_n^{TG} . In figure 5(a), we show λ_n^{TG} for the continuous model at $t = 39T/4$, for $N = \{10, 20, 30, 40\}$ particles respectively; as expected, bosons partially condense in the first natural orbital $\Phi_1^{\text{TG}}(x)$, and this is reflected in the fact that the peak in their momentum distribution is located at the minima of the bands irrespective of the filling, which is not the case for fermions (see figure 1). For $N = 40$, bosons are in the Mott state with $\lambda_n^{\text{TG}} \approx 1$ for every n . In figure 5(b), we show occupancies λ_n^{HCB} of the natural orbitals $\Phi_n^{\text{HCB}}(x)$ within the discrete model (calculated with the procedure outlined in [43, 44]). We find the same behavior as for the occupancies λ_n^{TG} shown in figure 5(a).

In figure 5(c), we show the real part of the first natural orbital $\Phi_1^{\text{TG}}(x, t)$ for $N = 10$ at $t = 39T/4$; we see the signature of the effective hopping phase $\phi_m = m\pi$, i.e. the real part of the first orbital changes the sign every two sites (the imaginary part follows the same behavior), and higher natural orbitals follow this behavior. For comparison, in figure 5(d) we show the first natural orbital $\Phi_1^{\text{HCB}}(x)$ of the ground state for $N = 10$ HCB on the discrete lattice. We see excellent agreement between the two models.

4. Conclusion and outlook

In conclusion, we investigated the applicability of the laser assisted tunneling in a strongly interacting Tonks–Girardeau gas. We found that the stroboscopic dynamics of the Tonks–Girardeau gas with laser assisted tunneling effectively realizes the ground state of 1D HCB on a discrete lattice with nontrivial hopping phases. Our strategy was to compare the quantum dynamics of the Tonks–Girardeau gas in a continuous Wannier–Stark-ladder potential with periodic driving, which simulated laser assisted tunneling, with the ground state of the discrete model. More specifically, we have compared the momentum distribution, natural orbitals and their occupancies at stroboscopic moments corresponding to the period of the driving potential, and found excellent agreement.

In the outlook, we would like to point out that by comparing continuous and discrete systems (see also [48]), as we have done here, one opens the way for the study of shallow optical lattices with periodic driving, where the tight-binding approximation is not applicable. These continuous systems could potentially be described as discrete models with the phase dependent next to nearest neighbors hopping amplitudes (multi-band models). It is reasonable to expect that such models, with interactions present, possess many intriguing quantum states yet to be explored. Next, we would like to point out that a similar study could be in principle performed for interactions of finite strength, where one could explore modulations of the interaction strength [49, 50] leading to systems with density dependent hopping parameters.

Acknowledgments

This work was supported by the QuantiXLie Center of Excellence.

References

- [1] Tsui D C, Stormer H L and Gossard A C 1982 *Phys. Rev. Lett.* **48** 1559
- [2] Laughlin R B 1983 *Phys. Rev. Lett.* **50** 1395
- [3] Chen X, Gu Z-C, Liu Z-X and Wen X-G 2012 *Science* **338** 1604
- [4] Bloch I, Dalibard J and Zwerger W 2008 *Rev. Mod. Phys.* **80** 885
- [5] Dalibard J, Gerbier F, Juzeliunas G and Öhberg P 2011 *Rev. Mod. Phys.* **83** 1523
- [6] Bloch I, Dalibard J and Nascimbene S 2012 *Nat. Phys.* **8** 267
- [7] Goldman N, Juzeliunas G, Öhberg P and Spielman I B 2014 *Rep. Prog. Phys.* **77** 126401
- [8] Madison K W, Chevy F, Wohlleben W and Dalibard J 2000 *Phys. Rev. Lett.* **84** 806
- [9] Abo-Shaeer J R, Raman C, Vogels J M and Ketterle W 2001 *Science* **292** 476
- [10] Lin Y-J, Compton R L, Jiménez-García K, Porto J V and Spielman I B 2009 *Nature* **462** 628
- [11] Aidelsburger M, Atala M, Nascimbene S, Trotzky S, Chen Y-A and Bloch I 2011 *Phys. Rev. Lett.* **107** 255301
- [12] Struck J, Ölschläger C, Weinberg M, Hauke P, Simonet J, Eckardt A, Lewenstein M, Sengstock K and Windpassinger P 2012 *Phys. Rev. Lett.* **108** 225304
- [13] Miyake H, Siviloglou G A, Kennedy C J, Cody Burton W and Ketterle W 2013 *Phys. Rev. Lett.* **111** 185302
- [14] Aidelsburger M, Atala M, Lohse M, Barreiro J T, Paredes B and Bloch I 2013 *Phys. Rev. Lett.* **111** 185301
- [15] Jaksch D and Zoller P 2003 *New J. Phys.* **5** 56
- [16] Kolovsky A R 2011 *Europhys. Lett.* **93** 003
- [17] Jotzu G, Messer M, Desbuquois R, Lebrat M, Uehlinger T, Greif D and Esslinger T 2014 *Nature* **515** 237
- [18] Kennedy C J, Burton W C, Chung W C and Ketterle W 2015 *Nat. Phys.* **11** 859
- [19] D'Alessio L and Polkovnikov A 2013 *Ann. Phys.* **333** 19
- [20] D'Alessio L and Rigol M 2014 *Phys. Rev. X* **4** 041048
- [21] Bukov M, Gopalakrishnan S, Knap M and Demler E 2015 *Phys. Rev. Lett.* **115** 205301
- [22] Kuwahara T, Mori T and Saito K 2016 *Ann. Phys.* **367** 96
- [23] Eckardt A and Anisimovas E 2015 *New J. Phys.* **17** 093039
- [24] Rigol M, Dunjko V and Olshanii M 2008 *Nature* **452** 854
- [25] Daley A J and Simon J 2014 *Phys. Rev. A* **89** 053619
- [26] Bermudez A and Porras D 2015 *New J. Phys.* **17** 103021
- [27] Natu S S, Mueller E J and Das Sarma S 2016 *Phys. Rev. A* **93** 063610
- [28] Fidkowski L and Kitaev A 2010 *Phys. Rev. B* **81** 134509
- [29] Girardeau M 1960 *J. Math. Phys.* **1** 516
- [30] Olshanii M 1998 *Phys. Rev. Lett.* **81** 938
- [31] Kinoshita T, Wenger T and Weiss D S 2004 *Science* **305** 1125
- [32] Paredes B, Widera A, Murg V, Mandel O, Fölling S, Cirac I, Shlyapnikov G V, Hänsch T W and Bloch I 2004 *Nature* **429** 277
- [33] Kinoshita T, Wenger T and Weiss D S 2006 *Nature* **440** 900
- [34] Lenard A 1964 *J. Math. Phys.* **5** 930
- [34] Schultz T D 1963 *J. Math. Phys.* **4** 666
- [34] Vaidya H G and Tracy C A 1979 *Phys. Rev. Lett.* **42** 3
- [35] Girardeau M D 2006 *Phys. Rev. Lett.* **97** 100402
- [36] del Campo A 2008 *Phys. Rev. A* **78** 045602
- [37] del Campo A 2011 *Phys. Rev. A* **84** 012113
- [38] Keilmann T, Lanzmich S, McCulloch I and Roncaglia M 2011 *Nat. Commun.* **2** 361

- [39] Girardeau M D and Wright E M 2000 *Phys. Rev. Lett.* **84** 5691
- [40] Girardeau M D, Wright E M and Triscari J M 2001 *Phys. Rev. A* **63** 033601
Lapeyre G J, Girardeau M D and Wright E M 2002 *Phys. Rev. A* **66** 023606
- [41] Pezer R and Buljan H 2007 *Phys. Rev. Lett.* **98** 240403
- [42] Jordan P and Wigner E 1928 *Z. Phys.* **47** 631
- [43] Rigol M and Muramatsu A 2004 *Phys. Rev. Lett.* **93** 230404
- [44] Rigol M and Muramatsu A 2005 *Mod. Phys. Lett. B* **19** 861
- [45] Walters R, Cotugno G, Johnson T H, Clark S R and Jaksch D 2013 *Phys. Rev. A* **87** 043613
- [46] Bukov M and Polkovnikov A 2014 *Phys. Rev. A* **90** 043613
- [47] Goldman N, Dalibard J, Aidelsburger M and Cooper N R 2015 *Phys. Rev. A* **91** 033632
- [48] Dubček T, Lelas K, Jukić D, Pezer R, Soljačić M and Buljan H 2015 *New J. Phys.* **17** 125002
- [49] Meinert F, Mark M J, Lauber K, Daley A J and Nägerl H-C 2016 *Phys. Rev. Lett.* **116** 205301
- [50] Rapp A, Deng X and Santos L 2012 *Phys. Rev. Lett.* **109** 203005